or the synergic action of insulin and endogenous cortisone may be a factor in our results (Henderson *et al.*, 1950).

It is stressed that the use of insulin hypoglycaemia should not be looked upon, at all events at present, as a new treatment for rheumatoid arthritis. This is only a report on a pilot series of cases, which require further control and follow-up. Further information is being sought on such problems as potassium excretion, fluid retention, correlation of results with A.C.T.H.-sensitivity, and the value of intermittent treatment.

Summary

Forty cases of active rheumatoid arthritis have been treated five times weekly for either three or four weeks with insulin hypoglycaemia.

Twenty-two of these cases showed marked clinical improvement, nine slight improvement, eight no change, and one deteriorated.

Ten maintained their marked improvement and three moderate improvement six weeks after completion of treatment.

In cases showing definite improvement, this usually began within a week of starting treatment.

All cases increased in weight, some remarkably.

The serum potassium was lowered in 34 out of 36 cases at the height of hypoglycaemia, and had risen again in all cases four and a half hours later, at the time of maximal eosinopenia—in 13 significantly above the control value.

Urinary uric acid excretion was significantly raised in 11 cases during treatment, but there was no increase in the neutral 17-ketosteroids in the eight cases in which these were estimated.

The sedimentation rate was not significantly changed.

Fourteen cases showed an increase and three a decrease of more than 1 g./100 ml. in haemoglobin.

Thirty-four cases showed a significant fall in circulating eosinophils four and a half hours after maximal hypoglycaemia. There was no marked correlation, however, between the degree of fall and the degree of hypoglycaemia or the clinical improvement. There was very little correlation between the eosinophil depression produced by adrenaline or A.C.T.H. and either the clinical response or the degree of eosinopenia following hypoglycaemia.

Eleven cases were treated by E.C.T. with an average of five shocks each. Three showed marked improvement, three slight improvement, and five no change.

No significant fall of eosinophils was found as the result of electric convulsion therapy.

We would like to thank Dr. Tyndall, of Messrs. Organon, Ltd., and the Medical Research Council for donations of A.C.T.H., the Empire Rheumatism Council for financial assistance in laboratory expenses, and Drs. J. B. Bennett, T. Northover, R. Pearson, L. Scott White, and J. Thoresby for their help and suggestions; also the nursing and secretarial staff of the unit.

REFERENCES

- Callow, N. H., *et al.* (1938). *Biochem. J.*, **32**, 1312.
 Cannon, W. B., *et al.* (1924). *Amer. J. Physiol.*, **69**, 46.
 Cantarow, A., and Trumper, M. (1945). *Clinical Biochemistry*, 3rd ed. Saunders, Philadelphia.
 Copeman, W. S. C. (1931). *British Medical Journal*, **2**, 1130.
 Godlowski, Z. Z. (1947). *Edinb. med. J.*, **54**, 239.
 — (1949). *Ann. rheum. Dis.*, **8**, 285.
 Henderson, E., *et al.* (1950). *J. clin. Endocrinol.*, **10**, 800.
 Kersley, G. D., *et al.* (1950). *Lancet*, **1**, 703, 1153.
 King, E. J. (1946). *Micro-analysis in Medical Biochemistry*, 1st ed. Churchill, London.
 Langecker, H. (1928). *Arch. exp. Path. Pharm.*, **134**, 155.
 Miller, R. A., *et al.* (1941). *Proc. Soc. exp. Biol., N.Y.*, **47**, 449.
 Rando'ph, T. G. (1949). *J. Lab. clin. Med.*, **34**, 1696.
 Recant, L., *et al.* (1950). *J. clin. Endocrinol.*, **10**, 187.
 Sargant, W., and Slater, E. (1948). *An Introduction to Physical Methods in Psychiatry*, 2nd ed. Livingstone, Edinburgh.
 Sprague, R. G., *et al.* (1950). *Arch. intern. Med.*, **85**, 199.
 Talbott, J. H. (1943). *Gout*. Oxford Univ. Press, London.

THE EFFECT OF CORTISONE ON THE CARDIOGRAM IN CHRONIC ADRENAL INSUFFICIENCY*

BY

WALTER SOMERVILLE, M.D., M.R.C.P.

Senior Registrar, Cardiographic Department, Middlesex Hospital; formerly Assistant in Medicine, Peter Bent Brigham Hospital, Boston, U.S.A.

The beneficial effects of cortisone (17-hydroxy-11-dehydrocorticosterone) on chronic adrenal insufficiency (Thorn, Bayles, Massell, Forsham, Hill, Smith, and Warren, 1949) suggested that the cardiovascular system, which is often involved in this condition, might take part in the improvement. In a typical and severe case the heart is smaller than normal, the blood pressure lower, and the cardiogram shows non-specific abnormalities. These are, in order of frequency, abnormally flat or inverted T waves, prolonged QT interval, low voltage, prolonged PR or QRS interval, and depressed ST segment. On the other hand, the cardiovascular system, including the cardiogram, may be normal (Somerville and Levine, 1950).

Present Series

This series comprises 11 patients (3 males and 8 females) with proved chronic adrenal insufficiency treated with cortisone under the direction of Dr. George W. Thorn. The ages ranged between 14 and 55 years. All had been treated previously with replacement therapy for periods of 2–28 years (average 9 years). The cardiograms of six patients showed one or more of the abnormalities mentioned above. In five instances the cardiograms were normal. The clinical and metabolic changes following cortisone therapy in these patients have been reported elsewhere (Thorn, Forsham, Bennett, Roche, Reiss, Slessor, Flink, and Somerville, 1949).

Cortisone was given by intramuscular injection in total doses of 163 to 850 mg. over periods of 5 to 18 days. In three instances cortisone pellets were implanted subcutaneously and the dosage was estimated from the predetermined rate of absorption of the pellets. Several of

Table showing changes Occurring in the Cardiogram during Cortisone Therapy

Case No.	Cortisone			Cardiogram					Result
	Dose (mg.)	Period of Admin. (Days)		Low Voltage	Prolonged PR	QT	De-pressed ST	Ab-normal T	
1	800	18	Before After					++	Improved
9	300	27	Before After					+	"
22	163	36	Before After	+	+			++	"
28	533	19	Before After	+		+		++	"
6	290	17	Before After	+		+	+	++	Unchanged
5	530	20	Before After	See note				++	"
12	375	9						++	
15	276	9						++	
49	275	4						++	
65a	850	5						++	
71	200	5						++	

Note.—Abnormalities persisting after cortisone may have been, in part, residue of recent pericarditis.

* From the Medical Clinic of the Peter Bent Brigham Hospital and the Department of Medicine, Harvard Medical School, Boston, U.S.A.

the patients were given additional replacement therapy in small amounts, but from previous experience with such measures in these patients no effect was expected on the cardiovascular system.

Results

The changes noted in the cardiogram during cortisone therapy are summarized in the Table. Four of the six abnormal cardiograms improved and two were unchanged. Improvement was represented by flat or inverted T waves

Changes in blood pressure were not remarkable. In two patients there was a slight increase. No significant alteration was noted in the remainder.

Discussion

The nature of the changes in the cardiovascular system in chronic adrenal insufficiency which give rise to the abnormal cardiogram is unknown. Thorn, Dorrance, and Day (1942) suggested that deficiency of adrenocortical carbohydrate-regulating factors might be responsible, since

After cortisone, the T waves are less inverted or upright, especially in the left ventricular leads (V4-6). In Case 28 the corrected QT interval, and in Case 22 the PR interval, have returned to normal.

becoming upright (see Fig.); shortening of a prolonged PR interval (one case) or QT (corrected) interval (one case). The five normal cardiograms were unchanged.

The duration of the cardiographic improvement after cortisone is uncertain. In one patient the cardiogram had partly regressed towards its original abnormal state two months after cortisone was discontinued. Throughout the observation period and for several weeks before cortisone was given this patient had been receiving small supportive doses of "percorten." In two other patients treated with cortisone pellets inserted subcutaneously the improvement was still present after two to five months.

The heart size, expressed in terms of the transverse diameter determined radiologically, was measured in six patients before and after cortisone therapy. In four the size of the heart increased, in one (normal at the beginning of treatment) it decreased, and in one it was unchanged. In no instance was the size increased beyond normal, and in general the greatest increase was observed in those patients in whom the heart size was smallest when cortisone therapy was instituted.

the cardiogram is usually abnormal in those patients who are unable to fast for 24 hours without a steady fall in blood-sugar levels or severe hypoglycaemic reactions. Cortisone and other 11, 17-oxysteroids are the most active carbohydrate-regulating adrenocortical steroids, and their administration increases the threshold at which hypoglycaemic symptoms occur (Thorn, Koepf, Lewis, and Olsen, 1940). The improvement in four of the six abnormal cardiograms in the present series may be thus explained in part. Yet normal cardiograms may be found in many patients with chronic adrenal insufficiency in whom deficiency of adrenocortical carbohydrate-regulating steroids may be demonstrated by the eosinophil reaction to the A.C.T.H. test described by Thorn, Forsham, Prunty, and Hills (1948). Therefore other factors must exist which lead to cardiographic abnormalities and which are influenced by cortisone, but at present they are unknown.

An increase in heart size following cortisone therapy is in itself unlikely to account for the cardiographic improvement. The incidence of normal and abnormal cardiograms was almost identical whether the heart size was normal, above normal, or below normal in 131 observations on 71 patients with chronic adrenal insufficiency (Somerville and Levine, 1950).

Perera *et al.* (1950) remarked on slight improvement in the cardiogram of a single patient with chronic adrenal insufficiency treated with 400 mg. of cortisone over a period of five days. Commenting on the small increase in blood pressure which they noted, these authors suggested that cortisone may act as a pressor substance in the absence of the adrenal cortex. In the present series cardiographic improvement occurred without changes in blood pressure.

Abnormal cardiograms recorded in acute rheumatic fever have reverted to normal in the course of cortisone therapy. Hench, Slocumb, Barnes, Smith, Polley, and Kendall (1949) have described instances in which prolonged PR intervals, ST segment deviation, and abnormal T waves returned to normal 7-26 days after the beginning

of cortisone therapy. No remarkable effects were noted by these workers on heart size or blood pressure, but in three cases tachycardia was replaced by bradycardia. The dosages of cortisone used were considerably greater than in the present series; on reducing the daily dosage from 200 mg. to 100 mg. the heart rate increased to normal levels. Hench and his associates offered no explanation of the cardiovascular action of cortisone.

Summary

Cortisone was administered to 11 patients (3 males and 8 females) with chronic adrenal insufficiency.

The drug was given by intramuscular injection or pellet implantation in doses of 163 to 850 mg. over 5 to 18 days.

All patients had been treated previously by various forms of replacement therapy.

The cardiograms in six of the patients showed the abnormalities often encountered in chronic adrenal insufficiency—namely, abnormal T waves or prolonged PR or QT intervals. In the remaining five patients the cardiograms were normal.

After cortisone therapy four of the six abnormal cardiograms improved, and two were unchanged. No effect was noted in the five normal cardiograms. The improvement was still present in one patient five months after the treatment was stopped. In two others in whom cortisone pellets had been implanted cardiographic improvement was maintained over a two-months observation period. The cardiogram in one patient had partly regressed two months after cortisone was suspended.

The way in which cortisone influences the cardiogram in chronic adrenal insufficiency is unknown.

This study was aided in part by a grant from the National Heart Institute, United States Public Health Service (Grant No. H-446).

REFERENCES

- Hench, P. S., Slocumb, C. H., Barnes, A. R., Smith, H. L., Polley, H. F., and Kendall, E. C. (1949). *Proc. Mayo Clin.*, **24**, 277.
 Perera, G. A., Pines, K. L., Hamilton, H. B., and Vislocky, K. (1949). *Amer. J. Med.*, **7**, 56.
 Somerville, W., and Levine, H. D. (1950). In press.
 Thorn, G. W., Bayles, T. B., Massell, B. F., Forsham, P. H., Hill, S. R., jun., Smith, S., III, and Warren, J. E. (1949). *New Engl. J. Med.*, **241**, 529.
 ———, Dorrance, S. S., and Day, E. (1942). *Ann. intern. Med.*, **16**, 1053.
 ———, Forsham, P. H., Bennett, L. L., Roche, M., Reiss, R. S., Sessor, A., Flink, E. B., and Somerville, W. (1949). *Trans. Assoc. Amer. Phys.*, **62**, 233.
 ———, Prunty, F. T. G., and Hills, A. G. (1948). *J. Amer. med. Ass.*, **137**, 1005.
 ———, Koepf, G. F., Lewis, R. A., and Olsen, E. F. (1940). *J. clin. Invest.*, **19**, 813.

The Ministry of Health has advised local authorities to make certain changes in the record cards for diphtheria immunization and vaccination. The Ministry suggests that the cards should be of postcard size with the address to which they are to be sent printed on the reverse. Immunization cards should be white and vaccination cards green. Some local health authorities have included on the cards a form of consent to be signed by the parent or guardian. This appears to be unnecessary where the vaccination or immunization is done by a practitioner at the instance of the parents in the ordinary course of his practice, and the Minister feels that it should be omitted. Other arrangements should be made when formal consent may be required for inoculations done at a local health authority centre. Among other changes suggested is a provision on the new card for the practitioner simply to indicate whether any unusual or general reaction or other sequelae to vaccination have occurred. Practitioners should be allowed to use rubber stamps in giving their name and address on the cards.

FURTHER OBSERVATIONS ON THE AGE OF THE MENARCHE

BY

DAGMAR C. WILSON, M.D., M.R.C.P., F.R.C.O.G.
D.P.H.

(From the Laboratory of Human Nutrition, University of Oxford)

AND

IAN SUTHERLAND, M.A., D.Phil.

(From the Institute of Social Medicine, Oxford)

In an earlier paper (Wilson and Sutherland, 1949) data collected from nine secondary schools in Oxford and Oxfordshire were used to study the onset of menstruation in relation to height and weight. In the first part of the present paper an analysis similar to the earlier one is made for seven schools elsewhere in the South of England, and the two sets of findings are compared. In the second part, serial records of height and weight, kept as a routine by two schools, are studied to see what information they give of growth processes in the adolescent girl.

First Investigation

The new data were obtained from five day schools and one boarding-school in Dorset and from one day school in Essex. The information consists of the ages, heights, and weights of 1,707 girls attending these schools in the summer term of 1949, together with a record of whether or not menstrual periods had started. Similar data were available for comparative purposes for 1,284 girls attending eight schools in Oxford and Oxfordshire in the autumn term of 1948.* In the case of children whose menstrual periods had already begun, the age at the menarche was also requested. But since it is possible to determine the average age of the menarche without using this extra information (which may not be fully reliable), it has been necessary to use it in only one analysis (Table III). In both surveys, children attending day schools were excluded if they had not been resident in the area for at least three years preceding the menarche or the date of the inquiry, whichever occurred earlier.

Age of the Menarche

A criticism of most studies relating to this subject is that they depend upon the girl's memory of her age at the menarche, and experience has shown that this cannot be wholly trusted even a short while after the event. A close examination of our data for Oxford and Oxfordshire shows certain discrepancies which may well be due to faulty memory. For this reason a different method of calculating the average age of the menarche, known technically as probit analysis, has been adopted. It can be used when, as in this instance, the age-range of the girls questioned is the same as that in which menstrual periods usually begin—namely, 9 to 18 years. Only two pieces of information are required from each girl—her present age and whether or not her periods have started—and these can both be recorded with a high degree of reliability.

Table I summarizes the findings concerning the age of the menarche in five groups of schools. In the day schools in Essex and in Dorset the average age of the menarche was 13.4 years, in the Oxford and Oxfordshire schools

*Information from one of the nine schools previously studied had to be discarded because it related only to those girls who had attained the menarche.