REFRESHER COURSE FOR GENERAL PRACTITIONERS**MIGRAINE**

BY

MACDONALD CRITCHLEY, M.D., F.R.C.P.*Neurologist, King's College Hospital; Physician, National Hospital, Queen Square, London*

What is migraine? Neurologists are apt to differ in their ideas upon this point. Some physicians would restrict this term to recurring unilateral headaches preceded by visual phenomena and associated with vomiting. Nowadays most would incline to a more catholic conception. More accurately one might define migraine as "a habit of recurring headaches, often one-sided, but not necessarily so; constitutional in nature, inherited, and bearing a vague association with personal or family 'allergic' tendencies and a still vaguer one with epilepsy." Taking an even broader viewpoint, however, we recall that in later life a migrainous sufferer may lose his headache; but recurring visual phenomena may betray the continuance of attacks, or attacks of headache may be replaced by bouts of Ménière-like vertigo. Thus it is that migraine without headache may occur.

Some neurologists isolate a type which might be called "symptomatic migraine." Here the patient develops migrainous attacks in middle life, associated with some morbid anatomical peculiarity, often of a vascular nature. In this connexion a cerebral aneurysm is important, but still more so a cortical angioma. For this reason some cases of migraine should be suspect, and demand special methods of study, if their true nature is to be revealed and the appropriate treatment instituted.

"Symptomatic Migraine"

What are the features which suggest a "symptomatic" as opposed to an idiopathic or constitutional migraine? In the first place, migraine-like headaches starting in the middle-aged or the elderly are not usual, and special scrutiny is advisable. Secondly, headache which is always strictly localized to one area of the skull (and does not change from one side to the other, as is so often the case in simple migraine) should be suspect. Thirdly, migrainous headaches associated with a pulsatile type of tinnitus suggest an angioma. Fourthly, migraine coupled with cranial nerve palsies, whether transient or persistent, should make one think of an aneurysm.

In all such cases there should be a particularly thorough examination of the nervous system, preferably repeated. The fundus oculi may show no abnormality at the first interview, but the wise physician will repeat his ophthalmoscopic examination from time to time. To the routine neurological testing should always be added a careful auscultation of the skull in an effort to pick up a cranial bruit. Still seeking to exclude or confirm the diagnosis of a vascular anomaly of the brain, one has recourse to various technical aids. Electro-encephalography may help by revealing a focus of abnormal cortical activity. Air-studies will occasionally prove informative. Most promising, however, is the result of percutaneous cerebral arteriography. These last two investigations will necessitate the patient entering a special neurological clinic for a few days.

The Migrainous Constitution

The belief is widespread that migraine represents an inborn type of reaction to a number of stresses, both

psychological and physical. Just as some rare persons seem to be immune from headache and may survive a long life without experiencing this particular affliction, so there are others—considerably many more—who belong to the category of "headachy subjects" like others in their family. Such persons are apt to develop their headache on every slight occasion; but in addition they may also experience a headache periodically, as it were, and without demonstrable cause. It is probable that the majority of these "headachy" subjects are victims of migraine in more or less outspoken fashion.

There have been various attempts to identify a migrainous habitus and a migrainous personality. None of these can be said to be altogether satisfying, and some of the descriptions indeed are mutually contradictory. Some would correlate migraine with a psychosomatic affection. There can be no doubt whatever that psychological factors may be effective in precipitating an attack of migraine in a migrainous subject. It is possible, too, that migrainous persons have certain identifiable psychological traits which distinguish them from the common herd. But neither of these arguments, singly or together, can justify the doctrine of migraine as a straightforward psychogenic affection, curable by psychotherapy.

Migraine as a Minor Malady

Some patients are fortunate, so far as their migraine is concerned, in that their symptoms scarcely warrant medical attention. Here belong those patients whose headache is mild in intensity and is not incapacitating. If the patient is able to continue his work with little if any falling off in efficiency, then the headache must be rated as mild. Where the headache is relieved by small doses of mild analgesics (e.g., aspirin 10 gr.—0.65 g.) the migraine must be regarded as trivial. When the attack is unaccompanied by vomiting the degree of incapacity is reduced, and unless the headache is intense the migraine may perhaps be regarded as minor in character. Very infrequent attacks—attacks which are promptly relieved by some simple common-sense measure such as the correction of refractive error—also belong within the cadre of minor migraine. Finally one may mention those cases in which the attack is represented by nothing more than short-lived attacks of teichopsia (without headache, dizziness, nausea, or vomiting).

Unfortunately, these foregoing cases would seem to be in a minority, unless it is that they exist without coming to the notice of the family doctor.

Migraine as a Major Malady

The other extreme may be considered—namely, where the migraine threatens the mental or physical health of the victim, upsets the tranquillity of the household, and interferes with working efficiency. Such cases may be spoken of as "migraine major," and they include:

(a) Cases where each attack causes much incapacity, by reason of intense and long-continued pain, loss of sleep, intractable vomiting, and anorexia.

(b) Cases of "status hemicanicus," where one attack follows another over the course of weeks.

(c) Cases of "migraine-epilepsy," in which a syncopal attack, or even an epileptiform seizure, arises in the course of the migraine.

(d) "Chronic migrainous debility," where severe attacks recur at frequent intervals and lead to a state of ill-health. The syndrome, which comprises loss of weight, poor appetite, anaemia, debility, lassitude, and depression, is perhaps most often seen in middle-aged women. It is a form of ill-health which may have far-reaching personal, domestic, and social repercussions, and the medical attendant often despairs of affording relief.

There is possibly some relationship between long-standing and severe migraine and a late development of cerebral arterial changes. This doctrine has been put forward, but so far no statistical evidence can be adduced either in support or in refutation.

When a middle-aged sufferer from migraine begins to show an increase in the severity or frequency of the attacks, or if the headache alters in quality and becomes more persistent even though perhaps milder, it is well to suspect an incipient hypertension complicating the picture.

Treatment

In the article on "Headache" contributed to this series by Dr. J. Purdon Martin the treatment of migraine was discussed. Only a few remarks are therefore necessary here. As regards the relief of an attack, one can state that in some mild cases acetylsalicylic acid will suffice. In more severe cases one will need recourse to tab. codein. co.—up to 10 tablets a day. High doses of this sort may dull the severe pain and allow the victim to continue at work. The resulting constipation and flatulency will also need palliative treatment.

The most efficacious measure is an intramuscular injection of ergotamine tartrate, given very early in the attack. It has certain drawbacks in that by reason of the natural history of the disease the first symptoms usually wake the patient unduly early—that is, at an hour when the medical attendant should still be in his bed. If the injection is delayed too long it may be relatively ineffectual. In many patients a good deal of malaise results, and vomiting is frequent. The attack of migraine may be curtailed—at least so far as headache is concerned—but the victim may have to lose a morning's work because of the drug-effect. Ergotamine tartrate by mouth, in hourly doses up to six, is often advocated, but rarely, I have found, with any great benefit. In my experience the same disappointing results have followed the oral use of drops of dihydro-ergotamine tartrate.

The patient with an acute attack of migraine will probably wish to rest in bed in a quiet and darkened room, but unfortunately sleep does not necessarily bring relief. Alcohol and smoking should be shunned during an attack, and indeed most sufferers will do their best to avoid a smoking compartment or any atmosphere tainted with tobacco smoke.

Prevention of attacks of migraine by medicinal means is often most disappointing. The total situation (including type of work, psychological stresses, visual strain) is the most potent adverse factor, which, if it cannot be avoided or suitable adjustments cannot be made, can rarely be neutralized by artificial—i.e., pharmacological—means.

Perhaps phenobarbitone is the safest bet. It should be prescribed in the form of 1 gr. (65 mg.) regularly each night on retiring. Little, if any, benefit results from the administration of this drug at the time of an attack. The patient

should persevere for two or three months before concluding that the drug is without avail. If "dopiness" is complained of in the morning, the tablet can be given a little earlier.

A large number of other preparations have been advocated for long-term treatment—thyroid, ovarian and pituitary hormones; urea; carbachol; liquor trinitrini; ergotamine tartrate; bile salts; antihistamine preparations; peptone injections; and so on. At times one or other of these seems to be beneficial, but only exceptionally. It is almost axiomatic that if a patient with migraine fails to respond dramatically to common-sense rules of hygiene, the correction of refractive errors, and phenobarbitone at night, then the doctor and the patient are both in for trouble: the case will probably prove most intractable.

Mention should, however, be made of autohaemotherapy, which is worth a trial in severe cases not responding to medicinal treatment. In some instances the benefit therefrom has been dramatic.

We are still not certain whether the so-called "histamine headache" represents a variety of migraine or not. In any case, for periodic headaches of the type described by Eszenyi-Halasy in the *British Medical Journal* of June 25, 1949 (p. 1121), a regime of desensitization with graded intramuscular doses of histamine is worth while. The patient needs to lie up during this time.

PAEDIATRICS IN THE HARVEIAN AGE

SIR LEONARD PARSONS'S HARVEIAN ORATION

Sir Leonard Parsons, F.R.S., delivered the Harveian Oration at the Royal College of Physicians on St. Luke's Day, October 18.

In taking as his subject "The Influence of Harvey and his Contemporaries on Paediatrics," Sir LEONARD PARSONS said that there was nothing he could add to the known details of Harvey's life, nor would he attempt to prove that Harvey was a paediatrician. Nevertheless, Harvey and certain of his contemporaries by their contributions to the study of children in health and disease had helped to further paediatrics. Harvey lived in a time of enormous outpouring of intellectual energy. He saw the last years of Shakespeare and the rise of Milton. A scientific revolution was in progress which overthrew mediaeval authority. The chief agents in the revolution were Galileo, Descartes, and Bacon, all of them Harvey's contemporaries. The base of the scientific revolution was Padua, which was in the sixteenth and seventeenth centuries the most famous university in the world—

"... the great desire I had

To see fair Padua, nursery of the arts."

(*Taming of the Shrew*)

It was from Padua, whose traditions had been carried on in this country by the Royal Society, that Harvey drew his inspiration.

Some idea of the state of paediatrics in the early part of the seventeenth century might be obtained from the writings of Shakespeare. Thus the mother of Coriolanus said to him: "Thy valiantness was mine; thou suck'dst it from me." Originally paediatrics formed part of obstetrics. The majority of children's diseases in the nineteenth and early part of the present century were, however, approached from the standpoint of the adult. It was true that a wide knowledge of general medicine was essential to the paediatrician, but the child was not an immature adult and required special study. When he (Sir Leonard Parsons) began the study of paediatrics at Great Ormond Street Hospital the number of babies admitted to the wards was extremely limited, and the great problems of infant morbidity and mortality were largely untouched. Within recent years it had been increasingly realized that mother and baby formed, before and after the birth, one biological unit. The trend was now towards the study of the premature child and the effect of antenatal conditions on the unborn and the newborn.