grow in tissue culture, but induced tumours in mice. This is probably due to the more favourable conditions *in vivo*, which offer better chances of recovery for the treated tumour cells than does the environment *in vitro*; it suggests that successful growth in tissue culture depends on the length of time of desiccation. Although the degree of desiccation reached after 75 minutes may be high, it is realized that dehydration is not complete. If the time of desiccation is more prolonged only few, if any, cells can be expected to survive.

It seems that three stages in the behaviour of tumour cells following different lengths of desiccation can be distinguished: (1) positive growth *in vitro* and *in vivo*; (2) growth *in vivo* but not *in vitro*; and (3) no growth *in vivo* or *in vitro*. Possibly these three stages are related to the amount of water remaining in the tumour material. In this connexion it may be mentioned that, according to Polge, Smith, and Parkes (1949), a high proportion of the water, about 90%, can be withdrawn at low temperatures from fowl semen mixed with equal quantities of 30% glycerol in Ringer's solution without killing the semen. Klinker showed by means of vital staining that the number of living cells in frozen tumour material diminished quickly after thawing followed by incubation in tissue culture. His observation suggests that our failure in tissue culture, especially after the more prolonged desiccations, may also have been due partly to the non-survival of tumour cells during incubation which otherwise would have survived under the more favourable conditions *in vivo*.

As can be seen from the table, the microscopical appearance of tumour cells, after freezing and desiccation and subsequent reconstitution in various solutions of glucose, depended probably on the individual resistance of different samples of tumour cells to the treatment, but to a greater extent on the time of desiccation. Even after the same length of desiccation the number of tumour cells which appeared damaged and the number of tumours induced in mice varied with different samples of tumour material. This may have been the result of the tumours having a different optimum time of freezing and desiccation, owing to their different general condition at the time of the experiment.

The present successful attempts show that tumour cells can withstand a high degree of desiccation when frozen and yet remain viable. They confirm the conclusion reached in the first paper—namely, that the induction of tumours in mice with frozen and desiccated suspensions of tumour cells is due to the proliferation of surviving tumour cells.

Summary and Conclusions

Suspensions of sarcoma cells which have been frozen in glucose solutions and desiccated may show active growth *in vitro* after reconstitution in glucose.

This indicates clearly that some tumour cells survive freezing and desiccation, and that the tumours which develop in mice injected with tumour material which has been frozen and desiccated arise from such surviving cells.

These experiments strengthen the conclusion reached in the first paper, that surviving cells and not a virus were responsible for the results obtained by Gye *et al.*

REFERENCES

- Gye, W. E., Begg, A. M., Mann, I., and Craigie, J. (1949). *Brit. J. Cancer*, **3**, 259.
Klinker, J. (1940). *Klin. Wschr.*, **19**, 585.
Passey, R. D., and Dmochowski, L. (1950). *British Medical Journal*, **2**, 1129.
Polge, G., Smith, A. U., and Parkes, A. S. (1949). *Nature, Lond.*, **164**, 666.

CELLULAR TRANSMISSION OF MOUSE SARCOMATA WITH FROZEN-DRIED TUMOUR TISSUES

BY

L. DMOCHOWSKI, M.D., Ph.D.

Reader in Cancer Research, School of Medicine, University of Leeds

AND

A. MILLARD

Technical Assistant in Electron Microscopy
(From the Department of Biomolecular Structure, University of Leeds)

[WITH SPECIAL PLATE]

Malignant cells of mouse tumours, after having been frozen to -79°C . and desiccated, have been grown in tissue cultures, as described in the preceding paper (Passey, Dmochowski, Lasnitzki, and Millard, 1950). This is unequivocal evidence of the viability of malignant cells after freezing and drying. There is therefore every justification for assuming that when frozen-dried tumour tissue is inoculated into mice the viable tumour cells present proliferate to form a tumour. Since the validity of this conception is crucial, it seemed desirable that the attempt should be made to confirm it experimentally. Our attention was drawn by Ludford to the advantages, for this purpose, of a technique, devised by Des Ligneris (1930), of grafting tumour tissue in plasma and serum which he described as "*in vitro* culture under *in vivo* conditions." Ludford recommended inoculating frozen-dried tumour tissue in clotted plasma, because he had found that it was possible, after inoculating cells embedded in plasma, to excise the clot after a few days and prepare from it histological sections in which the cells of the inoculum were clearly distinguishable from the surrounding tissues of the host. Experiments were therefore undertaken using this technique concurrently with those described by Passey *et al.* (1950).

Methods

The tumours used in the present experiments were the C3H and C48 sarcomata described in the original experiments of Gye *et al.* (1949) and in those of Passey and Dmochowski (1950) and of Passey *et al.* (1950). The tumour cells, in various concentrations of glucose, were submitted to storage at -79°C . and to desiccation for different lengths of time, followed by reconstitution in glucose of the same concentration as that used initially in each experiment. After reconstitution in glucose, the suspensions of tumour cells were centrifuged for three minutes at 7,000 times gravity, care being taken to keep the time of centrifugation as short as possible so as to avoid damage to cells. After discarding the supernatant fluid, equal amounts of a mixture of rat and fowl plasma (in a proportion of three parts of rat plasma and two parts of fowl plasma) and of chick embryo extract were added to the sediment (usually four to five drops of each) and gently shaken to obtain a suspension of cells in the medium. The suspension of tumour cells in the mixture of rat and fowl plasma and chick embryo extract was then put into an incubator at 37°C . for a few minutes to induce clotting. The clot was then cut into small pieces, some of which were taken for microscopical examination, and the rest grafted subcutaneously into mice by means of a trocar.

The medium used in the experiments was thus the same as that described in the tissue-culture experiments of Passey *et al.* (1950). Similarly, most of the suspensions of tumour cells were the same as those used for the tissue-culture work. Mice were sacrificed at various intervals after grafting and the clots excised and fixed in Bouin's fluid. Serial sections were stained with haematoxylin and eosin.

Eleven experiments with C3H sarcoma and one with C48 sarcoma were carried out. The accompanying Table gives the concentrations of glucose in which the tumour material was suspended, the times of freezing and desiccation, and the results of the microscopical examination of the clots excised at various intervals after grafting.

mass of the implant was necrotic and surrounded by the normal subcutaneous tissue of the mouse. By the fourth day much of the implant was absorbed, and on the fifth, sixth, and subsequent days they observed the beginnings of a new sarcoma in the connective tissue surrounding the necrotic mass. They concluded that "the appearances do not suggest survival of any of the implanted tissue, but rather a fresh sarcomatous reaction of the encapsulating subcutaneous tissues of the host."

In the present experiments these clotted suspensions of tumour cells, which on microscopical examination when first prepared showed few, if any, undamaged cells, appeared as necrotic masses in all subsequent preparations

Sarcoma and Concentration of Glucose	No. of Days Frozen	No. of Hours Desiccated	Result of Microscopical Examination of Clots at Various Time Intervals Following Grafting into Mice											No. of Positive Clots/ No. of Clots Injected
			Original Clot	6 Hours	18 Hours	1 Day	2 Days	3 Days	4 Days	5 Days	6 Days	7 Days	8 Days	
C3H 5.3%..	1	1	+	+	+	—	+	+	+	+	+	+	+	5/6
" ..	1	1	+	+	+	—	+	+	+	+	+	+	+	11/12
" ..	25	2	—	—	—	—	—	—	—	—	—	—	—	0/6
C3H 10% ..	1	1	+	+	+	+	+	+	+	+	+	+	+	0/6
" ..	25	2	—	—	—	—	—	—	—	—	—	—	—	3/6
C3H 20% ..	1	1	+	+	+	+	+	+	+	+	+	+	+	2/6
" ..	25	2	—	—	—	—	—	—	—	—	—	—	—	0/6
C3H 30% ..	1	1	+	+	+	+	+	+	+	+	+	+	+	3/6
" ..	25	2	—	—	—	—	—	—	—	—	—	—	—	3/6
C3H 30% ..	1	1	+	+	+	+	+	+	+	+	+	+	+	0/6
C48 5.3% ..	1	1	+	+	+	+	+	+	+	+	+	+	+	17/18

+ Positive clot: contains tumour cells looking apparently unharmed. — Negative clot: no tumour cells apparently unharmed. ± Indicates one positive and one negative clot. = or ≡ indicates two or three negative clots. + + or + + + indicates two or three positive clots. * Grown *in vitro* (Passey *et al.*, 1950).

As can be seen from the photomicrographs (Plate, Figs. 1–4), in many of the clots tumour cells apparently unharmed were found. An increasing number of tumour cells, some of them in division, clearly separated from the surrounding tissues of the mice, were seen in the clots taken out at intervals of 18, 24, 48, 72 hours, 4 days, and on the days following. These clots are designated as positive. Clots in which no apparently unharmed cells were found and which were ultimately absorbed are described as negative. It should be stressed that no actively proliferating cells outside the clots were seen in any of the serial sections. There is no doubt, from the appearance of the tumour cells in the first clots removed and the increasing number of cells in those taken out later, that large tumours would have resulted had growth been allowed to proceed uninterrupted. Growth in tissue culture was obtained only with the C3H tumour in 5.3% glucose frozen for 12 hours and desiccated for one hour, and with the same tumour in 20% glucose, frozen for 12 hours and dried for 30 minutes and 75 minutes (Passey *et al.*, 1950). With the grafted clots positive results were obtained with the same and also with other concentrations of glucose (see Table). The greater number of positive results obtained by inoculating the clotted suspensions of frozen and dried tumour cells into mice is not surprising in view of the more favourable conditions for growth in the animal body. According to Des Ligneris (1930) the medium in which the grafted tumour cells are embedded provides both better nourishment for the cells cut off from the normal circulation and better protection. The tumour cells within the clotted plasma induced no inflammatory reaction. Although a great many of them underwent necrosis, as could be seen in clots removed a day or two after grafting, yet a sufficient number survived and grew to form tumours.

Sections of implants of tumour tissues frozen at -79°C . were examined by Gye *et al.* (1949) at intervals of 2, 3, 4, 5, 6, and 7 days. They found, on the second day, that the

made after grafting. They were gradually absorbed *in vivo* and usually disappeared on the fifth or sixth day after transplantation, whereas, as can be seen from Figs. 1–4, clots containing cells looking apparently unharmed when grafted into mice show an increasing number of tumour cells, with many of them in division, when examined at successive intervals after transplantation. No cells outside the clots were actively proliferating, as one would expect if a virus were present capable of diffusing out and infecting the surrounding tissue of the host. Thus the experiments demonstrate that the propagation of tumours with frozen-dried tissue is entirely a process of cellular transmission.

Summary

Suspensions of frozen and desiccated sarcoma cells embedded in clots formed by the addition of chick embryo extract to a mixture of rat and fowl plasmas were inoculated subcutaneously into mice. Sections of clots excised at varying intervals after grafting revealed the presence of viable tumour cells, increasing in numbers with the age of the grafts.

We should like to express our thanks to Dr. R. J. Ludford for his valuable suggestions and helpful advice for carrying out these experiments.

Addendum.—During the preparation of the above paper for publication an additional experiment along similar lines with the C3H sarcoma (0.8 g. of tissue in 1.5 ml. of 5.3% glucose), frozen at -79°C . for 95 days and desiccated for 210 minutes, was carried out. In spite of the prolonged desiccation, an increasing number of tumour cells were seen in the clots taken out at various intervals after grafting. This observation strengthens beyond shadow of doubt the conclusion reached above.

REFERENCES

- Des Ligneris, M. J. A. (1930). *Brit. J. exp. Path.*, **11**, 249.
Gye, W. E., Begg, A. M., Mann, I., and Craigie, J. (1949). *Brit. J. Cancer*, **3**, 259.
Passey, R. D., and Dmochowski, L. (1950). *British Medical Journal*, **2**, 1129.
— — — Lasnitzki, I., and Millard, A. (1950). *Ibid.*, **2**, 1134.

L. DMOCHOWSKI AND A. MILLARD: TRANSMISSION OF SARCOMATA WITH FROZEN-DRIED TISSUES

FIG. 1.—C3H sarcoma. Minced tumour suspension in 5.3% glucose; frozen at -79°C . for 12 hours; desiccated for $\frac{1}{2}$ hour; reconstituted in 5.3% glucose; clotted in rat and fowl plasma and chick embryo extract. Appearance of clot 6 hours after grafting into mouse. H. and E. (Low power, approx. $\times 150$.)

FIG. 4.—C3H sarcoma. Minced tumour suspension in 5.3% glucose; frozen at -79°C . for 12 hours; desiccated for 1 hour; reconstituted in 5.3% glucose; clotted in rat and fowl plasma and chick embryo extract. Appearance of clot 5 days after grafting into mouse. H. and E. ($\times 1,100$.)

FIG. 2.—High-power view of area enclosed in rectangle in Fig. 1. Cells regarded as viable marked by arrows. ($\times 740$.)

FIG. 3.—C3H sarcoma. Minced tumour suspension in 5.3% glucose; frozen at -79°C . for 12 hours; desiccated for $\frac{1}{2}$ hour; reconstituted in 5.3% glucose; clotted in rat and fowl plasma and chick embryo extract. Appearance of clot 18 hours after grafting into mouse. Mitoses seen. Cells in mitosis marked by arrows. H. and E. ($\times 650$.)