on shaking with ether the blue colour could be transferred to the ether layer. (6) The acid solution formed a soluble magenta-coloured compound with diphenylcarbazine. Tests 5 and 6 were found to be the most sensitive.

Urine.—On routine testing of a specimen of urine from one of the victims a trace of albumin was found.

Test 6, which appeared to be the most sensitive test for chromium, was negative, but it was noted that, by using controlled urines, considerable quantities of chromium must be present before the above tests were positive. This is not an unusual finding in testing for some radicals in the presence of organic matter.

Source of Contamination.—Assurance is given that it did not occur in the victims' home. The package of oatmeal had travelled from Scotland to the South of England by parcel post, and it was known that it was necessary to repack it at a post office in Southern England, for a Post Office damage slip was affixed. With the above results and the possibility of legal proceedings, a sample was submitted for confirmation of the findings to Dr. J. H. Hamence, an expert analyst, who reported: "Potassium bichromate present to the extent of 0.25%."

Discussion

Fatalities following potassium bichromate poisoning, usually suicidal, are not uncommon, and Leschke (1934, p. 85) refers to a fatal case of suicidal poisoning in which a woman aged 21 took 15 g. of the substance. Taylor (1934, p. 548) records that the death of a woman aged 21 occurred in hospital about 24 hours after she drank 7 g. of potassium bichromate, which she had been incorrectly advised to try as a cure for neuralgia. Leschke (1934, p. 84) stated that 0.5 g. of potassium bichromate can cause serious symptoms of intoxication.

The early features are sore throat, vomiting, and diarrhoea. The urine frequently contains albumin and blood, and cases of only moderate severity may develop anuria with nitrogen retention, in spite of which the blood pressure remains abnormally low. Nervous symptoms, headache, stiffness of the back of the neck, and cramps are described. Haemorrhages into the intestinal tract, bronchi, and serous cavities may occur in severe cases. It is possible that these haemorrhages may be the outcome of a uraemia. It would seem that symptoms need not be proportional to the amount of potassium bichromate consumed, as the probable vomiting, and the time of its onset, must modify the amount of poison retained.

Sore throat was not a complaint in the cases described above. A trace of albumin was present in the urine, but there were no other features suggesting renal damage. The blue colour of the skin noticed by the practitioner in these cases is described (Taylor, 1934, p. 546) as a feature in some cases of potassium bichromate poisoning. The history of acute gastro-enteritis within so short a time of partaking of food (10 minutes) strongly suggests chemical poisoning. Staphylococcal food-poisoning is the form of bacterial food-poisoning with similar symptoms. Such cases, however, are typified by vomiting, diarrhoea, and acute prostration within two and a half to three hours, and are associated with well-recognized incriminating defects in the preparation of the food. The incidence, too, is much greater after certain foods, and excessively rare after oatmeal stored dry and eaten hot immediately after cooking.

Summary

Two cases of acute gastro-enteritis following food-poisoning by oatmeal contaminated with potassium bichromate are described. The features of chemical food-poisoning as compared with bacterial poisoning are mentioned, and the place of chemical food-poisoning in acute gastro-enteritis is considered.

Thanks are due to Dr. J. Hamence for his expert opinion, and to Dr. H. F. G. Hensel, who was first consulted by the patients.

REFERENCES

- Glaister, J. (1938). *Medical Jurisprudence and Toxicology*, 6th ed. Edinburgh.
Leschke, E. (1934). *Clinical Toxicology*. London.
Taylor's *Principles and Practice of Medical Jurisprudence*, 1934, edited by S. Smith, 9th ed., vol. 2. London.

Medical Memoranda

Dysgerminoma of the Ovary

The term "dysgerminoma" is applied to a type of solid ovarian carcinoma the histology of which is identical with that of a seminoma of the testis. The following is a report of two cases.

CASE 1

The patient, a girl aged 16, complained of feeling tired and of a lump in the right axilla. She had been told that "her tummy was sticking out." She had never menstruated. Examination revealed a large abdominal tumour and a firm gland in the right axilla. A radiograph of the chest was clear.

In March, 1947, a biopsy of the axillary gland was performed. Its centre was completely necrotic, and on section a few malignant cells were seen in one of the lymph sinuses, but it was not possible to make an exact diagnosis. A week later the abdomen was explored and a right-sided ovarian tumour attached to a long pedicle was found. There was some malignant infiltration of the pelvic floor and of the left side of the posterior abdominal wall. One or two enlarged glands were found in the mesentery. The tumour was removed by ligating and dividing the pedicle, and the abdomen was closed. The specimen measured 14 by 6 by 6 cm. and on microscopy was seen to be a dysgerminoma. In April the patient was given a course of x-ray treatment to the pelvic region and was discharged apparently quite well.

In December, 1947, a mass of secondary glands developed in the epigastrium. After x-ray treatment it regressed completely. In September, 1948, a mass of malignant glands developed in the left posterior triangle of the neck. A further course of x rays caused this to resolve. In March, 1949, a mass appeared in the central abdomen and left iliac fossa. With further x rays this disappeared. When last seen (October, 1949) the patient was very fit. She is, however, small and sexually infantile.

CASE 2

A girl aged 9 was admitted in April, 1949, six hours after she had been struck on the abdomen whilst playing. She had had severe epigastric pain and had vomited three times.

On admission the pain had largely passed off. A large mobile tumour was palpable in the lower abdomen. One pole of the tumour was tender. As her acute symptoms had disappeared, she was kept under observation and two days later laparotomy was performed.

An encapsulated tumour was found attached to a long pedicle arising from the left ovarian region. The ovary was incorporated in the medial aspect of the pedicle, but the growth was separate from it. The growth was removed by cutting across the pedicle, the ovary being left behind. One pole of the tumour was contused, presumably as a result of the injury. The growth was 14 by 9 by 6 cm. in size and on section was found to be a dysgerminoma.

The patient made an uneventful recovery from the operation and was seen from time to time in the out-patient department. Early in June she appeared perfectly well, but gave a history of an attack of colicky pain which had subsided spontaneously. Examination revealed nothing abnormal and she was told to report again in two months. However, she did not attend, and inquiries revealed that during July she had developed increasing abdominal distension and had died, presumably from metastases.

COMMENT

Case 1 is of interest in that it demonstrates the well-known radio-sensitivity of dysgerminomata. In spite of metastases when the patient was first seen, the disease was under control two-and-a-half years later.

Case 2 occurred in a patient comparatively young for this disease, though there have been several cases recorded in younger children. Russell (1938) reported a case in a child of 4 and McNally (1945) one in a child of 5. Novak's (1946) youngest patient was 6.

A few other extra-ovarian dysgerminomata have been reported, and, while the occurrence of the growth in such sites may be due to nests of embryonic tissue remaining behind during the course of development, it could also be explained by metastatic spread from a minute primary tumour. In Case 2 there did not seem to be any primary ovarian tumour.

The accepted treatment for localized dysgerminomata is excision of the affected tube and ovary, as recurrence in the other ovary is rare. Most authorities advise irradiation afterwards only if recurrence occurs, since it is obviously undesirable completely to destroy the patient's ovarian function, as the disease occurs in the young and the growth is usually of moderate malignancy only. The history of Case 2, however, suggests that this may not always be the wisest course. For those of adult age with hypoplasia genitalis there is obviously no objection to giving x-ray treatment post-operatively as a routine, nor is there when bilateral growths have been removed. Where metastases are present x-ray therapy gives comparatively good results, especially when one considers the poor prognosis of malignant disease arising in the young.

These cases were admitted under the care of Mr. Victor Riddell (who performed the operation on the first case) and Mr. M. F. Nicholls, and my thanks are due to them and to Mr. J. Jackson Richmond, to whom Case 1 was sent for x-ray treatment.

J. A. S. GREEN, F.R.C.S.

REFERENCES

- McNally, P. A. (1945). *J. Obstet. Gynaec. Brit. Emp.*, **52**, 193.
Novak, E. (1946). *Gynecological and Obstetrical Pathology*.
Saunders, Philadelphia.
Russell, P. M. G. (1938). *J. Obstet. Gynaec. Brit. Emp.*, **45**, 258.

Low Oesophageal Perforation at Gastroscopy

Low perforation of the oesophagus with the flexible gastro-scope is extremely rare (Schindler, 1950). The present case may therefore be of interest. It was the 320th of an otherwise uneventful series.

CASE REPORT

The patient was a wheelwright of 40 years, and of sthenic habitus. In about 1929 a gastro-enterostomy had been performed for pyloric stenosis. Since then he had been in good general health until six months before the gastroscopy, when high epigastric pain returned for two weeks. After a remission of four months a similar pain once more recurred, midline in site and passing through to the back. It was associated with epigastric fullness and belching half-way through a meal. The pain reached its maximum three hours after food; it also occurred nocturnally. There was no vomiting, no melaena, and no marked change in weight. His general condition called for no comment, his haemoglobin being 100% and a barium meal showing a well-functioning stoma and no evidence of ulceration. A stomal ulcer or persistent duodenal ulcer was, however, suspected.

Gastroscopy was attempted at 10.30 a.m. on April 13, 1949 (by C. F. J. C.), using a Hermon Taylor type with rubber

"finger-tip." Premedication consisted of "pethidine" 100 mg. with scopolamine 1/150 gr. (0.4 mg.) intramuscularly one hour beforehand; at 10 a.m. a Ryle's tube was passed and the patient given a tablet of "decicain" 50 mg. to suck. The instrument passed easily for over three-quarters of the distance, the patient showing no signs of distress. For about five minutes efforts were made by inflation, by flexion of the gastroscope, and by minor adjustments of the degree of extension of the head to obtain a field of vision, but none was forthcoming. During this period the patient, on being questioned, indicated that he was comfortable. The instrument was then withdrawn: there was a smear of blood on the rubber finger-tip.

At about noon the patient began to complain of increasing chest pain; by 2 p.m. he was very shocked and in great distress—sweating profusely, pale, restless, and with a thready pulse. His general and mental condition recalled the clinical picture of a recent coronary thrombosis. The pain appeared to be chiefly substernal, radiating to the back, up to the neck, and down to the left hypochondrium. There was marked epigastric tenderness and rigidity. In the left supraclavicular fossa surgical emphysema was visible and palpable, confirming our fears that instrumental perforation had taken place. Pressure over the sternum caused pain.

The abdominal pain with unyielding abdominal muscles suggested that the injury was a low one and probably close to the diaphragm. An x-ray film was taken in the erect posture, and no air could be seen under the cupulae, but the mediastinum was spongy with air bubbles, presumably pumped in at gastroscopy. A second film was taken after a small barium swallow (which enhanced the substernal pain): this showed an extravasation of the opaque medium just above the diaphragm into the mediastinal tissues on the right side. Operation revealed a ragged perforation in the gullet large enough to admit a fountain-pen, and about 1 in. from the diaphragm. The gullet wall at this point was thin and papery. There was no haematoma or other sign of injury, suggesting that the end of the gastroscope had temporarily stretched this point just on the diaphragmatic hiatus, piercing it, and had then been withdrawn. The tear was sutured, and the patient made a good recovery, but on the eighth day it became evident that a fistula had formed between gullet and operation track. The patient began to decline and died on the 26th day.

At necropsy the oesophagus showed a rent 1 in. (2.5 cm.) long on its antero-lateral aspect, at a point $\frac{1}{4}$ in. (1.9 cm.) above the diaphragm. The edges had separated and there was little sign of repair. On the anterior surface of the stomach 1 in. above the site of the previous gastro-jejunostomy was a recent perforation, $\frac{3}{4}$ in. across, of a chronic peptic ulcer. The splenic flexure of the colon, the tail of the pancreas, the hilum of the spleen, and the upper pole of the left kidney were tightly bound to one another and the stomach by dense fibrous tissue, in places almost 1 in. thick. This had abolished their mobility, and had bent and distorted the cardiac orifice of the stomach; which explained why the gastro-scope had been unable to pass beyond the oesophagus.

COMMENT

(1) Pathological immobility of the oesophagus or cardia in the region of the diaphragm adds to the inherent risks of gastroscopy, and, if possible, its presence and degree should be assessed before instrumentation. (2) A "barium swallow" is of assistance in locating the site of an oesophageal perforation. (3) It seems evident that the tear would not in this case have healed under a conservative regime.

C. F. J. CROPPER, F.R.C.P.Ed.,
Senior Physician, Ronkswood Hospital.

P. HOUGHTON, F.R.C.S.,
Senior Surgeon, Ronkswood Hospital.

P. KIDD, M.B., B.Ch.,
Pathologist, Worcester Royal Infirmary.

REFERENCE

- Schindler, R. (1950). *Gastroscopy*, p. 92. University of Chicago Press.