REFRESHER COURSE FOR GENERAL PRACTITIONERS

ENURESIS

BY

DOUGLAS HUBBLE, M.D., M.R.C.P.

Physician, Derbyshire Royal Infirmary, Derby City Hospital, and Derbyshire Hospital for Sick Children

Involuntary micturition is the normal infantile habit both by night and by day. Children should acquire day-time control in the second year, and during the third year the dry bed should also be achieved. If, in the absence of disease, repeated involuntary micturition occurs after the third year, then to this disorder of function the name enuresis is given. To this same symptom occurring in organic disease, whether in children or in adults, the description "incontinence of urine" is applied.

By its frequency alone, enuresis must be regarded as the most important disorder of function in childhood, and when the social consequences of bed-wetting are considered its importance as a potential disturber of development in childhood can hardly be exaggerated. Every family doctor is familiar with the unhappy story. Even in young children the failure to attain control of the bladder causes much distress to parents, and presents the child with a physical situation to which it has to make some mental adaptation. The mother's unhappiness is intensified by the washing of sheets and night-clothes—an unwelcome addition to her daily work. When the condition persists for a few years, as it often does, anxiety regarding the future is added to the depression concerning the present state.

Normal Control of Micturition

There is still uncertainty about the exact nature of the physiological control of micturition. In the automatic bladder of infancy and disease micturition is accomplished through the agency of a spinal reflex which causes relaxation of the sphincter followed by a mass contraction of the muscular coat of the bladder—the detrusor. In the adult the intravesical pressure is normally maintained at 15 ml. of water; as the bladder fills with urine this pressure is kept constant by relaxation of the detrusor. When the volume of urine exceeds a certain limit the pressure begins to rise, and at 18 ml. of water the impulse to micturate is recognized as a result of rhythmical bladder contractions. This impulse is inhibited, as Denny-Brown and Robertson showed in cystometric experiments on themselves, by cortical inhibition of the spinal reflexes. While the existence of this cortical inhibition is certain, other investigators have suggested that, in addition, cortical impulses will promote detrusor activity, while others report that stimulation of higher centres will cause relaxation of the bladder.

In cystometry the bladder is slowly filled through a urethral catheter attached to a manometer on which the intravesical pressure is recorded. In the adult the desire to micturate occurs when the bladder contains about 3 oz. (85 ml.), and again at about 6 oz. (170 ml.), but this impulse can be overcome by deep breathing and is not accompanied by a rise of pressure. As the bladder is increasingly filled the need becomes urgent and eventually cortical inhibition is overcome and involuntary micturition occurs. In the adult the impulse to micturate at these levels is inhibited by cortical impulses, but in the infant this cortical control is only slowly developed.

Ellison Nash describes four stages in this progress: the automatic bladder of infancy; the small bladder of increasing capacity in which the impulse to void is still uninhibited

(high-tension, uninhibited); the bladder which will hold 3 oz. or more, but in which the impulse is still succeeded by micturition (low-tension, uninhibited); and the "adult" bladder in which the contractions are inhibited by cortical activity (low-tension, inhibited). In the cystometric investigation of a large number of children suffering from functional enuresis, Nash found that nearly half of them showed the normal adult pattern, while the remainder fell in the other three groups. The only conclusion one can draw from this is that, while the bladder control in these children, as tested by cystometry, may show evidence of physiological retardation or regression, it may, on the other hand, be indistinguishable from that of normal persons, whether children or adults.

Incidence of Enuresis

The incidence of enuresis is variously estimated as being between 5 and 15% in the child population. If the higher figure is accepted, it must include children who are occasional and intermittent bed-wetters; if the lower figure, then it includes only those whose enuresis is unremitting for long periods. Most observers agree that boys are somewhat more often afflicted than girls, while children of all classes suffer from it. Enuresis may persist into adult life, but there is a tendency to spontaneous cure towards adolescence. There are no reliable figures for the incidence of enuresis amongst adults, but it is certainly less than 1%.

History and Examination

Children suffering from enuresis can immediately be separated into two groups—those whose bed-wetting has persisted from infancy, and those in whom nocturnal control of micturition had been completely established but in whom, at a later age, regression to an infantile habit has occurred. From published figures it may be assumed that about one in every three or four children brought to the family doctor on account of enuresis has for some reason relapsed from a state of adequate control. In many children there is no complaint of day-time disturbance of micturition, but in about one-third of them questioning will reveal that bladder control is not perfect by day. The most common symptom is frequency, the next urgency, and in some actual day-time enuresis occurs. Children in this group have been described as having "irritable bladders." Physical abnormalities in these children should be sought for, but when found they should not be too easily assumed to have aetiological significance. Thirty years ago some authorities regarded the presence of thread-worms or of spina bifida occulta as a pathogenetic factor. To-day they are held to be of no aetiological importance. Incontinence of faeces occurs occasionally in children who have enuresis.

Children suffering from enuresis should pass their urine in a free stream; if, from the mother's account, there is a possibility that they suffer from dribbling incontinence, they should be watched while micturating and suprapubic palpation and pressure should be applied after micturition. If a clean specimen of urine contains pus cells, then a catheter specimen must be obtained; if this, too, contains pus cells a complete urological investigation must be made.

The genitalia should be closely inspected and abnormalities noted. Evidence of skin inflammation and mucous membrane hyperaemia is not infrequent in chronic enuretics. There are some who maintain that such changes precede the enuresis or at least provide adequate reason for its persistence. Most observers, however, regard these changes as secondary and not causal. To describe an area of hyperaemia in the urethra or bladder-neck as urethritis or trigonitis (in the absence of pyuria) is as outmoded as to believe that hyperaemic gastric mucosa deserves the description of gastritis. Adhesions around the corona or the clitoris have been thought to be significant by a few clinicians.

Children who suffer from nocturnal enuresis are said to sleep deeply. This observation was first made by Trousseau, and most parents can be persuaded to agree with him. It should be said, however, that we have no index for profundity of sleep, and that most young children, whether bed-wetters or not, are awakened with difficulty.

History-taking should include an account of the child's physical development and mental growth. His psychological adjustments to his parents, siblings, companions, and teachers should be inquired into, and his performance at work and play assessed. The nature of his progressive adaptation to his disorder is important. Special consideration should be given to the personality of the parents, particularly of the mother. With American children it is thought that excessive emphasis by an over-anxious or obsessional mother on the early acquisition of a continent habit is the commonest cause of enuresis in children. Evidence of a similar sort may sometimes be obtained in this country. Conversely, in overcrowded homes and with negligent mothers, early training may occasionally be shown to have been defective. Parents whose personalities are not unusually deviated and who have an unequivocal affection for their children are not likely to harm them either by apparent neglect or by overemphasis on training. The occurrence of other manifestations of instability—negativism, temper-tantrums, masturbation, habit-spasms, day-dreaming, nail-biting, and stammering—is not infrequent. F. G. Still pointed out that acquired enuresis occurs most commonly in children aged from 5 to 8 years—at a time when such functional nervous disorders as habit-spasm and stuttering are apt to appear.

When the enuresis is acquired an opportunity arises to get an answer to one of Halliday's famous questions, "Why he took ill when he did." The answer usually lies in one of the normal stresses of childhood—an illness, an operation, the birth of a sibling, the temporary or permanent loss of a loved one, and so on. When the incapacitating environmental situation is disclosed the interest shifts to an attempt to discover why this particular form of somatic expression was chosen. Has the child shown any previous instability of bladder control? Did the parents or other relatives suffer from enuresis? A history of enuresis of some degree in the father or mother will be obtained in more than one-third of all cases. The mother of an enuretic child told me recently that she had wet the bed until she was 18 years old, that her brother had been in trouble while in the Army for the same reason, and that her mother had been enuretic when married.

Differential Diagnosis

If enuresis is defined as a functional disorder of micturition, then an obligation is laid upon both family doctor and paediatrician to exclude organic disease in all patients presenting with urinary incontinence. Fortunately, routine history-taking, physical examination, and microscopy of the

urine are sufficient to reveal those conditions which will require further investigation.

If the enuresis is acquired, then general disorders such as diabetes insipidus, diabetes mellitus, chronic nephritis, epilepsy, and disease of the spinal cord must be remembered and excluded. Petit mal may be accompanied by diurnal incontinence, but the story of the transient attacks of unconsciousness is not easily missed. Urinary infection is an occasional cause of acquired incontinence, and it requires continuous investigation until its aetiology and cure are established. When enuresis has persisted from infancy, whether intermittently or persistently, then organic disease of the brain such as a mental defect, with or without paresis, should be excluded. Enuresis is not more common in children with defective intelligence than in those with normal cerebration. Braithwaite found that the intelligence quotient in 60 enuretic children was of normal range. In enuresis which has persisted for many years, however, the percentage of persons with a low I.Q. is increased, and in such cases this must be regarded as a finding of aetiological significance.

The most important differential diagnosis is a lesion of the urinary tract. Much valuable work in the elucidation of these conditions has been done in recent years by urologists, and by their warning writings they have made those who treat enuresis in children suitably aware of the diagnostic problem. The actual incidence of these surgical conditions presenting as urinary incontinence is variously estimated. The urologists describe it as 10% of all cases of enuresis—this is certainly a much higher incidence than will be found in patients presenting in the ordinary hospital out-patient department or in the consulting-room. The family doctor will not miss these cases if he will make certain of three facts: (1) that the child is not suffering from dribbling incontinence; (2) that the bladder cannot be palpated in the suprapubic region after the child has micturated; and (3) that there are no pus cells in the child's urine.

Aetiology

The aetiology of enuresis is both debatable and debated, and these controversies will not be continued here.

In a functional disorder with such an extraordinarily high incidence it is certain that multiple aetiological factors are at work; these should be considered under three heads—constitutional, environmental, and local.

The constitutional factor in enuresis is ill defined (nor has it been much elucidated by cystometry), but it may reasonably be assumed to reside in some instability of neuromuscular control. The frequent finding of enuresis in relatives seems to support the existence of a constitutional factor, though the possibility that this finding depends on an environmental association cannot be excluded.

That the environmental factor is often a psychological one is clear from the histories of children who revert to enuresis after being "dry." But the environmental situation in the second and third years is not easily determined in retrospect. A disorder of function such as enuresis that has lasted for months and years has inevitable results which encourage its persistence: on the one hand are the neurological factors implicit in chronic habituation, and on the other the emotional reactions which ensue both in parents and in children. The disorder persists long after the precipitating situation has disappeared and been forgotten.

It is possible that conditions giving rise to reflex irritation may occasionally precipitate enuresis; it does not seem reasonable or necessary to assume that they could continue

as direct causes over a number of months or years, nor does it seem reasonable to accept such conditions as a tight foreskin, threadworms, coronal and clitoral adhesions, or a hypertrophied verumontanum as relevant in pathogenesis.

Treatment and Prognosis

The weapons in the doctor's armoury for the control of enuresis have changed but little in the course of 30 years, and there is not much in the following description of treatment that would not have been familiar to F. G. Still. These weapons are (1) psychotherapy—suggestion, explanation, and encouragement; (2) bladder-training; and (3) drugs—belladonna, ephedrine, phenobarbitone, and amphetamine.

Psychotherapy.—The family doctor should exchange his physicianly scepticism for a surgical confidence in the potentialities of successful therapy. Parents and children must be persuaded out of a wearied and fatalistic acceptance of their unhappy situation. Anxiety must give place to confidence, and hope must cast out shame. While the personalities of mother and child should be assessed, no suggestion of blame should be attached to either. The child should be encouraged to make his own record of the dry nights, and a system of rewards may be usefully instituted. No recourse should be made to the psychotherapist unless the situation plainly demands it or unless the methods outlined here have failed. It must not be forgotten that there is a bad habit to be broken, and until it is removed a profoundly dysgenic situation remains. The symptom itself is the cause of a psychological disturbance more important than any which may have produced it, and the doctor's plain duty is to get rid of it. The battle against enuresis is a long one, and the doctor should play his cards one by one, always keeping a therapeutic trick up his sleeve.

Bladder-training.—The restriction of the child's intake of fluid has not been shown to have any therapeutic value, but since there is some evidence that polyuria occurs in these children they should, if not asleep, be made to micturate within an hour of going to bed. They should later be thoroughly awakened at the parents' bed-time and made to pass water. Micturition should be made a fully conscious act, and to put a stuporous child on the pot is to perpetuate a condition which one is seeking to alter. Older children and adults may be given an alarm-clock to waken them at some time between 10.30 p.m. and rising. A system of day-time training should be instituted in the holidays. The child should be sent to the lavatory at fixed but gradually increasing intervals, beginning at one hour and increasing the interval by a quarter of an hour every second day. If the child can hold its water for three-hour periods in the day, dryness at night is facilitated. This method is most successful in children with irritable bladders who have day-time frequency.

Drugs.—These are much more useful in therapy than some allow them to be. Sometimes they fail because they are casually and inefficiently applied. Suggestion is valuable in the control of enuresis, and since these remedies have a rational basis the physician's belief in their reasonableness will undeniably enhance their therapeutic powers. Drugs should be pushed to the limit of the child's tolerance before they are discarded. It is not uncommon to find that some children can tolerate a dose of 1–2 gr. (65–130 mg.) of ephedrine or 15–20 min (0.9–1.2 ml.) of belladonna. Ephedrine is the most successful drug; it reinforces the sphincteric action. Methyl ephedrine is more prolonged in its action than ephedrine and has fewer side-effects. For a child aged 5–10 years, $\frac{3}{4}$ gr. (45 mg.) should to begin with

be given in the morning, at tea-time, and at the parents' bed-time. Belladonna may be used at the same time (if ephedrine fails when given alone) and in increasing dosage until it produces dryness of the mouth or paralysis of accommodation, when the dose should be reduced. Phenobarbitone in doses of $\frac{1}{4}$ or $\frac{1}{2}$ gr. (16 or 32 mg.) should be given to the nervous child who suffers from day-time frequency, together with ephedrine or belladonna. The child who is difficult to awaken may be given amphetamine, 2.5–10 mg., at bed-time, in order to lighten the depth of sleep. In some children ephedrine has a similar effect.

Associated conditions such as balanitis or vulvo-vaginitis should be treated whether or not they are thought to be affecting the enuresis. Stilboestrol in doses of 0.5–1 mg. given daily for two to three weeks is often effective in curing vaginitis. Enuresis should not be treated by circumcision. The occasional wet bed may be ignored, but a long relapse will require a fresh approach to the problem, with the institution of the old remedies and the introduction of new methods.

Chronic Enuresis

The measures outlined above may be expected to be successful in most cases of acquired or intermittent enuresis. There will remain a number of children with chronic enuresis—possibly representing one-third of all cases—who are unaffected by such treatment. They should be removed if possible to the home of a relative prepared to tackle the problem, to a residential school, or to a special boarding-school. It has often been found that measures which failed at home can be successfully applied elsewhere. In the days when hospital beds could be found for such children most of them achieved dryness during their stay in hospital. During the war, when enuretic evacuee children constituted a widespread social problem, Miss A. M. Maynard reported from the Quedley Hospital, Haslemere, where these children were taken for treatment, that "almost every case has been cured temporarily, and many permanently." There is little doubt that six months in a residential school would be expected to cure most of those children whose enuresis cannot be controlled at home.

Recruiting board and Service experience has shown that a considerable number of enuretic children carry their disability into adult life. The problem in enuretic adults and in children with chronic enuresis is essentially the same, for in all adults with functional incontinence of urine the disorder has persisted since childhood. It is obvious that the longer a bad habit has persisted the more difficult it is to break, but to compensate for this difficulty there are positive factors which will facilitate successful therapy in chronic enuresis. For example, there should be present, even in persons with I.Q.s between 70 and 100, a greater wish to be cured. It is also possible to apply sterner disciplinary measures, such as the regular waking through the night by alarm-clock. Davidson and Douglass* report favourably on an electric apparatus which wakens the patient by the ringing of a bell at the start of micturition when the bag on which he is lying becomes wet. The success of such a device, which may well be due partly to an increased confidence that there will be no soaking of the bed and partly to a natural distaste for being harshly called from sleep to get out of bed, to turn off a bell, to micturate, and to change a wet pad for a dry one, illuminates the nature of the mental processes which are perpetuating this disorder of function.

Ström-Olsen reports that amphetamine is very successful in the treatment of enuresis in adults. He emphasizes

*British Medical Journal, 1950, 1, 1345.

again that large doses may be required, varying from 10 to 30 mg. given at bed-time. Before any person is dubbed an incurable enuretic, almost any therapeutic method is justifiable, however fallacious its aetiological foundation. Such methods as the injection into the bladder of increasing quantities of fluid have been shown to be successful by some observers, and may well be tried before hope is abandoned. If all fails, the unfortunate is condemned in perpetuity to the nightly companionship of the mackintosh sheet and the rubber-bag urinal.

EIGHTIETH BIRTHDAY OF DR. DAIN

Dr. Guy Dain celebrated his eightieth birthday on November 5. His outstanding services to the profession in administration and medical politics have earned the gratitude of medical men and women increasingly legislated upon by successive Governments in the last 40 years. Dr. Dain was early a shrewd counsellor and later the protagonist in the struggle waged by doctors to prevent their traditions and professional independence from being submerged in State bureaucracy. Though now playing a smaller part in B.M.A. affairs, his great experience is still

often placed at the disposal of the Association, and he is a familiar figure at many committees.

Born and educated in Birmingham, where he has been in general practice all his life, Dr. Dain began his medico-political work when National Health Insurance was introduced in 1911, becoming a member of the first insurance committee and of the first panel committee. At that time his persuasiveness and tact did much to allay the hostility of many members of the insurance committee towards representatives of the medical profession.

By H. J. Whitlock, Birmingham

In 1917 he brought his knowledge of this work into the ambit of B.M.A. Headquarters when he became a member of the Insurance Acts Committee, of which he was chairman for 12 years. He also presided for five years over the annual conference of Local Medical and Panel Committees.

In 1936 he was awarded the Gold Medal of the Association for his services to the profession, though they were soon to be continued more strikingly. The following year he was elected to the chair of the Representative Body, which he filled at the Plymouth and Aberdeen meetings, and in 1943 he became Chairman of Council, holding that important office for six of the most stormy years through which the profession has ever passed.

Among other activities that he has packed into a busy life are his many years of service on the General Medical Council, to which he was first elected in 1934 as a direct representative of the medical practitioners of England and Wales, and his service on the Council of the Royal College of Surgeons as the general-practitioner representative: the College elected him an honorary Fellow in 1945.

The current interest—not only in this country—of hospital and health care on a regional basis has led to a revived demand from students of hospital administration and others for the *Interim Report on the Future Provision of Medical and Allied Services* (1920), commonly referred to as the Dawson Report. Since the Report has long been unobtainable from H.M. Stationery Office, King Edward's Hospital Fund for London has, with their permission, had a limited number reprinted. Copies may be obtained on application to the Fund at 10, Old Jewry, London, E.C.2.

PUBLIC HEALTH IN WESTERN UNION COUNTRIES

A CHADWICK TRUST LECTURE

A Chadwick Trust lecture was given at Westminster Medical School on October 24 by Dr. NEVILLE M. GOODMAN, a senior medical officer of the Ministry of Health, on public health administration in the countries—Belgium, France, Luxembourg, and the Netherlands—whose Governments, with the Government of Great Britain, signed in 1948 the treaty of economic, social, and cultural co-operation. Under that treaty a permanent commission with a small secretariat was set up in London, with committees of experts in various subjects, one of them public health administration, and it was as a member of that committee that Dr. Goodman recently visited the four countries in order to gain a general picture of their developments in this field.

National Outlooks on Administration

Dr. Goodman said that perhaps a distinction between the French and Belgian attitude towards administration and the British attitude was that France and Belgium regarded administration as a science and Great Britain regarded it as an art. The French and Belgian approach was logical though sometimes not very practical; the British approach was practical but nearly always illogical. The British put emphasis on personnel, the French on the proper definition of positions and functions. Again, the Dutch tradition of individual and local self-reliance and effort had fostered the conservation of voluntary effort in public health and other fields. The French and Belgian systems owed much to the Napoleonic concepts of administrative law and centralization. The British tradition of sanitary reform, as essentially a matter for local authorities, had fixed the broad lines of our public health system on a local basis.

In Luxembourg public health administration was entirely centralized. Its Minister of Health performed many duties which would be undertaken by the permanent head of the department in other countries. The medical director had no medical assistants except the director of a laboratory. He dealt with epidemiology, State hospitals for tuberculosis, mental health, and maternity services, and he was concerned with or advised on a great variety of other questions. He carried out all the duties which the medical officer of health of an English county would carry out. The ability and energy of the present Minister went very far to make up for exiguous staffing. In Luxembourg the coal and iron industries had their own medical services. All doctors were trained outside the country; 70% of students were trained in France, 15% in Belgium, 10% in Switzerland, and a few went to Germany. Medical and pharmaceutical practice was supervised by a medical college.

Structure of Health Ministries

In France there was a Ministry of Public Health and Population, in Belgium a Ministry of Public Health and the Family, in Holland a Ministry of Social Service. The British was the earliest and most stable of these central public health administrations. All four Ministries had functions additional to health—in Great Britain housing and relations with local authorities which elsewhere were questions for the Ministry of the Interior; in France, the Ministry handled demography and emigration, but not social security. In Belgium the Health Ministry dealt with housing, physical education, and sport.

Dr. Goodman went on to point out various differences in structure and organization of the Ministries. In all the countries except the United Kingdom, doctors and other technical health staff were eligible for and in fact held technical positions on a level with non-technical administrators. The permanent head of the Ministry in each country was a doctor. A special and important feature peculiar to France, which dated from pre-revolutionary days, was a corps of general inspectors. There were nine of these, five of them doctors