technique appears to be a safe and effective method of thoracic drainage post oesophagectomy. A large, randomised, controlled trial is required to assess fully the benefits and limitations of transhiatal chest drains.

References

1. Laws D, Neville E, Duffy J, and the British Thoracic Society Pleural Disease Group. BTS guidelines for insertion of a chest drain. *Thorax* 2003; **58** (Suppl II): ii53–9.
2. Gossage JA, Chukwuemeka AO, Dussek JE. Intercostal drain migration post esophagectomy. *Dis Esophagus* 2003; **16**: 268–9.
3. Law S, Boey J-P, Kwok K-F, Wong K-H, Chu K-M, Wong J. Pleural drainage after transthoracic esophagectomy: experience with a vacuum system. *Dis Esophagus* 2004; **17**: 81–6.

TECHNICAL TIPS

A useful connector for constant irrigation pressure in arthroscopic surgery

AM THIMMIAH, WILLEM SCHENK

Department of Orthopaedics, West Suffolk Hospital, Bury St Edmunds, Suffolk, UK

CORRESPONDENCE TO

Willem Schenk, Consultant in Trauma and Orthopaedics, West Suffolk Hospital, Hardwick Lane, Bury St Edmunds, Suffolk IP33 2QZ, UK

E: marion.gladwell@wsh.nhs.uk

therefore, developed a connector between the pressure infuser bag and one lead of the dual-channel, self-compensating, tourniquet inflation unit. We typically pressurise up to 100 mmHg for routine arthroscopies. This simple and cheap device has proven to be very useful by both surgeons and theatre staff in providing a constant clear view during arthroscopic procedures.

The 'horizontometer'

D MAHADEVAN¹, R BUCKINGHAM², MF GARGAN³

¹Department of Trauma and Orthopaedics, Derriford Hospital, Plymouth Hospitals NHS Trust, Plymouth, UK

²Department of Trauma and Orthopaedics, Avon Orthopaedic Centre, Southmead Hospital, Bristol, UK

³Department of Trauma and Orthopaedics, Bristol Royal Infirmary, Bristol, UK

CORRESPONDENCE TO

Devendra Mahadevan, Department of Trauma and Orthopaedics, Derriford Hospital, Plymouth Hospitals NHS Trust, Plymouth PL6 8DH, UK

E: m.deven@gmail.com or m_deven@hotmail.com

Maintaining constant pressure during arthroscopic surgery by manually pumping a pressure infusion bag is tedious, tiring and may cause repetitive strain injuries to the forearm; loss of pressure during the procedure is common. We have,

Figure 1 use of the connector.

Orientating fluoroscopic images during surgery can be testing when the visual field is small or lacking definite reference points. A simple and effective technique is described to assist orientation during hip arthrograms and peri-acetabular osteotomies. The anterior superior iliac spines are palpated and a metal ruler

Figure 1 Fluoroscopic image of right hemipelvis with metal ruler representing the true horizontal line of pelvis.

placed between these two landmarks. This projects a true horizontal reference line. The image viewed on the screen can then be rotated until the line produced by this radio-opaque marker is indeed horizontal (Fig. 1). This technique allows accurate orientation of the weight-bearing surface of the acetabulum and avoids miscalculation of the amount of coverage of the femoral head.

Safety first in the use of 'K'-wires

T NUNN, J KRIEK, P ROLLINSON

Department of Trauma and Orthopaedics, Ngwelezane Hospital, Empangeni, Kwa-Zulu Natal, South Africa

CORRESPONDENCE TO

T Nunn, Department of Trauma and Orthopaedics, Ngwelezane Hospital, Empangeni, Kwa-Zulu Natal, South Africa
E: timrachnunn@yahoo.co.uk

Kirshner ('K') wires are a useful tool in trauma and orthopaedics. The sharp cut end, however, risks intra-operative glove puncture and needle-stick injury to surgeon and assistant. This is a heightened problem where there is a high HIV seroprevalence. In Kwa-Zulu Natal, South Africa, we have instituted a policy to 'clip, cut, then condomise' (Fig. 1) the sharp cut wire end with a section of infant feeding catheter (IFC) appropriate to the

diameter of the 'K'-wire. We have found the following sizes to fit best: 1.2 mm 'K'-wire, CH 4/5 IFC; 1.4–1.8 mm 'K'-wire, CH 8 IFC; and 2 mm 'K'-wire, CH 10 IFC.

A simple manoeuvre that avoids kinking of chest tubes

SUSHIL DAWKA

Department of Surgery, Manipal Teaching Hospital, Pokhara, Nepal

CORRESPONDENCE TO

Sushil Dawka, Associate Professor, Department of Surgery, Manipal Teaching Hospital, Phulbari, Pokhara, Nepal.
E: sushil.dawka@gmail.com

A chest tube kinked within the pleural space can cause poor drainage, discomfort and trauma on removal, and may require repositioning. With declining usage of trocars for chest tube placement, kinking may occur more frequently. Rotation of the tube around its long axis during insertion ensures that any tendency of the tip to deviate will be annulled by its spiralling forward progression in the direction of advancement. After placement, a straight or gently curved tube can be rotated freely. An angulated tube resists rotation and will twist back when released unless, in the process, it straightens out.

Figure 1 Sections of 8CH feeding tube fit and protect the sharp cut ends of two 1.8 mm 'K'-wires during articular reconstruction.

The use of cough to sample detrusor muscle during flexible cystoscopic biopsy

HDI DE'ATH, O KAYES

Department of Surgery, Northwick Park Hospital, Harrow, UK

CORRESPONDENCE TO

Henry De'Ath, 137 Wymering Mansions, Wymering Road, London W9 2NF, UK
E: HenryDe-Ath@doctors.org.uk

During flexible cystoscopy, a biopsy may be required. Whilst it is useful to sample detrusor, controlling biopsy depth is frustrating and samples will frequently not yield muscle. We describe a simple technique that will ensure detrusor muscle sampling. During cystoscopy, the biopsy area is located and flexible forceps are introduced into the bladder. With the forceps open, push against the target region such that the bladder wall is slightly indented. The patient is asked to cough and, whilst simultaneously maintaining constant pressure against the wall, close the forceps. With the instrument's mouth closed, again ask the patient to cough and swiftly withdraw the forceps. This successful method is very easy to use and conjures no additional cost. Furthermore, we have observed that patients' subjective perception of pain is lowered.