Paramedic practitioners and emergency admissions

Evidence suggests a positive effect, but future programmes need rigorous assessment before being expanded

Research, p 919

Malcolm Woollard professor in prehospital and emergency care, Coventry University, Richard Crossman Building, Coventry CV1 5FB
Malcolm.woollard@coventry.ac.uk
Competing interests: None declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2007;335:893-4
[doi:10.1136/bmj.39356.700139.BE](https://doi.org/10.1136/bmj.39356.700139.BE)

In this week's *BMJ*, Mason and colleagues report a cluster randomised controlled trial examining the effects of a "paramedic practitioner" service in a UK urban setting.¹ The trial focused on managing older patients without life threatening conditions who accessed the emergency ambulance service. It aimed to increase the proportion receiving care in the community and reduce admissions to the emergency department. It found that people in the intervention group were less likely to attend the emergency department (relative risk 0.72, 95% confidence interval 0.68 to 0.75) or need hospital admission within 28 days (0.87, 0.81 to 0.94). However, use of secondary care services after the initial episode increased (1.21, 1.06 to 1.38).

Paramedic practitioners undertook a three week theory course followed by 45 days of supervised clinical experience. Their scope of practice was restricted to common presentations considered unlikely to result in serious injury, including falls, lacerations, epistaxis, and minor burns. Skills acquired beyond those normally practised by UK paramedics included wound care and suturing; examination of the joints; examination of the neurological, cardiovascular, respiratory, and ear, nose, and throat systems; social needs assessment; administration of antibiotics, simple analgesics, and tetanus toxoid; and referral of patients for radiography or to a general practitioner, district nurse, or social services.

As long ago as 1994, it was reported that services that deputised for general practitioners often could not cope with the demand for out of hours consultations.² In addition, calls for emergency ambulances have been rising persistently by as much as 8% each year,³ yet half of patients taken to emergency departments by ambulance are discharged without being treated or referred.⁴ The increased demand for emergency ambulances may have been exacerbated by the introduction of general practitioner contracts that do not require the provision of out of hours services, although this has not been proved. Ambulances often have to queue outside emergency departments because a lack of trolley bays means that patients cannot be admitted. This in turn results in a lack of ambulances to respond to further emergency calls. Similar problems have occurred in Australia, New Zealand, the United States, and Canada. This, along with the need to tackle the challenges of providing health care to remote populations, has led to the formation of

the International Roundtable on Community Paramedicine (www.ircp.info).

Emergency care practitioners were introduced with the aim of reducing admissions to the emergency department for a broader range of patients than those discussed by Mason and colleagues.⁵ A pilot programme began in the Warwickshire Ambulance Service in 2002, with the support of Coventry University and the changing workforce programme. It was subsequently expanded to encompass 17 pilot sites.⁶ Although this was intended to promote a standardised 15 week university based course, with all practitioners having a similar scope of practice, several centres opted for alternative approaches, including that described by Mason and colleagues.¹ Most emergency care practitioners are paramedics working in ambulance services, but some are nurses or physiotherapists, and most work in out of hours and primary care services or emergency departments. Similar paramedic based programmes in other countries include those of the Queensland and New South Wales Ambulance Services in Australia.

Few high quality evaluations of the emergency care practitioner scheme and other extended scope paramedic practitioner schemes have been published to date. A changing workforce programme review reported that emergency care practitioners could reduce admissions to emergency departments by 100 to 358 patients each year in rural and urban settings, respectively. For a training investment of £24 250 (£34 800; \$50 000), this would save the National Health Service £62 000-£72 000 each year for each practitioner.⁶ In Wales, autonomous "advanced paramedic practitioners," who are educated to masters degree level, responded to about 25% (n=635) of 999 calls received in one primary care trust area. Of these, 292 (46%) patients were treated and discharged at the scene; 75 (12%) were re-graded to non-emergency transport; and 34 (5%) were admitted to hospital destinations other than the emergency department. Although emergency care practitioners are typically targeted to calls triaged as "low priority" by ambulance dispatchers, the Welsh scheme found that the greatest proportion of patients with altered care pathways fell within high and medium priority categories (24% and 22%, respectively, v 17% in the lowest priority category).⁷

In a scheme in the west of England, 48/170 (28%)

patients were treated “on scene” by emergency care practitioners with a BSc in emergency care, compared with 60/331 (18%) attended by paramedics.⁸ A more recent evaluation found that 62% of patients seen by emergency care practitioners were not admitted and 38% were referred; practitioners self reported that their intervention prevented an acute admission in 66% of cases.⁹

The trial by Mason and colleagues is the first adequately powered randomised controlled trial to investigate the effect of an extended scope paramedic programme on admissions to the emergency department.¹ Although their results are largely positive, they cannot be generalised beyond the scheme evaluated. Perhaps, most importantly, the trial shows that high quality study designs are feasible in this setting. Such trials should, therefore, be used to evaluate the more widespread emergency care practitioner scheme and other extended scope paramedic programmes before further costly expansions take place.

- 1 Mason S, Knowles E, Colwell B, Dixon S, Wardrope J, Gorringer R, et al. Effectiveness of paramedic practitioners in attending 999 calls from elderly people in the community: cluster randomised controlled trial. *BMJ* 2007 doi: 10.1136/bmj.39343.649097.55.
- 2 Hallam L, Cragg D. Organisation of primary care services outside normal working hours. *BMJ* 1994;309:1621-3.
- 3 Department of Health. *Statistical bulletin ambulance services, England: 2003-4*. 2004. www.dh.gov.uk/en/Publicationsandstatistics/Statistics/StatisticalWorkAreas/Statisticalhealthcare/DH_4083989.
- 4 Volans AP. Use and abuse of the ambulance service. *Prehosp Immed Care* 1998;2:190-2.
- 5 Department of Health. *Reforming emergency care*. 2001. www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4058836.pdf.
- 6 Modernisation Agency/Department of Health. *The emergency care practitioner report*. 2004. www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4093088.pdf.
- 7 Woollard M. The role of the paramedic practitioner in the UK. *J Emerg Primary Health Care* 2006;4:990156. www.jephpc.com/full_article.cfm?content_id=337.
- 8 Cooper S, Barrett B, Black S, Evans C, Real C, Williams S, et al. The emerging role of the emergency care practitioner. *Emerg Med J* 2004;21:614-8.
- 9 Cooper S, O'Carroll J, Jenkin A, Badger B. Collaborative practices in unscheduled emergency care: role and impact of the emergency care practitioner—quantitative findings. *Emerg Med J* 2007;24:630-3.

Occupational therapy after stroke

Improves personal activities of daily living, but evidence remains sparse on other potential effects

RESEARCH, p 922

Kathryn M McPherson

professor of rehabilitation (Laura Fergusson chair), Health and Rehabilitation Research Centre, Auckland University of Technology, Private Bag 92006, Auckland 1142, New Zealand
Kathryn.McPherson@aut.ac.nz

Caroline Ellis-Hill senior lecturer in rehabilitation, School of Health Professions and Rehabilitation Sciences, University of Southampton, Southampton SO17 1JB

Competing interests: None declared.

Provenance and peer review: Commissioned, not externally peer reviewed

BMJ 2007;335:894-5

doi: 10.1136/bmj.39370.600729.BE

In this week's *BMJ*, Legg and colleagues present a systematic review of randomised trials assessing the effect of occupational therapy on personal activities of daily life in people who have had a stroke.¹

Stroke is a leading cause of death and disability² and sadly, despite medical advances and public health initiatives, its incidence is not declining.³ The sequelae are often devastating and can affect the full range of human life and functioning.⁴ The impact of stroke extends beyond the individual to the people closest to them—carers experience high rates of distress, depression, and social isolation; reduced health status; and even premature death.⁵⁻⁸ Prevention and acute medical management are a fundamental part of the response to the problem, but improving life after stroke is also important and, as increasing evidence indicates, possible.⁹

Occupational therapists have for many years been identified as key contributors to the rehabilitation of people with stroke. The 2004 national clinical guidelines for stroke¹⁰ stated that “a specialist stroke team should include staff with specialist knowledge of stroke including an occupational therapist.” Despite the fact that their input has long been seen as important, the level of evidence for this recommendation was limited to expert committee reports, opinions, or experience of respected authorities, with an indication that directly applicable clinical studies of good quality were absent.¹⁰

The review by Legg and colleagues¹ synthesises data collected over the past decade and finds that

occupational therapy targeted towards activities of daily living significantly increased performance on scores of personal activities of daily living (standardised mean difference 0.18, $P=0.01$) and reduced the risk of poor outcome (deterioration or dependency in personal activities of daily living: odds ratio 0.67, 95% confidence interval 0.51 to 0.87). The review, which includes a wide range of studies, provides strong evidence that occupational therapy after stroke improves outcome and prevents deterioration in functional performance. As many people struggle with the consequences of a stroke for years, policies should ensure a satisfactory level of service provision and appropriate staff capacity, and should optimise appropriate referral and management from health professionals.

Legg and colleagues¹ focused on the effect of occupational therapy targeted at enhancing activities of daily living. Although this is a key intervention, it is just one aspect of occupational therapy after stroke. Other activities undertaken by therapists aim to enhance participation in domestic and leisure activities and facilitate engagement in activities that people find meaningful—occupation in the widest sense of the word. As a result, assessing occupational therapy along with other aspects of rehabilitation is complex, and the evidence from Legg and colleagues focuses on an aspect that is perhaps the easiest to capture. The challenge ahead is to build on and maintain this evidence base.

As Legg and colleagues note,¹ key questions remain

about which occupational therapy interventions are most beneficial and for whom. In the meantime, occupational therapists, medical practitioners, and others involved in the care of people with stroke should be heartened by these findings. For therapists, the evidence clearly shows that occupational therapy does make a difference. For general practitioners and community service providers, referral to occupational therapy may make a real difference for stroke survivors and their families. Not everyone will benefit from occupational therapy, however, and challenges include identifying those who will benefit and deciding exactly how to provide services.

So what needs to happen next? Undoubtedly, more people who have had a stroke should be referred to occupational therapy for help with activities of daily life. We still don't know which people are most likely to benefit (age, type of stroke, location of insult, duration of time since stroke); whether occupational therapy makes a difference to outcomes such as participation in meaningful activities like work and other life roles; and which outcomes are most important to people who have had a stroke and whether they are accounted for in rehabilitation interventions.¹¹

Rehabilitation has often been labelled as comprising "black box" interventions¹² and being a "Cinderella" discipline.⁹ The promising results from Legg and colleagues' trial should stimulate research into other areas of occupational therapy practice

and encourage better provision of such services for people who have had a stroke.

- 1 Legg L, Drummond A, Leonardi-Bee J, Gladman JRF, Corr S, Donkervoort M, et al. Occupational therapy for patients with problems in personal activities of daily living after stroke: systematic review of randomised trials. *BMJ* 2007 doi: 10.1136/bmj.39343.466863.55.
- 2 Rosamond W, Flegal K, Friday G, Furie K, Go A, Greenlund K, et al. Heart disease and stroke statistics—2007 update: a report from the American Heart Association statistics committee and stroke statistics subcommittee. *Circulation* 2007;115:e69-171.
- 3 Kleindorfer D. The bad news: stroke incidence is stable. *Lancet Neurol* 2007;6:470-1.
- 4 Hochstenbach J, Prigatano G, Mulder T. Patients' and relatives' reports of disturbances 9 months after stroke: subjective changes in physical functioning, cognition, emotion, and behavior. *Arch Phys Med Rehabil* 2005;86:1587-93.
- 5 Berg A, Palomaki H, Lonnqvist J, Lehtihalmes M, Kaste M. Depression among caregivers of stroke survivors. *Stroke* 2005;36:639-43.
- 6 Van Heugten C, Visser-Meily A, Post M, Lindeman E. Care for carers of stroke patients: evidence-based clinical practice guidelines. *J Rehabil Med* 2006;38:153-8.
- 7 Grant JS, Glandon GL, Elliott TR, Giger JN, Weaver M. Caregiving problems and feelings experienced by family caregivers of stroke survivors the first month after discharge. *Int J Rehabil Res* 2004;27:105-11.
- 8 Christakis NA, Allison PD. Mortality after the hospitalization of a spouse. *N Engl J Med* 2006;354:719-30.
- 9 McPherson KM, Kersten P, McNaughton H, Turner-Stokes L. Background to neurorehabilitation. In: Candelise L, Hughes R, Liberati A, Uitendag B, Warlow C, eds. *Management of neurological disorders: an evidence-based approach*. Edinburgh: Blackwell Publishing, 2007:100-9.
- 10 Intercollegiate Stroke Working Party. *National clinical guidelines for stroke*. 2nd ed. London: Royal College of Physicians, 2004.
- 11 McPherson KM, Brander P, Taylor WJ, McNaughton HK. Consequences of stroke, arthritis and chronic pain—are there important similarities? *Disabil Rehabil* 2004;26:988-99.
- 12 Dejong G, Horn SD, Gassaway JA, Slavin MD, Dijkers MP. Toward a taxonomy of rehabilitation interventions: using an inductive approach to examine the "black box" of rehabilitation. *Arch Phys Med Rehabil* 2004;85:678-86.

Device therapy in heart failure

Whether combined cardiac resynchronisation and implantable cardioverter defibrillator therapy offers benefit above either device alone remains uncertain

RESEARCH, p 925

Finlay A McAlister associate professor, Division of General Internal Medicine, Department of Medicine, University of Alberta, Edmonton, AB, Canada T6G 2R7
Finlay.McAlister@ualberta.ca

Competing interests: None declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2007;335:895-6
doi: 10.1136/bmj.39353.546933.BE

This article was posted on bmj.com on 11 October 2007: <http://bmj.com/cgi/doi/10.1136/bmj.39353.546933.BE>

Over the past decade, two devices have emerged as adjuncts to optimal medical treatment for people with heart failure—cardiac resynchronisation therapy, which is delivered by atrial synchronised biventricular pacemakers, and implantable cardioverter defibrillators. The evidence base underpinning these treatments has evolved rapidly. In this week's *BMJ*, Lam and Owen have summarised mortality data from 12 randomised trials using a Bayesian network meta-analysis.¹

Their analysis confirms earlier meta-analyses showing that both devices independently improve survival in patients with heart failure (these earlier analyses also showed that resynchronisation therapy, but not implantable cardioverter defibrillators, improved functional status and reduced hospital admissions). More importantly, Lam and Owen found insufficient evidence to prove that combined devices were better than either device alone even using network meta-analysis incorporating all major trials. This analysis has four important implications for clinicians.

Firstly, while these devices work in selected patients (see table on bmj.com), their role in many patients with heart failure remains uncertain. Only 1-3% of people with heart failure meet the eligibility criteria for the trials that proved the efficacy of these devices.² The criteria were New York Heart Association (NYHA) class III or IV symptoms despite optimal medical management, left ventricular ejection fraction $\leq 35\%$, electrical dyssynchrony, and sinus rhythm for trials of cardiac resynchronisation, whereas criteria were NYHA class II or III symptoms and left ventricular ejection fraction $\leq 35\%$ for trials of implantable cardioverter defibrillators.

In addition, not all people eligible for trials will benefit. For example, 41% of trial participants had no improvement in functional status after successful implantation of their cardiac resynchronisation device,³ and two thirds of patients with implantable cardioverter defibrillators never received therapeutic defibrillation discharges.⁴ Only 10% of patients who receive an implantable cardioverter defibrillator for

primary prevention have been estimated to receive an appropriate shock and survive at least one year.⁵ Thus, we urgently need to develop and validate cheap and easily applied tools that can accurately identify those people most likely to benefit from either (or both) of these devices.

Secondly, although the benefits of both devices seemed to be relatively stable over time in the randomised trials (survival curves of patients with and without the devices did not converge as follow-up lengthened), these trials were of relatively short duration. Thus, long term follow-up data are essential to confirm that the benefits do not attenuate over time, particularly in light of the frequency with which Food and Drug Administration advisories and device recalls arise years after efficacy studies are completed.⁶ Everyone who has one of these devices implanted should therefore be entered into a registry and followed for long term risks and benefits. This would have the added benefit of tracking changes in the success rates and complications of implantation procedures over time as the sophistication of devices and technical expertise of those implanting them evolves.

Thirdly, data are lacking on the safety and effectiveness of either device when applied outside the confines of a clinical trial. These trials could possibly have overestimated the potential benefit-safety ratio because half of the people randomised were drawn from trials in which they were randomised after successful implantation of the device (thus, the 7% implantation failure seen for cardiac resynchronisation devices and 0.5-1% peri-procedural mortality seen with both devices are not factored into the effect estimates presented by Lam and Owen). Moreover, while the trials proving the efficacy of both devices enrolled relatively young patients, these devices are now being implanted into older people with more comorbidities.⁷ They are also being implanted by less experienced providers working in hospitals with lower implant volumes.⁸ Moreover, people with heart failure with less severe symptoms (NYHA class I or II) or common comorbidities (such as bradyarrhythmias, atrial fibrillation, or chronic kidney disease) were largely excluded from the published trials. Again, this emphasises the potential value of a prospective registry in defining the benefits and safety of these devices in a wider spectrum of patients.

Finally, the finding by Lam and Owen of uncertain incremental benefits for the combined device over single devices is important. Is it plausible that a combined device would not be better than either device alone? The answer is yes. Increases in the use of angiotensin converting enzyme inhibitors and β blockers over the past decade mean that people with heart failure are now more likely to die of progressive heart failure than sudden death.⁹ Also, cardiac resynchronisation alone seems to reduce the frequency of ventricular arrhythmias¹⁰ and—at least in the trial with the longest follow-up—the incidence of sudden death.¹¹

While several trials are ongoing to define the incremental benefits of combined devices over implantable cardioverter defibrillators alone in patients with heart failure, I agree with Lam and Owen that further randomised trials are needed to define the incremental benefit of combined devices over cardiac resynchronisation alone. As both devices have been shown to prevent morbidity and mortality in similar patient populations (see table on bmj.com), some may argue that such trials are not needed and would advocate the use of combined devices in all patients except those with severe comorbidities. However, consideration of the costs of the two options (£5074 (€7249; \$10 253) for cardiac resynchronisation therapy versus £17 266 for a combined device)¹² brings the matter clearly into focus—in publicly funded healthcare systems with finite budgets, should we reallocate resources from other areas to pay for expensive technologies that offer uncertain incremental benefits over cheaper options?¹² Until a trial definitively proves that a combined device is better than cardiac resynchronisation alone, such devices should be reserved for people with heart failure who are eligible for cardiac resynchronisation and have the highest risk of sudden cardiac death.

- 1 Lam SKH, Owen A. Combined resynchronisation and defibrillation therapy in left ventricular dysfunction: Bayesian network meta-analysis of randomised controlled trials. *BMJ* 2007 doi: 10.1136/bmj.39343.511389.BE.
- 2 McAlister FA, Tu JV, Newman A, Lee D, Kimber S, Cujec B, et al. How many patients with heart failure are eligible for cardiac resynchronization? Insights from 2 prospective cohorts. *Eur Heart J* 2006;27:323-9.
- 3 McAlister FA, Ezekowitz J, Hooton N, Vandermeer B, Dryden DM, Spooner C, et al. The role of cardiac resynchronization therapy in patients with left ventricular systolic dysfunction: a systematic review. *JAMA* 2007;297:2502-14.
- 4 Ezekowitz J, Dryden DM, Hooton N, Vandermeer B, Friesen C, Spooner C, et al. Implantable cardioverter defibrillators for adults with left ventricular systolic dysfunction: a systematic review. *Ann Intern Med* 2007;147:251-62.
- 5 Stevenson LW. Implantable cardioverter-defibrillators for primary prevention of sudden death in heart failure: are there enough bangs for the bucks? *Circulation* 2006;114:101-3.
- 6 Maisel WH. Safety issues involving medical devices: implications of recent implantable cardioverter-defibrillator malfunctions. *JAMA* 2005;294:955-8.
- 7 Reynolds MR, Cohen DJ, Kugelmass AD, Brown PP, Becker ER, Culler SD, et al. The frequency and incremental cost of major complications among Medicare beneficiaries receiving implantable cardioverter-defibrillators. *J Am Coll Cardiol* 2006;47:2493-7.
- 8 Al-Khatib SM, Lucas FL, Jollis JG, Mallenka DJ, Wennberg DE. The relation between patients' outcomes and the volume of cardioverter-defibrillator implantation procedures performed by physicians treating Medicare beneficiaries. *J Am Coll Cardiol* 2005;46:1536-50.
- 9 Teuteberg JJ, Lewis EF, Nohria A, Tsang SW, Fang JC, Givertz MM, et al. Characteristics of patients who die with heart failure and a low ejection fraction in the new millennium. *J Cardiac Failure* 2006;12:47-53.
- 10 Ermis C, Seutter R, Zhu AX, Benditt LC, VanHeel L, Sakaguchi S, et al. Impact of upgrade to cardiac resynchronization therapy on ventricular arrhythmia frequency in patients with implantable cardioverter-defibrillators. *J Am Coll Cardiol* 2005;46:2258-63.
- 11 Cleland JGF, Daubert JC, Erdmann E, Freemantle N, Gras D, Kappenberger L, et al. Longer-term effects of cardiac resynchronization therapy on mortality in heart failure (the cardiac resynchronization-heart failure (CARE-HF) trial extension phase). *Eur Heart J* 2006;27:1928-32.
- 12 Fox M, Mealing S, Anderson R, Dean J, Stein K, Price A, et al. The effectiveness and cost-effectiveness of cardiac resynchronization therapy (biventricular pacing) for heart failure: a systematic review and economic model. *Health Technol Assess* (in press).

Diet and the risk of cancer

New report shows obesity and alcohol are the strongest risk factors

Tim Key professor, Cancer Research UK Epidemiology Unit, Oxford University, Oxford OX3 7LF
tim.key@ceu.ox.ac.uk

Competing interests: TK is a member of the Vegan Society.

Provenance and peer review: Commissioned based on an idea from the author; not externally peer reviewed.

BMJ 2007;335:897
doi: 10.1136/bmj.39373.676204.BE

This week, the World Cancer Research Fund (WCRF) launches its second report on diet and cancer through simultaneous conferences in London and Washington. Entitled *Food, Nutrition, Physical Activity, and the Prevention of Cancer*, the report updates the previous publication from the charity. It is the culmination of five years' work by scientists in nine universities from four countries who have assessed the original research according to a specially developed standardised review protocol.¹

The possible influence of diet on the risk of cancer is constantly topical. The subject is important because people can change their diets, and even a moderate effect on risk could prevent several thousand cancers each year in a country the size of the United Kingdom. However, apart from the confirmed adverse effects of alcohol and obesity on the risk for some types of cancer,²⁻⁵ progress in understanding has been slow and the evidence remains confusing.

The conclusions in the WCRF report about the effects of obesity and alcohol are similar to those reached by other expert consultations.¹⁻⁵ Obesity increases the risk of cancer of the oesophagus, colorectum, pancreas, breast, endometrium, and kidney. The report's public health goal for obesity is for the median body mass index of the population to be between 21 and 23. Mean body mass index in adults in the UK is now about 27.⁶ Enormous efforts by individuals, society, and government will be needed to reverse the current trend and approach the goal proposed—mean body mass index in the UK has not fallen into the target range since the 1940s. The mean body mass index is also in the overweight range in most other Western countries and is rising rapidly from previously low levels in many developing countries.⁷

Alcohol increases the risk for cancers of the mouth, pharynx, larynx, oesophagus, colorectum, and breast and also causes cirrhosis, which predisposes to liver cancer. Globally, alcohol consumption has increased in recent decades, with the highest consumption in Europe and North America.⁷ The WCRF panel made recommendations that men should not drink more than two units of alcohol a day and that women should not consume more than one unit a day. These recommendations are much lower than current government advice in Britain of up to three or four drinks a day for men and two or three for women.⁸ A substantial shift in drinking habits would be needed to achieve these goals.

The role of fruit and vegetables remains poorly understood. The panel concluded that the evidence that diets rich in vegetables and fruits protect against cancer is overall less compelling than it was in the mid-1990s. They judged that vegetables, fruits, and other foods containing dietary fibre probably protect against several cancers, but that the evidence was not convincing for any specific cancer. Despite the absence of convincing evidence for a protective effect,

the panel reiterated the previous recommendation that people should eat at least five portions of vegetables and fruits each day, and pointed out that, to meet this individual recommendation, the average consumption in the population would need to be about 7.5 portions a day. Currently, only a minority of the world's population consumes the recommended five portions of fruit and vegetables a day, and consumption is low in many regions of the developing world.⁵ The average consumption in Britain is currently about three portions a day,⁹ so the consumption of fruit and vegetables would need to be more than doubled.

The major new conclusion in the report is that red and processed meat convincingly cause colorectal cancer. The previous WCRF report concluded that the evidence for this link was "probable," and the 1998 Department of Health report concluded that the evidence was "moderate."^{2,3} This new conclusion is based mainly on results from prospective observational studies, supported by results from case-control studies. On average, people who ate the most red or processed meat had about a 30% increased risk compared with those who ate the least. On the basis of their conclusion, the panel proposed a public health goal that the average intake of red and processed meat in the population should be no more than 300 g each week. Consumption of meat has been rising in most of the world, and total meat consumption is well above the 300 g per week goal in most regions except Africa and South Asia.⁵ The current mean intake of red and processed meat in Britain is about 970 g a week in men and about 550 g a week in women,⁹ so reaching this goal would require a large reduction in meat consumption.

Many challenges remain. Obesity and excessive alcohol intake definitely increase the risk of cancer but are hard to control, while our understanding of whether and how particular foods affect risk is still rudimentary. The goal for intake of vegetables and fruits requires careful consideration—achieving the goal may not reduce rates of cancer, whereas the financial and environmental costs could be large. Vegetables and fruits provide only small amounts of energy and protein, so an increase in intake may be largely on top of the rest of the diet. This could be expensive for people, and it has implications for the optimal use of agricultural land, water, pesticides, transport, etc. It may be better to concentrate efforts on increasing the consumption of plant foods such as cereals and beans, which supply energy and protein and can therefore partially replace meat. The production of staple plant foods is usually more efficient than the production of meat, so replacing some meat with plant foods should reduce the demands on the environment.¹⁰ These wider questions will be dealt with in the WCRF's policy report, due to be published next year.

All references are on bmj.com

Mental Capacity Act 2005

Should guide doctors to help protect vulnerable people

Andrew Alonzi senior lecturer in law, Nottingham Trent University, Nottingham NG1 5LP

Mike Pringle professor of general practice, University of Nottingham, Nottingham NG7 2RD
mike.pringle@nottingham.ac.uk

Competing interests: None declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2007;335:898

doi: 10.1136/bmj.39372.691076.80

The Mental Capacity Act 2005, which was fully implemented on 1 October 2007, is intended to protect people who lack capacity to make decisions and to encourage them to participate in the decisions that are intended to help them. It presents a new range of challenges for doctors, but it should help clarify actions in difficult situations.

A helpful code of practice that supports the act guides people treating or caring for adults who lack the capacity to make decisions about their treatment or care,¹ and it is all most doctors will need. This code has statutory force, which means that practitioners now have a legal duty to take note of the code when working with or caring for adults who lack capacity to make decisions for themselves. However, there is no legal duty to comply with the code, although evidence of non-compliance may be used by a court or tribunal—for example, in a claim for damages for clinical negligence.

Several principles underpin the act. A person must be assumed to have capacity to make decisions unless evidence to the contrary is available. All practical steps must be taken to help someone make a decision before incapacity is assumed. Just because someone makes an unwise decision does not mean that they are incapacitated. If you make a decision on someone's behalf, it must be in their best interests, and there must be no better way of achieving the same outcome.

Who lacks mental capacity? The answer is any person who is unable to make a particular decision or take a particular action at the time the decision or action needs to be taken. A doctor has to judge whether the functioning of the mind or brain is sufficiently impaired or disturbed, permanently or temporarily, that the person lacks the capacity to make a particular decision.

How does a doctor act in the patient's best interest? To determine what is in a person's best interests, the doctor should encourage the person to participate in the decision making process; try to identify the views and factors that would have influenced them if they had capacity to make the decision (such as evidence of past and present wishes and feelings; religious, cultural, or moral beliefs and values); avoid assumptions about best interests based on age, appearance, condition, or behaviour; consider whether they are likely to regain capacity after medical treatment (which implies that the decision could be delayed until then); and not be motivated by a desire to bring about death or be influenced by assumptions about quality of life, if the decision is about life sustaining medical treatment.

The doctor should consult with others—while being careful not to breach confidentiality—for their views when trying to determine what is in the person's best interests, and to see if they have any information about the person's wishes, feelings, beliefs, and values. Whatever is decided must also be the least restrictive option available to the person, in terms of their rights and freedoms. The

doctor must record how the decision on best interests was reached (for example, the reasons, who was consulted, and what special factors were considered).

Can a doctor be liable for his or her decision? If certain conditions are met, doctors are protected from civil or criminal liability when performing tasks in the best interests of a person who lacks capacity. In emergency situations, for example, it will almost always be in the person's best interests to give urgent treatment without delay, unless a valid and applicable advance decision to refuse medical treatment is in place.

This protection does not, however, provide defence against a claim of negligence. For example, if a person who lacks capacity is restrained, the act does not protect against liability unless it can reasonably be believed that restraint is needed to prevent harm to the person. The amount and type of restraint used and the time that it lasts must also be proportionate to the likelihood and seriousness of harm.

The act also covers advance decisions to refuse medical treatment. If a valid, applicable advance decision to refuse treatment exists a doctor must respect it. A doctor following an advance decision will be protected from liability. If the doctor knows such a decision exists and does not follow it, the doctor will not be covered. Even if an advance decision is not valid or applicable to the circumstances, it still should be considered by healthcare staff as an expression of previous wishes when working out the person's best interests.

Whenever treating a person who might have impaired capacity, the code needs to form a conscious part of a doctor's decision making process. Clear evidence of alignment with the code's guidance must exist, particularly if decisions or treatments are called into question. This will require education for staff at all levels. Health professionals should be encouraged to reflect on their practice in the context of the code's guidance and to discuss experiences openly and frankly.

1 Department of Constitutional Affairs. *Mental Capacity Act 2005. Code of practice*. www.opsi.gov.uk/acts/acts2005/related/ukpgacop_20050009_en.pdf.