Summary

Chronic mastitis is a term which has lost all precision and value and should be abolished. There are two common non-malignant conditions of the breast met with clinically—cystic disease and the pain syndrome—which are separate and distinct conditions. Cystic disease presents clear objective physical signs from which a clinical diagnosis can be made, the cause is unknown but is possibly a hormonal upset, there is no special liability to carcinoma of the breasts affected, and tapping is a satisfactory treatment. The pain syndrome is a purely subjective disorder, although sometimes superimposed on exaggerated physiological premenstrual engorgement of the breasts, and the breasts affected present no objective abnormality. The cause of the fixation on the breast is usually fear of carcinoma, but in a small proportion of cases is a psychosexual upset. Treatment by explanation and reassurance usually results in cure of the pain.

REFERENCES

- Atkins, H. J. B. (1940). *Lancet*, **2**, 411.
 Birkett, J. (1850). *The Diseases of the Breast*. London.
 Bloodgood, J. C. (1931). *Ann. Surg.*, **93**, 235.
 Brodie, B. C. (1846). *Lectures in Pathology and Surgery*. London.
 Bryant, T. (1887). *The Diseases of the Breast*. London.
 Bucher, N. L. R., and Geschickter, C. F. (1941). *J. clin. Endocrinol.*, **1**, 58.
 Campbell, O. J. (1934). *Arch. Surg.*, **28**, 1001.
 Cheate, G. L., and Cutler, M. (1931). *Tumours of the Breast*. London.
 Cooper, Astley (1928). *Illustrations of the Diseases of the Breast*. London.
 Deaver, J. B., and McFarland, J. (1918). *The Breast, its Anomalies, its Diseases, and their Treatment*. Philadelphia.
 Geschickter, C. F. (1943). *Disease of the Breast*. Philadelphia.
 König (1893). *Cbl. Chir.*, **20**, 49.
 Paget, J. (1870). *Lectures on Surgical Pathology*. London.
 Patey, D. H. (1939). *Surg. Gynec. Obstet.*, **68** (Int. Abstr. Surg.), 575.
 Réclus, P. (1893). *Sem. méd.*, **13**, 353.
 Schimmelbusch, C. (1892). *Arch. klin. Chir.*, **44**, 117.
 Spence, A. W. (1940). *Lancet*, **2**, 387.
 Taylor, H. C. (1942). *Surg. Gynec. Obstet.*, **74**, 326.
 Velpeau, A. (1856). *A Treatise on the Diseases of the Breast and Mammary Region*. London.
 Willis, R. A. (1948). *The Pathology of Tumours*. London.

AMETHOCAINE HYDROCHLORIDE

SEVERE TOXIC EFFECTS WHEN USED FOR BRONCHOSCOPY

BY

C. A. JACKSON, F.R.C.S.

Chief Assistant, St. Bartholomew's Hospital Chest Unit,
 Hill End Hospital and Clinic, St. Albans

Amethocaine hydrochloride is increasing in popularity in this country as a surface analgesic. The occurrence of two severe toxic reactions arising from its use in a thoracic unit while securing analgesia for bronchoscopy has emphasized again the danger of overdosage when applied to a mucous surface, together with a specific sensitivity shown by certain individuals. These near tragedies have prompted me to review similar cases reported in the literature and to study the properties of this drug. Since amethocaine has been used for surface analgesia twelve deaths at bronchoscopy and gastroscopy have been reported by various authors and a further three deaths during preparation for urethral instrumentation.

Whether it be due to sensitivity or overdosage, the nature of the reaction reveals a striking similarity in the majority of cases and is the typical classic picture of cocaine or procaine intoxication. Within a few moments of local application the patient becomes faint and giddy, and unconsciousness follows. There may be wandering or deviation of the eyes. A trembling and twitching of the extremities occurs

which rapidly develops into generalized clonic convulsions, epileptiform in nature, during which cyanosis, foaming at the mouth, and an accelerated weakened pulse are noted. This seizure may subside with recovery or may return with a greater violence, when the patient may succumb from respiratory or cardiac failure.

Cases in the Literature

There have been a number of case reports in German and American journals since 1935 giving instances of severe constitutional disturbances and even fatalities. Apart from two deaths recorded by Hancock (1939), the only British report is found in a statistical review of causes of sudden death in which two fatalities are recorded as having occurred when this analgesic was being used in bronchoscopy.

In 1939 Fasselt reported two reactions to amethocaine 2% in anaesthetizing the throat before the removal of foreign bodies by bronchoscopy. All symptoms, however, disappeared spontaneously in these cases. Phillips and his associates (1941) reported a similar reaction in a patient who was undergoing a bronchoscopic examination. In this case generalized tonic contractions occurred which were controlled by "dial," 2 ml., given intramuscularly; the strength of the solution was not mentioned. The patient recovered.

Schoen (1939) reported three fatal reactions to local analgesia using amethocaine prior to bronchoscopy. Wagener is reported by him to have quoted a case in which 6 ml. of 2.5% solution was applied to the larynx for the removal of a wart and in which general convulsions occurred, leading to death in six to eight minutes.

In a case quoted by Cazzaniga (Schoen, 1939) a total dose of less than 130 mg. of amethocaine was used, yet the patient, a 27-year-old woman with pulmonary tuberculosis, developed severe convulsions some seconds after the initial administration and death occurred thirty minutes later.

Freeman (1939) reported four serious accidents with amethocaine employed as a mucous membrane analgesic for gastroscopy. Hancock (1939) had two deaths from the drug, also used for gastroscopy. His patients were both in advanced cachexia from malignant disease of the stomach.

Pfeiffer (1937) and Shumacher (1941) reported sensitivity to 0.5 and 1% solutions when administered in the eye. Videbeck (1936) mentioned three cases of amethocaine eczema in doctors who used the drug in their practices.

Thomas and Fenton (1943) provide a series of cases of their own in which amethocaine was shown to possess definite toxic properties. The first patient gave a history of asthma but no history of drug sensitivity. During bronchoscopy, while receiving a second spray of amethocaine 2% five minutes after the first, he had a sudden severe asthmatic attack and died thirty minutes later. A woman giving a history of pulmonary tuberculosis of one year's duration and with no known allergy, after receiving 5 ml. of amethocaine solution by pharyngeal spray became faint and suffered a generalized convulsive seizure with frothing at the mouth, which persisted until respirations ceased. Two other patients, both women, underwent bronchoscopy with amethocaine local analgesia for investigation of bronchiectasis. Feeble respirations, cyanosis, weak pulse, and clonic convulsions developed in both patients. Only after the administration of adrenaline and oxygen did these patients return to consciousness. In both cases skin tests for amethocaine sensitivity proved positive when performed afterwards. A 58-year-old man, after a gastric resection for carcinoma, was being prepared for gastroscopy with a 2% amethocaine gargle, to be followed by a 5-ml. spray, this to be repeated

in ten minutes. About fifteen minutes later the patient became stuporous, suffering clonic convulsions involving the arms and legs, lacrimated, and there was drooling of saliva and grinding of the teeth. The skin was cold and clammy, the lips were cyanotic. This reaction lasted ten minutes, but the patient recovered. One patient, ten minutes after receiving 5 ml. of 2% amethocaine prior to gastroscopy, became pulseless and convulsive, developed jerky respirations, and died. Three other cases were marked by nystagmus and convulsive seizures, but all three patients survived.

Doane and Cohn (1945) reported the case of a 48-year-old woman who suffered severely from asthma. Sixteen days before admission under their care a successful bronchoscopy had been performed, using 2% amethocaine as a spray. Five days after admission she was being prepared for a bronchogram; 2 ml. of a 2% solution were instilled by a nasal catheter, no premedication being given. A further 1 ml. of the solution was being instilled eight minutes later, when the patient slumped forward unconscious and in another three minutes slight convulsive movements occurred. This seizure was relieved by 2 g. of sodium phenobarbitone intravenously, but soon returned with greater violence. A double dose of barbiturate and cardiac restorative measures failed and the patient died thirty minutes later.

Richards (1947) recorded two cases of sudden death following the application of a 2% amethocaine spray for the purpose of bronchoscopy—one in a patient with a lung tumour and the other in a case of bronchiectasis. In both cases generalized convulsions set in within a few minutes and death occurred very shortly afterwards.

Shindler (1940) and Holt (1945) each report a fatality.

Properties of the Drug

Amethocaine, known variously as "pantocaine," "ponto-caine," "anethaine," "dikaine," "tetracaine," and "decicain," belongs like procaine to the para-aminobenzoyl group of anaesthetics and was introduced as decicain about eighteen years ago by Bayers.

Advantages.—It is resistant to boiling, it is stable in solution, it is lethal to all non-spore-forming organisms, it is ten times more potent than procaine or cocaine, and it is relatively quick in achieving its action and provides prolonged analgesia. Although the onset of analgesia is slower than with cocaine, the effect is more persistent.

Disadvantages.—It penetrates the tissues more rapidly than procaine and does not produce the vasoconstrictive action of cocaine. It is therefore essential to use it in conjunction with adrenaline, which reduces its toxicity by one-fifth. The relative toxicity of amethocaine, cocaine, and procaine is 4.5, 3, and 1 respectively, but this disadvantage is mitigated by the effective dose being one-tenth that of cocaine. It is detoxicated more slowly than procaine. It is less soluble than procaine 1 : 7 to 1 : 1.

For injection 0.1 and 0.05% solutions are recommended. For surface analgesia 2% solutions are advised (James, 1943). In Cazzaniga's case death occurred with a dose of 130 mg.; but James has used as much as 300 mg., although he advises that this amount should never be exceeded. The recommended therapeutic dose in man is 2 mg. per lb. of body weight. Hancock (1939) refers to a series of 2,000 cases of gastroscopy performed using 10 ml. of 2% solution without mishap.

For four months we have given anethaine, a 2% solution of amethocaine hydrochloride (4 ml. in conjunction with 1 ml. of 1 in 1,000 adrenaline), preparatory to bronchography and bronchoscopy. For the latter investigation the following was the procedure adopted.

The patient is given two "euphagin" tablets (benzocaine, 200 mg.) to suck forty minutes pre-operatively, followed ten minutes later with $\frac{1}{4}$ or $\frac{1}{2}$ gr. (11 or 16 mg.) of morphine. In the anaesthetic room the fauces, posterior pharyngeal wall, and both pyriform fossae are painted with the above solution, using about 2 ml., but only a portion of this amount would actually be expended. A final injection of 2 ml. between the cords under direct vision usually effects adequate analgesia for the examination, although a further supply of the local analgesic is available for direct bronchial spray if required.

Case Reports

Case 1.—The patient was a 62-year-old man with superior mediastinal obstruction believed to be secondary to a right upper lobe bronchial carcinoma. Local analgesia for bronchoscopy was being induced by the above method with 5 ml. of mixed solution. Within 30 seconds of the intratracheal injection the patient, who was sitting up, complained of giddiness and then fell back unconscious; convulsions followed as described above. Endotracheal oxygen and carbon dioxide was instituted immediately. The head was lowered and venepuncture performed, 500 ml. of blood being withdrawn. Recovery took place after three convulsive seizures. There were no sequelae.

Case 2.—A young man aged 21 with left lower lobe bronchiectasis was having preparation of the pharynx and larynx by this method, but 4 ml. of the solution instead of the usual 2 ml. was injected between the cords. (The first 2 ml. was thought to have been swallowed.) Within a few moments the patient became pale and a cold sweat broke out on the face; he lay down and at once started jactitating, at first in the lips and then in the limbs. The pulse was 80–90, regular, and of good volume. He then had a convulsion, during which he became cyanosed. The clonic spasm subsided and respirations returned, but were shallow and rapid. Almost immediately convulsions were renewed with increased intensity. The patient became deeply cyanosed, and during the clonic stage it was noticed that the pulse had ceased. Intracardiac adrenaline, 2 ml., was injected, while artificial respiration was begun and oxygen was administered through a face-mask. There being no response to these measures, the abdomen was opened through an upper midline incision and direct cardiac massage through the diaphragm was performed. A flaccid immobile heart was massaged without response for about one minute. A few rhythmic contractions occurred, ceased, and then in response to further massage recurred. The heart subsequently started and continued to beat strongly with occasional extrasystoles. Although the total period of cardiac arrest was estimated at more than four minutes and unconsciousness persisted for four days, complete recovery took place.

Comment

The first patient was seriously ill with gross congestion of the head, neck, and shoulder girdle. He proved difficult to anaesthetize and was uncooperative. Five millilitres of the normal mixture (80 mg.) of amethocaine was used and must have been rapidly absorbed. The second patient was a relatively fit young man. In this case 4 ml. (64 mg.) of amethocaine mixture was projected between the cords into the trachea. In an endeavour to co-operate the patient was hyperventilating, so that the full quantity of this drug may have directly reached the alveoli and been rapidly absorbed from this highly vascular surface.

Neither patient gave a history of allergy, and, although in each case more than the usual 4 ml. of amethocaine-adrenaline mixture was used, both patients would appear to have been unusually sensitive to the drug.

No satisfactory sensitivity test has been described, but Thomas and Fenton (1943) recommended the use of a patch skin test in suspected or doubtful cases. They admit that such a test is not entirely reliable. They also suggest that

sensitivity to the drug may be provoked by its previous use or the use of procaine, which is closely related to it.

Certainly in two fatal cases (Ahroon, 1946; Doane and Cohn, 1945) there had been recent previous administration of the drug. Thomas and Fenton draw attention to the intolerance to amethocaine manifested by allergic individuals, bronchial asthmatics, etc., as well as by cachectic patients.

Following the method of Hancock (1939), the technique in preparation for bronchoscopy has been modified. Each patient is now given a 1-gr. (65-mg.) amethocaine pastille to suck thirty minutes before instrumentation. This produces a much more complete analgesia of the mouth and pharynx than benzocaine, and it is considered that any idiosyncrasy possessed by the patient would be displayed while the relatively small dose contained in the pastille was being absorbed.

Apart from the severe constitutional reactions to amethocaine on record other less serious reactions have been recognized. These include attacks of dyspnoea and asthmatic symptoms (Fujikawa, Neves, Brasher, and Buckingham, 1948), blepharoconjunctivitis (McAlpine and Berens, 1942), and occupational dermatitis and eczema in doctors and oculists (Videbeck, 1936; Cameron, 1937; Hollander, 1939; Shimkin, 1939).

Prevention of Severe Reactions.—The following prophylactic measures are suggested: (a) A barbiturate should be given by mouth in pre-operative preparation—e.g., phenobarbitone 3 gr. (0.2 g.) or sodium amylal or nembutal 1½ gr. (0.1 g.) 40 minutes before examination. (b) An amethocaine pastille, 1 gr. (65 mg.), should be given to the patient to suck 30 minutes before the examination. (c) Amethocaine when used for surface analgesia should have adrenaline 1 in 1,000 solution added in the proportion of 4 to 1 by volume, and the total dose of amethocaine should not exceed 4 ml. of a 2% solution (80 mg.) or a corresponding amount in a more dilute solution. (d) Application by means of a laryngeal spray should be avoided, since its use can easily lead to overdosage. (e) Amethocaine should never be applied to an inflamed, traumatized, or highly vascular epithelial surface—especially so in the urethra (Knepper, 1937). (f) The drug should be avoided in patients giving a history of allergy or in severely debilitated or cachectic cases.

Control of the Established Reaction.—(i) General:—Secure an unobstructed airway by endotracheal tube if necessary. Begin artificial respiration with an oxygen-carbon-dioxide mixture. (ii) To control convulsions:—The immediate intravenous administration of a rapidly acting barbiturate—20 ml. of 2½% sodium thiopentone (0.5 g.)—is usually sufficient to control convulsions. (Picrotoxin should be available in case too much barbiturate is given.) (iii) Administration of respiratory and cardiac stimulants:—nikethamide, 2–5 ml.; caffeine sodium benzoate, 7½ gr. (0.5 g.), etc.

In using amethocaine in surface analgesia, as with many other local analgesics, from time to time the unusually susceptible individual may be met with. Reactions come with devastating suddenness and the results may be disastrous. By excluding those patients mentioned under (e) and (f) above, sensitive patients should be even more rarely met with. An analysis of the fatal cases reported reveals overdosage, faulty technique, or failure to use adrenaline to be more often the cause of death than individual idiosyncrasy.

Summary

A report is given of two toxic reactions with the use of amethocaine hydrochloride as an analgesic for bronchoscopy.

A review of the literature reveals twelve reported deaths at bronchoscopy and gastroscopy, while there are many cases of severe constitutional disturbance.

The advantages and disadvantages of the drug are assessed, the difficulty of applying an adequate sensitivity test accepted, and suggestions made for the prevention and treatment of toxic reactions.

I wish to express my thanks to Mr. O. S. Tubbs for permission to report these cases and for helpful advice in the preparation of this paper.

REFERENCES

- Ahroon, W. A. (1946). *Laryngoscope*, **56**, 320.
Cameron, O. J. (1937). *Nebraska med. J.*, **22**, 229.
Doane, J. C., and Cohn, E. M. (1945). *Anesthesiology*, **6**, 421.
Fasselt, H. (1939). *Arch. Ohr.-Nas.-KehlkHeilk.*, **146**, 217.
Freeman, E. B. (1939). *J. Amer. med. Ass.*, **112**, 217.
Fujikawa, F., Neves, A., Brasher, C. A., and Buckingham, W. W. (1948). *J. thorac. Surg.*, **17**, 123.
Hancock, P. E. T. (1939). *Proc. R. Soc. Med.*, **32**, 538.
Hollander, L. (1939). *Arch. Derm. Syph.*, **40**, 92.
Holt, T. (1945). *Arch. Otolaryng.*, **Chicago**, **42**, 280.
James, N. R. (1943). *Regional Anaesthesia for Intra-abdominal Surgery*, p. 2. Churchill, London.
Knepper, R. (1937). *Munch. med. Wschr.*, **84**, 1572.
McAlpine, P. T., and Berens, C. (1942). *Amer. J. Ophthalm.*, **25**, 206.
Pfeiffer, R. L. (1937). *Arch. Ophthalm.*, **Chicago**, **18**, 62.
Phillips, F. J., Congleton, V. L., and Tuttle, W. (1941). *Curr. Res. Anesth.*, **20**, 233.
Richards, R. (1947). *British Medical Journal*, **2**, 51.
Schindler, R. (1940). *Amer. J. digest. Dis.*, **7**, 293.
Schoen, F. (1939). *Wien. klin. Wschr.*, **52**, 505.
Shimkin, N. (1939). *Ann. Oculist.*, **Paris**, **176**, 198.
Shumacher, H. B. (1941). *Surgery*, **10**, 134.
Thomas, J. W., and Fenton, M. M. (1943). *J. Allergy*, **14**, 145.
Videbeck, H. I. (1936). *Ugeskr. Laeg.*, **98**, 278.

"GAMMEXANE" AND MOSQUITO CONTROL IN THE BELGIAN CONGO*

BY

G. DAVIDSON, B.Sc.

(From the Ross Institute of Tropical Hygiene)

Experiments are now being carried out in a large palm-oil plantation at Yaligimba, in the Belgian Congo, on the control of malaria by adult mosquito destruction, using the residual insecticide "gammexane" in the houses in the labour camps and in several native villages around the plantation. The plantation, which occupies an area of some 30 square miles (77.7 km.²), is situated in a forest belt in the northern part of the Belgian Congo. The main species of anopheline mosquito concerned is *A. moucheti*, which breeds at the edges of the large rivers in the vicinity. There are also a few *A. gambiae* and *A. paludis*.

Mosquito densities in the houses before and after treatment with the insecticide are being determined by the following four methods:

1. The number of mosquitoes resting in the houses during the daytime is being estimated by laying white sheets over the whole floor surface of a house and spraying the closed house with a pyrethrum-in-kerosene spray.
2. The number of mosquitoes leaving the house each morning is being determined by the use of window-traps.
3. The number of mosquitoes entering during the night to bite the occupants of any one bed in the house is being determined by the use of mosquito-net traps, which mosquitoes can enter but not leave.
4. The mosquito-nets of the inhabitants of the camps and villages are being examined to determine the percentage

*This work has been made possible by grants from United Africa Company, Limited, and from Imperial Chemical Industries, Limited.