attacks; prominent among these are headache and a desire to sleep, signs of injury, tongue-biting, incontinence of urine, vomiting, and soreness of muscles.

Then we can hear from the patient of the time of attacks, in particular their relation to sleep and to menstruation. As to sleep, the attacks may (1) occur during sleep, which is virtually diagnostic—of these the patient usually knows by his condition on waking; (2) if diurnal, show a marked predilection for the early morning hours; (3) occasionally occur just as the patient falls asleep. Further, an account of morning myoclonus is common and helpful.

Finally, though the patient cannot tell us of his actions in a state of automatism, he is often aware of this condition through circumstantial evidence, such as finding he has wandered into some strange place. The second part of the history, from relatives, may be decisive, but is often singularly inconclusive.

Aids to Diagnosis

The Electro-encephalogram.—Only if attacks are induced artificially, or if the patient is having frequent attacks under observation, is it possible to observe the distinctive seizure patterns. Confirmation of a clinical diagnosis, or help in a doubtful case, therefore depends on any variations from the normal which may be found in the resting record. Certain paroxysmal discharges seen in this record are virtually diagnostic of epilepsy: clearly defined “larval” attacks—i.e., electro-encephalographic but not clinical attacks—were never found by Williams (1944) in non-epileptic subjects, and the only case in 8,000 in which a wave-and-spike outburst (the type associated chiefly with petit mal) was seen in a non-epileptic was the identical twin of an epileptic. The Harvard workers found this type of outburst in only 1 in 500 of non-epileptic subjects but in 77% of patients with a clinical diagnosis of petit mal. These observers (Gibbs, Gibbs, and Lennox, 1943) conclude that in about 48% of persons with a history of seizures a routine electro-encephalogram is of great value in diagnosis, and in about 42% it is of little or no value. The results depend largely on the frequency of examinations. They are of most value in the diagnosis of petit mal, in which class of case clinical diagnosis is most often in doubt. It must also be recognized that not infrequently successive records in the same patient are inconsistent—a fact which has disturbed a number of clinicians (Holmes, 1946; Rook, 1947). The interpretation of records is a specialized and skilled procedure, and the results must be evaluated only in conjunction with the clinical picture.

Induction of Fits.—Two methods may be used to induce fits for the purpose of diagnosis. The first is the production of alkalosis by overbreathing, the second to cause a positive water balance by forced ingestion of water and the administration of pitressin (Blyth, 1943). A couple of minutes’ overbreathing may confirm a diagnosis of petit mal or bring out paroxysmal discharges in the electro-encephalogram. Otherwise these tests are formidable and have little place in ordinary practice.

The Clinical Decision.—Doubts and mistakes are inevitable even after the most painstaking analysis of the evidence. In civil practice, which is free from the complications in diagnosis encountered in the Services, the bias in a doubtful case should, in my view, be towards the diagnosis of epilepsy. If there is a history of recurrent attacks strongly suggestive of epilepsy, though certainty is lacking, the correct diagnosis is more likely to be epilepsy than anything else. Final judgment may be reserved, but appropriate treatment should be instituted.

[The Second Lecture, with the list of references, will appear in our next issue.]

ANTIHISTAMINE DRUG TREATMENT OF ACUTE NEPHRITIS

BY

JOHN CRAIG, M.B., Ch.B., F.R.C.P.Ed.

Professor of Child Health, Aberdeen University

N. S. CLARK, M.B., Ch.B., B.Sc., M.R.C.P., D.C.H.

Lecturer in Child Health, Aberdeen University

AND

J. D. CHALMERS, M.B., Ch.B.

Supernumerary House-Physician, Royal Aberdeen Hospital for Sick Children

This paper is a preliminary report on the use of “anthisan” in the treatment of acute nephritis in children. The number of cases so far treated is small, but the results have been so striking that we feel it is advisable to publish them at this stage in order to stimulate further work on the subject.

No specific treatment has yet been found which will abort acute nephritis or even appreciably modify its course. Treatment up till now has been directed towards providing rest both for the patient and, so far as this can be achieved by dietary restrictions, for his kidneys, in the hope that these organs have not already been irreparably damaged and will eventually recover spontaneously. If the response we have observed to treatment with anthisan can be confirmed over a larger series of cases a very material advance will have been made in the treatment of acute nephritis. In a centre such as this it will be some years before we see enough cases for the proper evaluation of the treatment with adequate controls. Therefore it is advisable that this work should also be undertaken at other and larger centres.

The well-known tendency for acute nephritis to occur at an interval of a week or two after an acute bacterial infection suggests that nephritis may be in the nature of an allergic response to bacterial toxins. The relationship which seems to exist between acute nephritis and some cases of anaphylactoid purpura lends further support to this theory. It would therefore appear reasonable to test the effect of the antihistamine drugs in this condition. So far as we are aware the only published work on this subject is that of Reubi (1946), who has claimed good results from treatment with “antistin” both in a few cases of experimentally induced nephritis in rabbits and in a small series of cases of acute nephritis in human beings. As might be expected, he found that antistin had no effect on chronic nephritis where the kidneys are already irreparably damaged.

We have now used anthisan in the treatment of eight successive cases of acute nephritis admitted to the Royal Aberdeen Hospital for Sick Children. The course of these cases has been compared with that of the nine consecutive cases of this condition admitted immediately preceding the institution of this treatment. Two cases of nephritis which were nephrotic in type from the outset have not been included, but, apart from them, all fresh cases of nephritis admitted to the hospital between October, 1946, and October, 1948, are included in either the anthisan or the control groups.

The groups differ in one important respect—i.e., they were not admitted during the same period of time—and it is of course possible that the later cases were suffering from a relatively benign form of nephritis. The two groups are, however, closely comparable as regards the duration of illness before admission and the clinical assessment of severity at the onset, based on albuminuria, haematuria,

oedema, hypertension, blood urea, and urinary output. The average age of the control group was 9 years and of the anthesis group 6 years.

Four cases in the control group and two in the anthesis group showed signs of active infection in the throat or ears on admission; the same number of cases in each group showed evidence of recurrent or persistent throat infection during the course of treatment. Whether or not these persistent infections had any effect on the course of the disease it is difficult to say. All patients in both groups who showed signs of active infection were given courses of penicillin or sulphonamide, but this did not seem to influence materially the course of the nephritis. Hypertensive convulsions, which occurred in one control and two anthesis cases, were treated on the usually accepted lines by lumbar puncture, hypertonic magnesium sulphate enemata, and sedatives. No other treatment was employed in either group apart from complete rest in bed until abnormal constituents had remained absent from the urine for several weeks and restriction of protein intake as long as azotaemia persisted.

For the purposes of comparison the two groups have been subdivided into "mild" and "moderate or severe" cases. Cases were classified as "mild" if there was no hypertension or oedema other than slight puffiness around the eyes and if azotaemia was absent or minimal. On this basis three cases in the control group and two cases in the anthesis group were mild, leaving six cases in each group which were moderate or severe.

Brief summaries of the important clinical and laboratory findings and of the course of all cases in both groups are given. The figure for "duration" shown at the end of each case history signifies the duration of activity of the nephritic process from the first day of treatment until the last abnormal signs had disappeared. Albuminuria or microscopical haematuria were the most persistent abnormal signs in all cases except No. 16, in which the blood urea remained raised after the urine had returned to normal.

Case Histories

Control Group: Mild Cases

Case 1.—A boy aged 11 was admitted on Feb. 28, 1948, with a six-day history of headache, giddiness, and haematuria. He had had a similar but milder attack in February, 1946. Since then, however, he had been very well, and no albumin had been found in his urine. Examination showed no oedema apart from slight puffiness around his eyes. The throat was healthy. B.P. 110/65. No other abnormality was detected. Urine: smoky, albumin ++, output normal. Blood urea 32 mg. per 100 ml. Sixteen days later his urine was free from albumin, and no red blood cells were seen on microscopical examination. Only once since then has his urine contained any abnormal constituents—namely, a few granular casts. He was discharged on April 3, and was very well when seen at the follow-up clinic two months later. *Duration:* 16 days.

Case 2.—A girl aged 10 was admitted on Feb. 6, 1948, with a history of having developed a sore throat ten days previously. Seven days later she had persistent headache, and on the day before admission her B.P. was 140/90 and her urine contained albumin ++ and scanty red blood cells and pus cells. Examination showed no oedema; the throat was healthy; there were signs of a congenital cardiac defect—probably a patent ductus arteriosus. B.P. 120/80. Blood urea 29 mg. per 100 ml. Urine: no albumin, scanty pus cells, culture sterile. Next day her urine again contained albumin, and this persisted for seven days. At no time during her stay in hospital were casts or red blood cells noted. She was discharged on Feb. 27. *Duration:* 8 days.

Case 3.—A girl aged 8 was admitted on June 19, 1947, with a history of haematuria for two days. There was no history of a throat infection, but she had been vaguely unwell for a

week before admission. Examination showed no oedema: tonsils were mildly inflamed; physical examination otherwise negative. B.P. 115/85. Blood urea 58 mg. per 100 ml. Urine: macroscopic blood, albumin + + +, occasional granular casts. Her urine became free from albumin 26 days after admission, but a few red blood cells were present microscopically until the 39th day. Tonsillectomy was performed on Aug. 8 under cover of a course of penicillin, and she was discharged on Aug. 14. *Duration:* 39 days.

Anthesis Group: Mild Cases

Case 4.—A girl aged 5 was admitted on June 13, 1948, with a history of haematuria and puffiness of her eyelids for two days. She had had an attack of otitis media two months previously, and this had been followed by haematuria, but her symptoms had cleared in ten days, and she had been back at school for two weeks. Examination showed slight puffiness below the eyes. Tonsils were moderately enlarged but not inflamed. No other abnormality was detected. B.P. 132/70. Blood urea 50 mg. per 100 ml. Urine: macroscopic blood, albumin + +.

Anthesis, 0.1 g. t.i.d., was given from June 13 to June 21 inclusive. By the latter date her symptoms had completely subsided and her urine was free from albumin and red blood cells. She was discharged on June 29 to rest in bed at home for one month. She returned to the follow-up clinic on Aug. 2, having just recovered from an attack of mumps. Her urine showed a trace of albumin, but this had disappeared by her next visit to the clinic on Aug. 30, and she has had no albuminuria since. *Duration:* 8 days.

Case 5.—A boy aged 2 was admitted on Aug. 30, 1948, with a history of listlessness, pallor, and puffiness of the eyelids for two days. There was no history of an upper respiratory infection or of haematuria. On examination he was a pale, listless child with puffy eyelids but no oedema elsewhere. Apart from a soft apical systolic murmur and some enlargement of the tonsils, no other abnormality was detected. B.P. 106/76. Blood urea 20 mg. per 100 ml. Urine: albumin +, scanty red blood cells. Haemoglobin 54%.

Anthesis, 0.1 g. t.i.d., was given from Aug. 30 to Sept. 3 inclusive, and within four days his urine was free from albumin and red blood cells. On Sept. 4 albumin was again present in his urine, but thereafter no abnormality was detected. He was discharged on Sept. 15, and has since been seen twice at the follow-up clinic, his urine being normal on both occasions. This attack of nephritis was accompanied by a fairly severe microcytic hypochromic anaemia, which responded to iron therapy. *Duration:* 6 days.

Control Group: Moderate or Severe Cases

Case 6.—A girl aged 4 was admitted on June 2, 1948. Two weeks before admission she developed a respiratory infection which responded to treatment with sulphathiazole; her right ear then became painful and started to discharge. Two days before admission her urine became dark brown in colour. Examination showed no oedema apart from some puffiness around her eyes. Her throat was healthy, but her right tympanic membrane showed a loss of lustre. An x-ray film of her right mastoid showed a loss of translucency, suggesting a mastoiditis. No other abnormality was found on physical examination. B.P. 125/80. Blood urea 66 mg. per 100 ml. Urine: albumin + + +, no cellular deposit, output 7 oz. (197 ml.) in first 24 hours. On June 7 gross haematuria was present and there was a purulent discharge from the right ear. The discharge from her ear gradually dried up, her urinary output increased, and her haematuria disappeared. She was discharged on June 22, with a trace of albumin still present in the urine. At the follow-up clinic two months later the albuminuria had disappeared. B.P. 110/70. *Duration:* over 20 days.

Case 7.—A girl aged 11 was admitted on Dec. 8, 1947. Four weeks before admission she developed a sore throat and a generalized scarlatiniform rash. She returned to school after two weeks. On the day before admission her eyelids became puffy and she passed a small quantity of dark-brown urine. On examination there was some puffiness of the lower eyelids.

Her throat was healthy. A few rales were present at her left lung base. No other abnormality was detected. B.P. 176/80. Blood urea 51 mg. per 100 ml. Urine : albumin + + +, macroscopic blood. On Dec. 21 the blood urea was 38 mg. per 100 ml., B.P. 110/65; albuminuria was still marked, but no red blood cells were present. The albumin persisted until Feb. 26, 1948, when for the first time the urine showed no abnormality. She was discharged on March 4. *Duration* : 80 days.

Case 8.—A girl aged 4 was admitted on March 20, 1947, with a history of swelling of her face and abdomen and of passing dark-coloured urine for three days. On examination oedema of the face and limbs was present and the throat was inflamed. Physical examination was otherwise negative. B.P. 102/65. Blood urea 156 mg. per 100 ml. Urine : albumin + + +, macroscopic blood, output 8 oz. (227 ml.) in first 24 hours. By April 1 the oedema had practically disappeared; the blood urea was 82 mg. per 100 ml. and frank haematuria had ceased. Her blood urea on April 27 was 23 mg. per 100 ml. An acute tonsillitis on May 3 caused an increase in the haematuria and albuminuria. The red blood cells gradually diminished, and on May 19 the urine was free from albumin. A few red blood cells, average 2 per high-power field, persisted till the date of her discharge on July 2. These were still present at her first visit to the follow-up clinic on July 24, but had disappeared by Sept. 4. Her urine has been normal on several occasions since that date. *Duration* : over 126 days.

Case 9.—A girl aged 10 was admitted on Oct. 2, 1946. Three weeks before admission she developed swelling of the eyelids and face. A week later haematuria was first noticed; this gradually became more marked, being accompanied by oliguria. On examination no oedema was present and her throat was healthy. B.P. 155/95. No other abnormality was detected. Blood urea 52 mg. per 100 ml. Urine : albumin + +, macroscopic blood, numerous granular casts, output 15 oz. (425 ml.) in first 24 hours. Blood urea on Oct. 19 was 28 mg. per 100 ml. The albuminuria and haematuria persisted, and the blood pressure was 140/90 on Dec. 15. On March 14, 1947, for the first time red blood cells were not seen on microscopical examination of the urine, but the albuminuria persisted. B.P. 120/70. A trace of albumin was still present in the urine when she was discharged on June 22, and again on her return to the follow-up clinic on July 16. *Duration* : over 287 days.

Case 10.—A boy aged 9 was admitted on Nov. 16, 1946. Three weeks before admission he developed a sore throat, followed a week later by a purulent discharge from his left ear. Five days before admission swelling of his eyelids was noted, and three days later there was haematuria and oliguria. On examination oedema of his face and ankles was present, tonsils infected, left suppurative otitis media. No other abnormality was detected. B.P. 140/100. Blood urea 221 mg. per 100 ml. Urine : albumin + + +, macroscopic blood, numerous granular casts, output 14 oz. (397 ml.) in first 24 hours. On Dec. 4 he developed chicken-pox and was transferred to an isolation hospital. At this time his blood urea was 50 mg. per 100 ml. and his B.P. 126/74. Frank haematuria persisted and was present on his readmission to this hospital on Dec. 27. A flare-up of his chronic tonsillar sepsis caused a further increase in the haematuria, and on Feb. 3, 1947, tonsillectomy was performed under cover of a course of penicillin. Gradual improvement took place in spite of an attack of measles starting on March 5. Red blood cells were last seen in the urine on April 28, but traces of albumin persisted until May 25. He was discharged on June 7, and until his last visit to the follow-up clinic on Dec. 24, 1947, his urine remained normal. *Duration* : 190 days.

Case 11.—A girl aged 11 was admitted on Dec. 31, 1946, having developed a sore throat three weeks previously. Four days before admission her face became swollen, and two days later her urine became dark brown in colour and she had several convulsions. There were four further convulsions on the day of her admission. On examination she was drowsy and showed marked generalized oedema; the throat was healthy. No other abnormality was detected. B.P. 170/120. Blood urea 56 mg. per 100 ml. Her urinary output was not diminished.

Urine : albumin + +, macroscopic blood. She had no further fits after admission, and by Jan. 12, 1947, her blood pressure had fallen to 120/80 : the urine still contained albumin and red blood cells, but there was no longer obvious haematuria. Albumin and red blood cells were persistently present in the urine till March 8. Thereafter scanty red blood cells and traces of albumin were occasionally present. She was discharged on April 14. *Duration* : 67 days.

Anthisan Group: Moderate or Severe Cases

Case 12.—A girl aged 5 was admitted on June 21, 1948, with a history of having developed a sore throat two weeks previously. On the night before admission her face became puffy and she passed dark urine. On examination there was definite oedema of her face and slight oedema of her legs : follicular tonsillitis was present with enlarged tonsillar nodes. Physical examination was otherwise negative. B.P. 154/100. Blood urea 58 mg. per 100 ml. Urine : albumin + +, macroscopic blood, output in first 24 hours 12 oz. (340 ml.). A throat swab showed haemolytic streptococci.

Anthisan, 0.1 g. t.i.d., was given from June 21 to July 6. The oedema disappeared and the blood pressure rapidly returned to within normal limits. On July 1 her urine was free from albumin and no red blood cells or casts were seen on microscopic examination. She was discharged on July 6, and at her subsequent visits to the follow-up clinic on Aug. 16 and Sept. 28 she remained well and her urine was normal. *Duration* : 10 days.

Case 13.—A boy aged 9 was admitted on July 15, 1948, with a history of having developed a sore throat two weeks previously. Eleven days later he complained of headache and sickness, and his urine was dark brown in colour. Two days before admission his face became swollen and his urinary output was considerably diminished. On examination he was drowsy, and oedema of his face and sacrum was noted; his tonsils were enlarged and inflamed and the tonsillar nodes were palpable; no other abnormality was detected. B.P. 130/108. Blood urea 138 mg. per 100 ml. Urine : albumin + +, 20 red blood cells per high-power field (uncentrifuged specimen), output 13 oz. (368 ml.) in first 24 hours.

He was given anthisan, 0.1 g. four-hourly, from July 15 to 23. His oliguria persisted, and on July 18 his blood pressure rose to 158/120 and he developed a series of convulsions. He had another short convulsion on July 20, but next day his urinary output increased and his blood pressure began to drop. He had no further convulsions, but still complained of headache and vomited occasionally. By July 28 he was symptom-free, his blood urea was 41 mg. per 100 ml., his blood pressure 130/75, and his urine showed no abnormal constituents. He was discharged on Aug. 10. *Duration* : 13 days.

Case 14.—A boy aged 7 was admitted on Sept. 29, 1948, with a history of puffiness of his face and haematuria for three days. Three weeks previously he had complained of a sore throat accompanied by earache. On examination oedema of his face and ankles was present; his throat and ears were healthy and no other abnormality was found. B.P. 150/116. Blood urea 25 mg. per 100 ml. Urine : albumin + +; macroscopic blood.

Anthisan, 0.1 g. t.i.d., was given from Sept. 29 to Nov. 11. On Oct. 6 his urine was still smoky and contained albumin, but his oedema had subsided and his B.P. was 110/70. On Oct. 20 the albuminuria had cleared up and no red blood cells were present on microscopical examination. Routine examinations of his urine since Oct. 20 have on only two occasions shown an abnormality. Once a trace of albumin was noted, and five days later 6 red blood cells per high-power field were found on microscopical examination. *Duration* : 21 days.

Case 15.—A boy aged 5 was admitted on Oct. 21, 1948, with a history of swelling of his face for six days. Three days before admission the swelling became more pronounced and was accompanied by headache and vomiting. There was no history of sore throat or haematuria. On examination his face was oedematous, his throat was infected, and his tonsillar nodes enlarged; no other abnormality was found. B.P. 150/110. Blood urea 45 mg. per 100 ml. Urine : albumin + +,

macroscopic blood, occasional red cell and granular casts. His urinary output was normal.

He was given anthisan, 0.1 g. t.i.d., from Oct. 21 to Nov. 12. On Oct. 23 his urine contained no microscopical blood, but albumin was still present: the facial oedema had almost completely subsided. On Nov. 3 his urine showed no abnormality, and his blood pressure was 105/70. On only one occasion since Nov. 3 has a trace of albumin been found in his urine. *Duration*: 13 days.

Case 16.—A girl aged 6 was admitted on Oct. 22, 1948, with a history of swelling of her face and abdomen and some diminution in her urinary output for seven days. She had recently been a patient in the ear, nose, and throat ward of this hospital, suffering from otitis media, and had been discharged only seven days before her present admission. On examination there was some oedema of her face; her throat and ears were healthy. No other abnormality was noted. B.P. 185/145. Blood-urea 60 mg. per 100 ml. Urine: albumin +, no cellular deposit, output 9 oz. (255 ml.) in first 24 hours.

She was given anthisan, 0.1 g. t.i.d., from Oct. 22. On Oct. 30 the blood pressure was 95/55; examination of her urine showed no abnormality, but her blood urea was still 60 mg. per 100 ml. On Nov. 12 the blood urea was 39 mg. per 100 ml. At the time of writing (Nov. 17) she is still having anthisan, and her urine has remained normal on daily examination. *Duration*: 20 days.

Case 17.—A girl aged 9 was admitted on July 5, 1948, with a history of vomiting and abdominal pain for three days. Physical examination on admission was negative, but her urine was found to contain albumin and a few pus cells. Three days after admission she complained of headache, and there was a suggestion of swelling of her face. Her blood pressure was 140/98. On July 10 she developed convulsions. B.P. 182/130. Blood urea 43 mg. per 100 ml. Her urinary output was 16 oz. (454 ml.) in 24 hours, and the urine continued to show albumin and a few pus cells with occasionally a few red blood cells.

Anthisan, 0.1 g. t.i.d., was given from July 11 to Aug. 3. On July 27 her blood pressure was 90/60 and her urine contained no abnormal constituents. Her urine remained normal until Aug. 27, when albumin and red blood cells were again found. Anthisan was again given in dosage of 0.1 g. t.i.d. from Aug. 27 to 31 and 0.1 g. four-hourly from Sept. 1 to 13. From Sept. 6 till her discharge on Oct. 29 no abnormality was found on daily examination of her urine, and her blood pressure and blood urea remained normal. *Duration of initial attack*: 16 days. *Duration of relapse*: 10 days.

Comment

After the first one or two cases had been treated with anthisan the clinical impression was formed that these cases were clearing up more quickly than would have been expected, and this impression became stronger as more cases were treated. A study of the figures given for the duration of the signs of activity in each case confirms this clinical impression, for not only is the average duration in the anthisan group (13 days) much less than that in the

control group (92 days) but if the mild and the moderate or severe cases are considered separately the least satisfactory case in the anthisan group has cleared up as quickly as the most satisfactory in the control group (see accompanying Table).

After the conclusion of anthisan treatment urine examination was made daily in every case up to the date of discharge from hospital, and occasional examination at the follow-up clinic thereafter.

In Case 17 the patient had a recurrence of albuminuria and microscopical haematuria after her urine had remained normal for one month. She was given another course of anthisan, and her urine again cleared after ten days; it has now remained normal for over two months.

In two other anthisan cases a trace of albumin was present on one occasion, and in one of these scanty red blood cells were seen on another occasion. All other urine examinations were normal.

It is of course too early as yet to speak of complete cure in any of the cases in the anthisan group, and all will be followed up so far as is possible. It seems probable, however, that as a general rule the longer signs of activity persist the greater are the chances of permanent renal damage. There would appear, therefore, to be reasonable grounds for hope that the further progress of the cases in the anthisan group will prove equally satisfactory.

Anthisan did not prevent the development of hypertensive convulsions in Case 13, in which they developed two days after the beginning of treatment. The patient's subsequent progress was, however, unusually satisfactory for such a severe case.

We have so far formed no definite opinions on optimum dosage or duration of treatment. In some cases the administration of anthisan was stopped when all abnormal signs disappeared; in others it was continued for a further week or two.

Acute nephritis is a disease which may run a very variable course, and with such a small series of cases it would obviously be unjustifiable and premature to advance any definite claims with respect to the efficacy of a particular line of treatment. Much further work will be necessary before the place of anthisan in the treatment of acute nephritis can be accurately assessed, and it is obviously desirable that this work should go forward as speedily as possible. Treatment with anthisan is simple, and toxic manifestations with this drug seem to be few and relatively unimportant. If its value in acute nephritis can be proved a considerable advance in the management of this disease will have been achieved.

Summary

An account is given of the treatment with anthisan of eight cases of acute nephritis in children.

The course of these cases is compared with that of the nine consecutive cases of this condition seen in the same hospital during the period immediately preceding the institution of this treatment.

The duration of activity of the disease, as assessed by abnormal findings in the urine, is notably shorter in the cases treated with anthisan than in those of the control group.

REFERENCE

Reubi, F. (1946). *Helv. med. Acta*, **13**, Suppl. 18, 1.

Table Showing Duration of Activity of the Nephritis in Each Case

	Control Group		Anthisan Group	
	Case No.	Duration	Case No.	Duration
Mild cases	1	16 days	4	8 days
	2	8 "	5	6 "
	3	39 "		
	Average 21 days		Average 7 days	
Moderate or severe cases	6	Over 20 days	12	10 days
	7	80 "	13	13 "
	8	Over 126 "	14	21 "
	9	Over 287 "	15	13 "
	10	190 "	16	20 "
	11	67 "	17	16 "
	Average 128 days		Average 15 days	
Average duration of whole group		92 days	13 days	

A recommendation for the building of a health centre, at an estimated cost of £38,000, has been passed by the Birmingham City Council. This first health centre will be experimental and will serve a population of about 20,000. There is to be accommodation for five general practitioners. Eventually it is hoped that there will be between 40 and 50 health centres in the city, each serving about 15,000 people.