single daily injection. On the 27th day slight morning nausea appeared and four days later she began to vomit. Benadryl, 50 mg. eight-hourly, was started on the following day. During the next twelve days there was no nausea and she vomited only once. Identical capsules containing phenobarbitone were substituted. Twenty-four hours later the vomiting reappeared. The next day the same capsules containing 100 mg. of antistim were substituted; vomiting ceased after the first dose. Antistim in tablet form was continued for ten days. After its withdrawal neither nausea nor vomiting reappeared and she remained free from these symptoms throughout the remainder of the 32 weeks of streptomycin treatment.

In at least three of these cases we would have had to stop the streptomycin if we had not been able to control the nausea and vomiting. With the continuous administration of benadryl it was possible to complete the full four-months course. We would emphasize, however, that giddiness due to streptomycin was not relieved by the antihistamine drugs.

Summary

Antihistamine drugs were found to relieve the nausea and vomiting which sometimes occur during treatment with streptomycin. This conclusion was supported by control tests in four severe cases. Giddiness due to streptomycin was not relieved.

We are grateful for the co-operation of the physicians of the Brompton Hospital under whose care the patients were admitted. We also wish to thank the hospital pharmacists for help with the control tests and Messrs. Parke, Davis and Co. for a supply of empty capsules.

REFERENCES

- Committee on Chemotherapeutics, National Research Council (1946). *J. Amer. med. Ass.*, **132**, 70.
Farrington, R. F., Hull-Smith, H., Bunn, P. A., and McDermott, W. (1947). *Ibid.*, **134**, 679.
Heilman, D. H., Heilman, F. R., Hinshaw, H. C., Nichols, D. R., and Herrell, W. E. (1945). *Amer. J. med. Sci.*, **210**, 576.
Medical Research Council (1948). *British Medical Journal*, **2**, 769.
Nichols, D. R., and Herrell, W. E. (1946). *J. Amer. med. Ass.*, **132**, 200.
Streptomycin Committee of Veterans Administration (1947). *Amer. Rev. Tuberc.*, **56**, 485.

TETANY FOLLOWING REMOVAL OF A PARATHYROID ADENOMA WITH BONE DISEASE: FINALLY ALLEVIATED WITH CALCIFEROL

BY

NANCY S. CONWAY, M.B., Ch.B.

Hall Tutorial Fellow in Medicine, University of Glasgow
(From the Gardiner Institute of Medicine, University of Glasgow, and the Western Infirmary, Glasgow)

When a parathyroid adenoma has caused extensive bone disease its removal is followed by severe tetany. This report illustrates the value of calciferol and the uselessness of parathormone in the treatment of such a case.

Case History

The patient, a woman aged 43, was admitted on June 1, 1947, with backache dating from 1940 and general weakness which had caused her to become bedridden. She had kyphosis and sacral deformity, and radiographs revealed extreme decalcification of the entire skeleton. The blood calcium was 12.2 mg. per 100 ml., phosphorus 3 mg. per 100 ml.; alkaline phosphatase 45 King-Armstrong units per 100 ml., and urea 32 mg. per 100 ml.

A diagnosis of hyperparathyroidism was made, and on July 15 Professor C. F. W. Illingworth removed a small adenoma from one parathyroid gland.

After operation the serum calcium fell abruptly (Fig. 1) and tetany developed. Parathormone was started on the day after operation, 80 units being given intramuscularly. The dose was

increased until 200 units were given daily, but no rise in the serum calcium occurred even after the initial doses. During this period the patient received intravenous calcium gluconate (10-20 ml. of a 20% solution two or three times a day) and 12 g. of calcium lactate orally per day. In order to facilitate

FIG. 1.—Showing the immediate fall in the calcium and phosphate and rise in the phosphatase which occurred after operation. During administration of calciferol the calcium and phosphate gradually rose, and the phosphatase fell, to normal.

absorption of calcium a low phosphorus diet (0.56 g.) and 40,000 units of calciferol were given daily. In spite of these measures the patient's condition rapidly deteriorated. In addition, she became uncooperative, refusing to have injections, and parathormone was stopped on the 14th post-operative day, when the serum calcium was still less than 5 mg. per 100 ml. As a last resort a transfusion of 2½ pints (1.42 litres) of fresh citrated

FIG. 2.—Showing rise of blood urea after operation and during calciferol treatment. It fell again after cessation of the drug.

blood (Group A) was given. The result was disastrous. Four hours later tetany, more severe than had hitherto occurred, developed, although examination of the blood showed that there had been no further drop in calcium.

An attempt was then made to raise the serum calcium by increasing the daily dose of calciferol to 300,000 units (6 ostelin tablets, each of 50,000 units). Three days later definite improvement was noted; thereafter the serum calcium gradually rose (Fig. 1), and signs of tetany disappeared and were not again elicited. At no time did the patient show any intolerance to the calciferol, which was stopped on Oct. 22 after a total of 19,000,000 units had been given. The blood urea, however, which rose after operation, remained high until some time after the drug had been stopped (Fig. 2). The patient was discharged

TABLE I.—The Serum Calcium, Phosphate, and Alkaline Phosphatase after Cessation of Calciferol Treatment

Date	Serum Calcium (mg./100 ml.)	Serum Phosphate (mg./100 ml.)	Serum Alkaline Phosphatase (King-Armstrong Units per 100 ml.)
30/10/47 ..	9.1	4.0	10.2
21/11/47 ..	8.6	3.8	9.0
19/12/47 ..	8.5	—	9.0
16/1/48 ..	10.2	3.5	8.6
24/2/48 ..	8.8	3.1	—
10/3/48 ..	9.0	3.2	8.6

on Oct. 31, and since then the blood chemistry has remained satisfactory (Table I). In March, 1948, she was symptom-free, had gained 30 lb. (13.6 kg.) in weight, and was able to walk unsupported.

Discussion

The post-operative tetany was due to an abrupt fall in serum calcium. Efforts to raise the latter by parathormone (total 1,260 units) were unsuccessful even with different batches of the extract. That it was physiologically active was certain, since its effect in producing a diuresis was well marked. It was felt that failure to absorb calcium might account for the lack of response to the hormone, but faecal analysis for calcium showed that this was not so. It was concluded that the fall in serum calcium and the resultant tetany were due not to too little parathyroid tissue but to the "hungry bones" (Cope, 1941). After operation the decalcified skeleton absorbs calcium and phosphorus from the plasma with so much avidity that hypocalcaemia and hypophosphataemia result. The biochemical findings support this theory, the blood calcium and phosphorus both falling immediately after operation. The rise in alkaline phosphatase which occurred is interpreted as being due to increased osteoblastic activity associated with this recalcification.

The severe and almost fatal tetany which followed the transfusion of blood was produced in a different way. The introduction of citrate (the anticoagulant) would on account of its avidity for calcium ions cause a further reduction in the ionized calcium without altering the total calcium. "Citrate tetany" has been produced experimentally, but its occurrence after transfusion must indeed be rare unless, as was the case here, the serum calcium is already depleted.

The rise in serum calcium which started three days after large doses of calciferol had been prescribed was due to the specific effect of the drug and not to a spontaneous recovery, since the calcium fell again when the dosage of calciferol was temporarily reduced (Fig. 1). During its administration, also, the serum phosphorus rose slowly, while the alkaline phosphatase fell with remarkable rapidity and was normal eight weeks after operation. The calciferol did not appear to have any inhibitory effect on deposition of calcium or phosphorus in bone, for during a five-weeks period 69.25 g. of calcium and 14.84 g. of phosphorus were retained.

Not every case of parathyroid adenoma will require massive calciferol therapy in the post-operative phase. It will definitely be indicated when there is well-marked bone disease and should be started immediately after removal of the adenoma. In addition, large doses of calcium must of course be given orally, and until the calciferol takes effect symptoms of tetany may be alleviated by intravenous calcium gluconate; parathormone should not be used.

I wish to thank Professor C. F. W. Illingworth for allowing me to investigate this patient and Professors J. Norman Davidson and J. W. McNee for advice in the preparation of the paper.

REFERENCE

Cope, O. (1941). *Ann. Surg.*, **114**, 706.

The Ministry of Health has issued a pamphlet, "Nursing and Domestic Staff in Hospitals," intended as a guide to hospital management committees on the recruitment, training, and conditions of service of nurses. It includes an example of how to work a shift system in a ward of 34 beds with a staff of one sister, two staff nurses, seven students for day duty, and two nurses for night duty, as well as some observations on what the proper task of a nurse should be and how some of the ward duties at present performed by nurses could be undertaken by domestic orderlies. These include arranging flowers, answering telephones and running messages, preparing meal trays, and handling clothing and laundry. The pamphlet has been sent to boards of governors of teaching hospitals and hospital management committees.

AGGLUTININ ANTI-P IN PREGNANCY REPORT ON TWO CASES

BY

I. DUNSFORD

(From the National Blood Transfusion Service, Sheffield)

The detection of an antibody in the serum during pregnancy has had much significance since Levine and his co-workers (1941) first propounded the theory (since amply confirmed) that haemolytic disease of the newborn is caused by the action on the foetal erythrocytes of immune antibodies formed in the mother's serum during pregnancy and passed through the placenta into the foetal circulation. The formation of these antibodies is stimulated by an antigen present in the foetal red cells but absent from the maternal red cells, the antigen having been inherited from the father. In a very large proportion of cases it has been demonstrated (Boorman *et al.*, 1942) that the rhesus complex has been involved, and in a few the ABO system (Aubert *et al.*, 1945).

There are not many references in the literature to immune or naturally occurring anti-P, and a leading article in the *British Medical Journal* of Oct. 20, 1945 (p. 535), refers to a case in which anti-P was associated with pregnancy. Wintrobe (1946) states: "In rare cases iso-immunization apparently occurs against the agglutinogens A and B, and possibly even M and P may cause immune bodies to be formed." Nigg (1930) has described two human sera which were later found to contain natural anti-P iso-agglutinins, whilst Wiener (1943) records having found only three or four such sera during a period of ten years. The latter serologist also found this iso-agglutinin in two post-transfusion sera he investigated; he states that "while the possibility exists that this factor could on rare occasions give rise to haemolytic transfusion reactions, to date no such reactions have been reported."

During the course of routine investigations carried out in this laboratory, which included testing the patient's serum at 37° C. against a series of Group O cells chosen not only for their rhesus type but also for their M, N, P, Kellecher, Lewis, and Lutheran types, the agglutinin anti-P was discovered in the sera of two pregnant women. In view of the extreme rarity of such an occurrence the history and the serological findings in these two cases are presented below.

Case 1

Mrs. X, a 2-gravida aged 36, in 1940 had a full-term clinically normal child weighing 6 lb. 12 oz. (3.06 kg.), and was again due for delivery on July 31, 1947. There was no history of transfusion or injection of blood. A blood sample was sent from the antenatal clinic for rhesus typing.

Serological Findings

March 7.—Mrs. X:—Group O Rh-positive. Her serum agglutinated some Group O cells weakly at 37° C., giving much stronger reactions at room temperature and at 4° C. At the latter temperature it appeared specific for P cells with a titre of 4. These reactions were compared with those of a proved anti-P serum and were found to be identical, suggesting that Mrs. X's serum contained anti-P. The serum gave a strong Diamond (Diamond and Abelson, 1945) slide reaction. A sample of serum was submitted to the Blood Group Reference Laboratory, Lister Institute, and confirmed as anti-P.

May 9.—Mrs. X:—Group O; genotype cDE/cde; M, N, p. The titre of the antibody against the same Group O p cell was 16 at room temperature and there was no evidence of enhancing when pooled serum from Group O p persons was used in place of saline. Mr. X:—Group A Rh-positive; genotype CDe/cDE; M, N, P. His cells were strongly agglutinated by his wife's serum at 4° C. and room temperature, and weakly at 37° C after absorption with A p cells.

June 20.—Mrs. X:—The titre of the antibody was now 4 with the standard O p cell. There was no evidence of an