The Puerperium.—The patients are allowed out of bed on the seventh day and normal cases are discharged on the tenth day. Active movement and breathing exercises are encouraged from the very beginning. Each ward is visited daily by a physiotherapist, who conducts the exercises. During a period of eight years, with 12,830 deliveries, there have been only two deaths from pulmonary embolism.

Discussion

In addition to the series here presented there were from 1944 to 1946 in the General Hospital three separate series, of 1,050, 1,346, and 1,730 deliveries, without a maternal death, and it must be stressed that such results are the outcome of a co-operative effort on the part of the whole staff of both parts of the Municipal Service and of the general practitioners in the area.

Particularly must I lay emphasis on the loyalty and diligence of the resident and nursing staff, who have given service far beyond the power of gold to buy. The excellence of the general administrative arrangements and the ancillary services provided have contributed greatly. Other factors of importance are the large proportion of antenatal beds and the continuity of specialist supervision for abnormal cases.

Of the therapeutic agents, blood, plasma, penicillin, and the sulphonamides have played a great part. It is important to give blood in adequate amount; for example, one greatly exsanguinated woman received 10 pints (5.68 litres), and at one stage two veins were being used at the same time to give the blood under pressure of oxygen. Such treatment has been made possible by the wonderful facilities provided by the Leeds Regional Blood Transfusion Service. Glucose 20% solution, too, has played its part, and Dieckmann (1942) has stated that the proper use of hypertonic glucose solution in pregnancy is one of the outstanding accomplishments of modern obstetrics. In third-stage and post-partum haemorrhage, ergometrine, given intravenously, has proved life-saving and deserves to rank in importance in obstetrics with penicillin.

These potent remedies together with modern methods of anaesthesia are great advances, but the most important single factor is the skill and ready availability of the senior staff to deal personally with abnormalities. It can be shown that the maternal mortality rate in the cases delivered in the General Hospital is lower than the rate for the borough as a whole.

In the last ten years the borough deliveries numbered 14,833, with 43 maternal deaths—a rate of 2.89. In the same period the General Hospital, dealing with almost all the emergencies from inside and outside the borough, had 14,324 deliveries with 24 deaths, giving a rate of 1.67. And it is noteworthy that part of this decade was before the introduction of blood banks and penicillin, and that as late as 1946 (the last year for which complete figures are available) the national rate was 1.43.

In the Ministry of Health Report (1937) Halifax was shown to be the county borough with the blackest maternal mortality record in the country. It occupied an almost equally unenviable position in the R.C.O.G. Report (1944), in which for the years 1939 to 1942 it was placed seventh in a list of county boroughs with the highest rate in England and Wales. This, however, was largely due to the eleven maternal deaths in Halifax in 1940; only two of these occurred in the General Hospital and one on the Domiciliary Service.

It is true, then, to say that the high rate for the borough in a list of county boroughs with the highest rate in deaths in the General Hospital or on the Domiciliary Service, where in the same ten-year period there were 4,389 deliveries with the death of only one woman.

Summary and Conclusions

Along the lines suggested by the Royal College of Obstetricians and Gynaecologists (1944) we have attempted to bring about the integration of the hospital, antenatal, domiciliary, and specialist services in the Halifax Municipal Obstetric Organization.

Between Sept. 12, 1946, and Oct. 5, 1948, there were 5,000 deliveries after the twenty-eighth week without a maternal death due to pregnancy.

The stillbirth rate in the series was 23.07 and the neonatal death rate 11.91.

A brief summary of the treatment of some major complications has been given.

The Halifax General Hospital has a large proportion of cases from the county area, and therefore the number of its deliveries in the last decade almost equals the total ascribed to the borough.

In this ten-year period the maternal mortality rate in the General Hospital was 1.67; in the Halifax Borough it was 2.89.

The belief is expressed that the most important single factor in reducing maternal mortality is the provision of skilled staff to deal with abnormalities.

This series began after a maternal death from pulmonary embolism and continues without a maternal death attributable to pregnancy at the time of writing (Oct. 6, 1948).

I am grateful to the M.O.H. for Halifax, Dr. G. C. F. Roe, for permission to publish these results and for the statistics relating to the Borough. My thanks are due to Mr. H. I. Deitch, medical superintendent and surgeon to the General Hospital, for his great help in many difficult cases and for encouraging the development of the maternity department of the hospital.

REFERENCES

- Dieckmann, W. J. (1942). *The Toxaemias of Pregnancy*, p. 397. London.
Ministry of Health (1937). Report of an Investigation into Maternal Mortality. H.M.S.O., London.
R.C.O.G. (1944). Report on a National Maternity Service. London.
Thomas, R. C. (1940a). *British Medical Journal*, 1, 562.
— (1940b). *Ibid.*, 1, 745.

A STATISTICAL AND CLINICAL REVIEW OF 107 CASES OF ECTOPIC GESTATION

BY

J. S. MacVINE, M.B., B.S., F.R.C.S.Ed., M.R.C.O.G.

AND

D. H. LEES, M.B., Ch.B., F.R.C.S.Ed., M.R.C.O.G.

(From the Central Middlesex Hospital, London)

Ectopic pregnancy is a condition encountered by all clinicians engaged in general medical and surgical practice. The management usually falls to the gynaecologist, but the problem of the differential diagnosis concerns all, and in many cases the operative treatment has to be done by the general surgeon.

Keeping that in mind, we have undertaken a survey of all cases of ectopic pregnancy occurring in a busy municipal hospital over a ten-year period. The findings are instructive in showing that the question of aetiology has not finally been settled. It is, however, in the sphere of diagnosis that we believe the value of such a survey lies. The various symptoms and physical signs are evaluated in the light of the case records and an attempt is made to ascribe to each its true place and importance. The treatment and general results are also considered.

Despite the enormous literature on ectopic gestation, few recent surveys of this type have been made, and nearly

all of these were in America. Several will be referred to subsequently.

Present Series

The cases in this series numbered 107. The average age of the patients was 29.9 years. This is in accordance with previously noted figures. The ages varied between 20 and 42. On an average the patients had had 1.25 pregnancies, and 34% of all the pregnancies ended in abortion—a rather high figure.

The time interval between marriage and ectopic pregnancy, or between the ectopic and the last intrauterine pregnancy, averaged 5.8 years. This finding agrees with the figures of Woodhouse (1940), Ware and Winn (1941), Marchetti *et al.* (1946), and MacFarlane and Sparling (1946). It supports the view that sterile or relatively sterile women are more prone to ectopic gestation. Whether this is a result of pelvic inflammation is another matter, and will be dealt with more fully later.

Right- and left-sided tubal pregnancy occurred in 55 and 50 cases respectively. There is thus no real preponderance of the right side as has been suggested by Mayo and Strassmann (1938) and by Nucci (1946). The theory is attractive, as the right side is that more likely to be involved following lesions of the appendix. Ovarian pregnancy occurred in 2 (1.87%) of the 107 cases. The usually quoted figure is 1%. There were six cases of recurrent ectopic gestation—an incidence of 5.6%. Nucci's figure is 6.6%.

At operation it is extremely difficult to say where the ovum originally embedded. This is particularly so if there is an old rupture with haematocele formation or if a tubal abortion is in progress. In this series fully 50% were definitely ampullary, about 10% were fimbrial, and a further 20% might have been either of these. Isthmic forms made up 10% of the total, and at least two of the most severe type were interstitial.

Causation of Extrauterine Pregnancy

We have considered the causation under six headings, really working on a basis of Novak's (1947) classification, and we attempted to evaluate this as a criterion.

1. *Pelvic Inflammation.*—A history of anything possibly classifiable as genital infection was obtained in only seven cases. In 18 cases there was a history of one or more previous abortions, most of which took place some years before the ectopic pregnancy, and in no case was an abortion followed by prolonged subsequent upset. No patient gave a history of puerperal sepsis, but it is realized that some degree of puerperal or post-abortive morbidity may have been overlooked or forgotten. In several cases there was a normal delivery in the interval from the last abortion.

The operative findings were carefully recorded and gave the following information. In a total of eight cases there were some peritubal adhesions, but a certain proportion of these dated from the first blood leakage and were associated with the haematocele present. Hydrosalpinx was found in one case, and bilateral tubal occlusion following an old ileo-caecal tuberculous process in another. A chronically inflamed appendix was encountered at one operation for right tubal pregnancy.

These figures would suggest that pelvic inflammatory disease, particularly gonorrhoea, but also post-abortive and puerperal sepsis, has been much overrated as a cause or predisposing factor in ectopic pregnancy.

Many of the authorities are absolutely convinced of its importance, and Dougal (1937) estimates pelvic infection as a cause in 50% of cases. Novak (1947) supports this view, and Woodhouse (1940) puts the figure as high as 65.8%. These latter were cases giving a definite history, or showing evidence of pelvic inflammation at operation. MacFarlane and Sparling (1946) find a history of previous abortion in 32.7% of their cases and say that this is of unquestionable importance in the causation of ectopic pregnancy. They had a total of 17.2%

proved cases of pelvic infection. Marchetti *et al.* (1946) do not agree about the role of infection, and say that in most instances the aetiology is not clear. Dodds (1940) considered that in his series the possible predisposing causes were unconvincing.

2. *Previous Pelvic Operations.*—It does seem that previous operations on the pelvis may have a bearing, and it would be reasonable to expect conditions complicated by infection or peritonitis to be most prolific of ectopic gestation. In this series six patients had had appendicectomy, the appendix being ruptured, with local peritonitis, in one case. Six patients had had ectopic pregnancies, two had had ovariectomies, and two had had pelvic operations for tuberculous glands. Thus 16 of the 107 cases had previously had an operation. Farrell and Scheffey (1943) found that 26% of their cases had had previous abdominal operations. Dodds's (1940) figures for previous pelvic operations would work out at 20%.

3. *Congenital Abnormalities.*—The importance of congenital abnormalities or defects must be very difficult to estimate, and would be small.

4. *External Factors Altering the Lumen of the Tube.*—Alteration in calibre of the Fallopian tube and distortion due to external factors would seem to be a more likely cause. In this series we found the following:

Cystic or enlarged ovaries noted (not corpus luteum)	16 cases
Fimbrial or parovarian cysts	3 "
Uterine fibroids in a position to obstruct the affected tube	1 case
Tubal fibroid on uterine side of ectopic sac	1 "
Endometrioma of broad ligament	1 "

The enlarged or cystic ovaries were usually studded with a series of follicular cysts. These were seldom larger than $\frac{1}{4}$ in. (1.9 cm.) in diameter, but occasionally needed surgical attention. Many cases were not carefully examined, so that the incidence of these cysts is not fully recorded. The latter were the severe intraperitoneal rupture cases. Certain cases show this relationship very clearly. We have seen a tubal abortion in which the ectopic sac extended medially to a kink in the Fallopian tube caused by the pressure of a simple ovarian cyst the size of a pullet's egg.

5. *External Transmigration of the Ovum.*—One case afforded evidence of this. The ovary had previously been removed on the affected side and the tube on the opposite side.

6. *Tubal Spasm.*—This is held to be a very important factor, but we have not been able to collect any helpful information. In only two cases had a tubal insufflation been done—in both some time previously. A history of attempted abortion was given in three or four cases, but quite what the true incidence of this would be we do not know. From the very high rate of induced abortion in this particular district it would seem reasonable to regard it as a possible factor.

Diagnosis in Relation to Symptomatology

The proportion of cases in which particular symptoms appeared are as follows:

Abdominal pain	98%
Vomiting associated with the pain	41%
Faintness	42%
Anomalous uterine "bleeding"	76%
Amenorrhoea (including late or overdue period in a woman normally of regular cycle)	71%
Shoulder pain (including pain in the upper arm and in the upper chest or pectoral region)	37%
Bladder symptoms	26%
Rectal symptoms	16%

It will be seen that pain is by far the most reliable and constant symptom. It may be of any type, depending on the pathology. Pain sufficient to cause vomiting is a common description, and there is no doubt that it is a help in diagnosis. Nausea is not included in this estimate. The incidence of shoulder pain is higher than has previously been reported, and is a valuable diagnostic guide. The importance of bladder symptoms in diagnosis has never been fully recognized. These usually take the form of dysuria and frequency due to bladder irritation. In many of these cases urinary investigations were done and failed to reveal

any evidence of infection. Rectal irritation was evidenced as pain on defaecation, sacral pain, or diarrhoea.

There is general agreement on the importance of abdominal pain, amenorrhoea, and anomalous bleeding as the important triad. MacFarlane and Sparling (1946) found nausea and vomiting due to pain in 30% of their cases, while 35.4% complained of faintness due to pain; in 11.8% there were bladder symptoms. Marchetti *et al.* (1946) did not find nausea and vomiting with any regularity, although they noted faintness due to pain in 29.1% of cases and shoulder pain in 8.5%. Dodds (1940) reports that seven of his 19 acute cases had shoulder pain.

Diagnosis in Relation to Pathological Types

Cases of ectopic pregnancy may come under one's care at any stage of the pathological process, and the findings differ so much that some attempt at classification must be made. We have taken as a basis that of Dougal (1937). Briefly, the four types are: (a) where the ovum is still in the tube and there is little or no haemorrhage into the abdomen; (b) tubal rupture or abortion with diffuse intraperitoneal haemorrhage; (c) tubal rupture or abortion with encysted haemorrhage; and (d) advanced ectopic pregnancy. We have taken (a), (b), and (c) and termed them Types 1, 2, and 3, respectively. Our Type 1 group includes cases with a somewhat more generous blood leakage, while under Type 3 are included the subacute cases with haematocoele formation, even if not actually encysted.

Diagnosis in Relation to Physical Signs

The physical findings are considered in relation to the pathological type of ectopic gestation. Taking them in order, the incidence was as follows:

Type 1 (29 Cases)			
Tenderness in iliac fossa or suprapubically	89%		
Softening of the cervix	89%		
Tender swelling felt per vaginam	82%		
Uterine enlargement	62%		
Pain on moving the cervix	58%		
Breast activity	53%		
Pulsation felt in fornices	2%		

Type 2 (48 Cases)			
Tender on abdominal examination	100%		
Pain on vaginal examination (done in 39 cases only) ..	97%		
Fullness felt on vaginal examination (ditto) ..	87%		
Evidence of internal haemorrhage	85%		
Distended abdomen	77%		
Patient prostrate	56%		
Abdominal resistance or guarding	50%		
Free fluid in abdomen	41%		
Tube palpable	Nil		

Type 3 (28 Cases)			
Mass felt on vaginal examination (with or without general anaesthesia)	100%		
Tender on vaginal examination	96%		
Abdomen tender on examination	89%		
Softening of the cervix	88%		
Pain on moving the cervix	84%		
Uterus enlarged	65%		
Uterus felt separate from the swelling	61%		
Breast activity	54%		
Patient pale in appearance	54%		
Abdomen distended	50%		
Patient apparently ill	46%		
Abdominal guarding or resistance	32%		
General feeling of tumour on abdominal examination ..	32%		

Thus it is seen that all types, and particularly Type 2, have abdominal tenderness.

Pain on vaginal examination is the other common finding. This is usually elicited in the posterior or lateral fornix. A tender swelling is the usual accompaniment in Type 1, a generalized fullness is all that can be made out in Type 2, while a vague mass in the cul-de-sac is the common finding in Type 3.

Softening of the cervix is a very important diagnostic point and occurs in nearly 90% of Type 1 and Type 3

cases; it is of less diagnostic importance in Type 2, and is often overlooked in the obvious cases of that group. Pain on moving the cervix with the finger would appear to be a less reliable finding than tenderness in the fornices.

The presence of breast activity was a great diagnostic help in the less acute cases. Where there had been frank rupture and diagnosis was obvious it was surprising how often the breasts were not examined or at least no record of their condition was made.

Enlargement of the uterus was noted in 60–65% of Type 1 and Type 3 cases; it cannot be demonstrated in Type 2 cases.

In Type 2 there is very often evidence of internal haemorrhage, the abdomen is rather distended, and free fluid may be demonstrated. There is some degree of resistance or guarding in about half of the cases.

In Type 3 the patient often looks pale or ill. The average pulse rate was worked out for this group, and was 104 per minute. The temperature averaged between 99 and 100° F. (37.2 and 37.8° C.), with evening readings as high as 101° F. (38.3° C.). There may be some distension of the abdomen, but a tumour is rarely felt, and rigidity is also rare. In fully half of the cases the uterus was felt separate from the pelvic mass on vaginal examination.

The other signs, the blood investigations, pregnancy diagnosis tests, and curettage are usually too slow and unwieldy to be of value. We must admit, however, that the pathologist's report led to our exploring one of the cases.

Reference to the literature shows no great disagreement on these matters. Some authors stress the importance of one sign and some favour another. It must be remembered that diagnostic criteria are comparable only when a similar stage of the pathological process is encountered. There is a tendency both here and in America to discount the value of colpotomy, curettage, and haematological investigations in diagnosis.

Treatment

Blood Transfusion.—It has been our custom to give blood at once to the Type 2 cases admitted in a state of collapse. The theatre is meantime prepared and the patient is taken there with the blood-drip transfusion running. This is continued during the operation, and afterwards for as long as is necessary. We do not agree with those who say that blood should not be given before the operation is begun. Neither do we agree that operation should be postponed (in Type 2) till the general condition is greatly improved. The number of cases requiring blood transfusion was 41, and the average amount of blood given to each transfused patient was 1,080 ml. A reaction in the form of raised temperature and slight jaundice was observed in one case, but was quickly overcome. In nearly every case group O stored blood was given.

Williams and Corbit (1944), investigating 101 deaths from ectopic pregnancy, found that only 37 of the patients had received blood transfusion. Only four had more than one transfusion, although the time interval could have allowed of others. MacFarlane and Sparling (1946) transfused 57% of their cases, while Marchetti *et al.* (1946) transfused 42.16% of theirs.

Operative Treatment.—It was Lawson Tait (1888) who thoroughly established the treatment of ectopic pregnancy. The treatment is surgical and the earlier it is done the better. We think that if the patient is exsanguine and shocked it is reasonable to start blood replacement at once, with the other resuscitation measures. So far as the actual operation is concerned, it is advisable to do the least possible when the patient is in a serious condition. The affected tube is removed and haemostasis secured. In this series the following operations were done:

Salpingectomy	77
Salpingo-oophorectomy (one bilateral)	24
Oophorectomy	2
Removal of sac of pregnancy only	1
Partial salpingectomy	2
Laparotomy only	1
Cysts necessitating ovariectomy or cystectomy	7

Thus it will be seen that salpingectomy is usually sufficient, and in common with others we advocate total salpingectomy. The risk of recurrence in the same tube is otherwise said to be high. It is mainly in Type 3 cases that salpingo-oophorectomy is necessary, and is due to involvement of the ovary in the haematocoele to such an extent that it cannot be dissected clear. Ovariectomy is necessary in cases of ovarian pregnancy. The comparatively large number of cases with some cystic involvement of the ovaries will be noted, and was mentioned previously.

The above procedure is fairly standard. Mayo and Strassmann (1938) stress the importance of conservative measures. Dodds (1940) preserves the ovaries wherever possible, while Dougal's (1927) figures show that he was less conservative in treatment. There has apparently been a tendency in America to do operations for other abnormalities discovered at the time of dealing with the ectopic pregnancy. Williams and Corbit's (1944) investigations of mortality figures have led them to condemn the doing of multiple non-essential operations.

There was one death in this series, giving a mortality rate of 0.9%.

The number of days in hospital before operation was taken as a method of assessing the accuracy of diagnosis. All the days the patient spent in hospital before the operation are counted. The average time was 2.8 days. Marchetti *et al.* (1946) record that 27% of their cases were operated on within 24 hours of admission, while 70.2% had operation within 48 hours.

Summary

A series of 107 consecutive ectopic pregnancies occurring over a ten-year period is reviewed. The findings are compared with those of recent reviews.

In respect of aetiology we find little support for the time-honoured view that previous pelvic inflammation is all-important. We believe that previous operations on the pelvis are a predisposing factor, and we are strongly of the view that extratubal factors causing alterations in calibre of the lumen are an important cause.

Diagnosis is considered in relation to symptomatology, pathological types, and physical findings. Our figures show that carefully taken histories reveal a very much higher incidence of helpful diagnostic symptoms than has previously been reported. For example, we find a high percentage of those patients complaining of vomiting and faintness due to pain, of shoulder pain, and of irritative bladder symptoms.

The treatment of these cases has been orthodox in every way. We attach particular importance to the free use of blood pre-operatively, without delaying the operation once the diagnosis has been made.

REFERENCES

- Dodds, R. L. (1940). *Proc. R. Soc. Med.*, **33**, 783.
 Dougal, D. (1927). *British Medical Journal*, **2**, 1074.
 — (1937). *British Encyclopaedia of Medical Practice*, **5**, 258.
 Butterworth, London.
 Farell, D. M., and Scheffey, L. C. (1943). *Amer. J. Obstet. Gynec.*, **46**, 686.
 MacFarlane, K. T., and Sparling, D. W. (1946). *Ibid.*, **51**, 343.
 Marchetti, A. A., Kuder, K., and Kuder A. (1946). *Ibid.*, **52**, 544.
 Mayo, C. W., and Straussmann, E. O. (1938). *Surg. Gynec. Obstet.*, **67**, 46.
 Novak, E. (1947). *Gynecological and Obstetrical Pathology*, 2nd ed., p. 437. Saunders, Philadelphia.
 Nucci, R. C. (1946). *Penn. med. J.*, **49**, 953.
 Tait, R. Lawson (1888). *Lectures on Ectopic Pregnancy and Pelvic Haematocoele*. Birmingham.
 Ware, H. H., jun., and Winn, W. C. (1941). *Amer. J. Obstet. Gynec.*, **42**, 33.
 Williams, P. F., and Corbit, J. D. (1944). *Ibid.*, **48**, 841.
 Woodhouse, K. W. (1940). *Amer. J. Surg.*, **49**, 3.

ENDOGENOUS GAS GANGRENE COMPLICATING CARCINOMA OF COLON

REPORT OF A CASE

BY

A. L. WYMAN, M.D., M.R.C.P.

Late Assistant Medical Officer, Lambeth Hospital

Clostridium welchii is commonly present in the human intestine. Its pathogenic qualities are well known, yet it is rare for gas gangrene to complicate diseases or surgery of the alimentary canal unless introduced from outside. Rare though it is, this complication has long been recognized. Welch (1900) gave three examples of primary emphysematous gangrene of the abdominal wall which he believed started from the intestine. Greeley (1918) described two cases of "idiopathic *Bacillus aerogenes capsulatus* infection." In one of his patients the blood culture was positive three months before death. In the case described below, the organism was grown from the blood 17 days before death.

Several writers have reported endogenous gas gangrene arising after operation, especially after appendicectomy (Eckhoff, 1930; Gamble, 1935; Johnson, 1936; Smith and Zimring, 1939; Breslin, 1940). Jennings (1931) found that 90% of cultures made from the contents of the lumen of appendices removed at operation grew *Cl. welchii*.

Gas gangrene may arise even when no operation has taken place. Gordon (1936) treated conservatively an appendix abscess which ruptured and formed a pelvic abscess. Gas gangrene of the right thigh developed, and at necropsy it was seen that the peritoneum and fascia overlying the iliacus muscle had been eroded and that the gas gangrene had spread down this muscle to the lower limb.

It may also arise after typhoid ulceration and tuberculous ulceration of the small intestine (Knapper, 1929; Galiana and Balaguer, 1946). Dawson and Hardy (1948) reported emphysematous cellulitis of the left thigh resulting from diverticulitis.

Case Report

The patient, a man aged 65, had all his teeth extracted in September, 1947. A few days afterwards he felt some pain in the lower abdomen on the left side. He had to strain to defaecate, and when he was successful the pain disappeared. Later he felt as though his bowels were not completely emptied after defaecation, and the pain was not completely relieved. The motions were soft, without blood or slime. His appetite was poor and he had lost weight. There was no nausea or vomiting, and urination was normal. He was admitted to hospital on Nov. 14, having been febrile for four days. He had had an operation for bilateral inguinal hernia in 1915.

On examination he was flushed and looked ill. The temperature was 102° F. (38.9° C.), pulse 80, and respiration 20. The gums were healthy, the throat clean, and the tongue furred. Some petechiae were seen on the floor of the mouth. The abdomen was tumid and there was a tender indurated area with slight erythema above the left Poupart's ligament. The herniotomy scars were healthy. Neither spleen nor liver was felt. Intestinal movements were audible. Rectal examination revealed nothing abnormal. There was a slight degree of kyphoscoliosis of the spine but no local tenderness. The patient kept his hips flexed, and there was some pain in the left hip on external rotation. A blood culture was taken before any treatment was given.

The next day the reddening of the skin was more apparent in a line 6-8 in. (15-20 cm.) long, about 1 in. (2.5 cm.) above and parallel to the left inguinal ligament. He was very tender near the left anterior-superior iliac spine. Penicillin and sulphamezathine were given.