Salpingectomy	77
Salpingo-oophorectomy (one bilateral)	24
Oophorectomy	2
Removal of sac of pregnancy only	1
Partial salpingectomy	2
Laparotomy only	1
Cysts necessitating ovariectomy or cystectomy	7

Thus it will be seen that salpingectomy is usually sufficient, and in common with others we advocate total salpingectomy. The risk of recurrence in the same tube is otherwise said to be high. It is mainly in Type 3 cases that salpingo-oophorectomy is necessary, and is due to involvement of the ovary in the haematocoele to such an extent that it cannot be dissected clear. Ovariectomy is necessary in cases of ovarian pregnancy. The comparatively large number of cases with some cystic involvement of the ovaries will be noted, and was mentioned previously.

The above procedure is fairly standard. Mayo and Strassmann (1938) stress the importance of conservative measures. Dodds (1940) preserves the ovaries wherever possible, while Dougal's (1927) figures show that he was less conservative in treatment. There has apparently been a tendency in America to do operations for other abnormalities discovered at the time of dealing with the ectopic pregnancy. Williams and Corbit's (1944) investigations of mortality figures have led them to condemn the doing of multiple non-essential operations.

There was one death in this series, giving a mortality rate of 0.9%.

The number of days in hospital before operation was taken as a method of assessing the accuracy of diagnosis. All the days the patient spent in hospital before the operation are counted. The average time was 2.8 days. Marchetti *et al.* (1946) record that 27% of their cases were operated on within 24 hours of admission, while 70.2% had operation within 48 hours.

Summary

A series of 107 consecutive ectopic pregnancies occurring over a ten-year period is reviewed. The findings are compared with those of recent reviews.

In respect of aetiology we find little support for the time-honoured view that previous pelvic inflammation is all-important. We believe that previous operations on the pelvis are a predisposing factor, and we are strongly of the view that extratubal factors causing alterations in calibre of the lumen are an important cause.

Diagnosis is considered in relation to symptomatology, pathological types, and physical findings. Our figures show that carefully taken histories reveal a very much higher incidence of helpful diagnostic symptoms than has previously been reported. For example, we find a high percentage of those patients complaining of vomiting and faintness due to pain, of shoulder pain, and of irritative bladder symptoms.

The treatment of these cases has been orthodox in every way. We attach particular importance to the free use of blood pre-operatively, without delaying the operation once the diagnosis has been made.

REFERENCES

- Dodds, R. L. (1940). *Proc. R. Soc. Med.*, **33**, 783.
 Dougal, D. (1927). *British Medical Journal*, **2**, 1074.
 — (1937). *British Encyclopaedia of Medical Practice*, **5**, 258.
 Butterworth, London.
 Farell, D. M., and Scheffey, L. C. (1943). *Amer. J. Obstet. Gynec.*, **46**, 686.
 MacFarlane, K. T., and Sparling, D. W. (1946). *Ibid.*, **51**, 343.
 Marchetti, A. A., Kuder, K., and Kuder A. (1946). *Ibid.*, **52**, 544.
 Mayo, C. W., and Straussmann, E. O. (1938). *Surg. Gynec. Obstet.*, **67**, 46.
 Novak, E. (1947). *Gynecological and Obstetrical Pathology*, 2nd ed., p. 437. Saunders, Philadelphia.
 Nucci, R. C. (1946). *Penn. med. J.*, **49**, 953.
 Tait, R. Lawson (1888). *Lectures on Ectopic Pregnancy and Pelvic Haematocoele*. Birmingham.
 Ware, H. H., jun., and Winn, W. C. (1941). *Amer. J. Obstet. Gynec.*, **42**, 33.
 Williams, P. F., and Corbit, J. D. (1944). *Ibid.*, **48**, 841.
 Woodhouse, K. W. (1940). *Amer. J. Surg.*, **49**, 3.

ENDOGENOUS GAS GANGRENE COMPLICATING CARCINOMA OF COLON

REPORT OF A CASE

BY

A. L. WYMAN, M.D., M.R.C.P.

Late Assistant Medical Officer, Lambeth Hospital

Clostridium welchii is commonly present in the human intestine. Its pathogenic qualities are well known, yet it is rare for gas gangrene to complicate diseases or surgery of the alimentary canal unless introduced from outside. Rare though it is, this complication has long been recognized. Welch (1900) gave three examples of primary emphysematous gangrene of the abdominal wall which he believed started from the intestine. Greeley (1918) described two cases of "idiopathic *Bacillus aerogenes capsulatus* infection." In one of his patients the blood culture was positive three months before death. In the case described below, the organism was grown from the blood 17 days before death.

Several writers have reported endogenous gas gangrene arising after operation, especially after appendicectomy (Eckhoff, 1930; Gamble, 1935; Johnson, 1936; Smith and Zimring, 1939; Breslin, 1940). Jennings (1931) found that 90% of cultures made from the contents of the lumen of appendices removed at operation grew *Cl. welchii*.

Gas gangrene may arise even when no operation has taken place. Gordon (1936) treated conservatively an appendix abscess which ruptured and formed a pelvic abscess. Gas gangrene of the right thigh developed, and at necropsy it was seen that the peritoneum and fascia overlying the iliacus muscle had been eroded and that the gas gangrene had spread down this muscle to the lower limb.

It may also arise after typhoid ulceration and tuberculous ulceration of the small intestine (Knapper, 1929; Galiana and Balaguer, 1946). Dawson and Hardy (1948) reported emphysematous cellulitis of the left thigh resulting from diverticulitis.

Case Report

The patient, a man aged 65, had all his teeth extracted in September, 1947. A few days afterwards he felt some pain in the lower abdomen on the left side. He had to strain to defaecate, and when he was successful the pain disappeared. Later he felt as though his bowels were not completely emptied after defaecation, and the pain was not completely relieved. The motions were soft, without blood or slime. His appetite was poor and he had lost weight. There was no nausea or vomiting, and urination was normal. He was admitted to hospital on Nov. 14, having been febrile for four days. He had had an operation for bilateral inguinal hernia in 1915.

On examination he was flushed and looked ill. The temperature was 102° F. (38.9° C.), pulse 80, and respiration 20. The gums were healthy, the throat clean, and the tongue furred. Some petechiae were seen on the floor of the mouth. The abdomen was tumid and there was a tender indurated area with slight erythema above the left Poupart's ligament. The herniotomy scars were healthy. Neither spleen nor liver was felt. Intestinal movements were audible. Rectal examination revealed nothing abnormal. There was a slight degree of kyphoscoliosis of the spine but no local tenderness. The patient kept his hips flexed, and there was some pain in the left hip on external rotation. A blood culture was taken before any treatment was given.

The next day the reddening of the skin was more apparent in a line 6-8 in. (15-20 cm.) long, about 1 in. (2.5 cm.) above and parallel to the left inguinal ligament. He was very tender near the left anterior-superior iliac spine. Penicillin and sulphamezathine were given.

In the course of the next few days the inflammation spread down the thigh, chiefly on its anterior and medial aspects, as far as the knee. The skin was red and brawny, with a small necrotic patch in the mid-thigh. A fine crepitation in the region of the anterior-superior spine was palpable. On Nov. 19 Mr. W. K. Targett incised the thigh. "With a large aspirating needle, foul-smelling greyish-brown pus was obtained fairly superficially in the thigh (inner side). An incision in the same area opened a superficial 'space' containing gas and more pus. This was lined with greyish necrotic material. A drain was inserted. A second incision lower down was made. More gas and pus were found. The whole process was more like a generalized infiltration than a true abscess." Anti-gas-gangrene serum was given.

After the operation the patient's general condition improved a little, but locally there was an extension of the inflammation below the knee, with effusion in that joint and oedema of the leg. A black bulla was present in the thigh. Soon a necrotic area joined the two incisions. Subcutaneous crepitus could be felt in the upper and outer part of the thigh (Fig. 1).

FIG. 1.—Wound in left thigh and extent of inflammation one week after operation

On Nov. 27 a fresh spontaneous sinus appeared near the knee posteriorly. The following day a morbilliform rash attributed to the sulphamezathine appeared on the right thigh and then spread generally. By Dec. 1 he looked worse; he became drowsy and dyspnoeic, and died the next day.

Investigations.—Nov. 15:—blood culture: growth of *Cl. welchii*; blood count: Hb, 90%; red cells, 5,140,000; white cells, 17,000 (polymorphs 81%). Nov. 17:—radiography: chest normal; spine and pelvis, no destructive bone changes seen; jaws edentulous. Nov. 19:—swab from wound: direct film—many Gram-positive and Gram-negative bacilli and cocci; aerobic culture gave a growth of non-haemolytic streptococci, coliform bacilli, and Gram-positive sporing bacilli. Anaerobic culture: no anaerobes isolated. Nov. 25:—radiography of left thigh: considerable gas present in the soft tissues, consistent with gas gangrene (Fig. 2).

FIG. 2.—Radiograph of thigh showing gas in soft tissues

Post-mortem Examination.—A fungating carcinoma of the descending colon, 2 by 3 cm., was found just below the anterior-superior spine of the ilium. The carcinoma extended three-quarters of the way round the bowel and constricted the lumen but did not occlude it. There was a small stercoral ulcer 1 cm. proximal to the growth. The carcinoma was adherent to the psoas sheath and there was a track from the bowel into the psoas sheath, the contents of which were gangrenous. The superficial tissues over Poupart's ligament were also necrotic. Sections showed an anaplastic spheroidal-cell carcinoma of the colon with a marked trabecular formation. It had eroded through all layers of the gut wall. The muscle showed fragmentation with much oedema and hyaline degeneration. There were numerous clumps of *Cl. welchii* in the spleen. All swabs taken grew *Cl. welchii*.

Summary

An account is given of a patient with carcinoma of the colon presenting as a gas gangrene of the left thigh. The infection had spread from the intestine to the psoas muscle by means of a track connecting the carcinoma to the psoas sheath.

Similar examples of gas gangrene resulting from diseases of the bowel are quoted from the literature.

I am greatly indebted to Mr. W. K. Targett for the surgical treatment of the patient, and to Dr. J. Hewlett and Dr. G. Vincent for the pathological investigations.

REFERENCES

- Breslin, F. J. (1940). *Amer. J. Surg.*, **49**, 501.
 Dawson, R. L. G., and Hardy, R. H. (1948). *British Medical Journal*, **1**, 498.
 Eckhoff, N. L. (1930). *Brit. J. Surg.*, **18**, 38.
 Galiana, S. C., and Balaguer, J. P. (1946). *Rev. clin. esp.*, **23**, 405.
 Gamble, H. A. (1935). *Amer. J. Surg.*, **28**, 389.
 Gordon, S. (1936). *Brit. J. Surg.*, **24**, 399.
 Greeley, H. (1918). *Bull. Johns Hopk. Hosp.*, **29**, 231.
 Jennings, J. E. (1931). *Ann. Surg.*, **93**, 828.
 Johnson, W. W. (1936). *Amer. J. Surg.*, **33**, 141.
 Knapper, C. (1929). *Ned. Tijdschr. Geneesk.*, **2**, 4792.
 Smith, P. E., and Zimring, J. G. (1939). *Amer. J. Surg.*, **45**, 351.
 Welch, W. H. (1900). *Bull. Johns Hopk. Hosp.*, **11**, 185. Reprinted in *Medical Classics*, 1941.

ENTERITIS NECROTICANS DUE TO CLOSTRIDIUM WELCHII TYPE F

BY

J. ZEISSLER, M.D.

AND

L. RASSFELD-STERBERG

(From the Bacteriological Institute, General Hospital, Hamburg-Altona)

In Hamburg on Oct. 8, 1946, Beckermann and Laas gave an account of a number of cases of a severe illness, often fatal, due to a necrotic inflammation of several areas of the intestine, especially in the jejunum. The onset was acute, with very severe pain and slight rigidity, mostly in the left lower abdomen, without obvious guarding. Vomiting and profuse diarrhoea resulted in general dehydration; within a few days extreme circulatory collapse occurred, with general cyanosis. Increased blood urea, low blood chloride, indicanaemia, and indicanuria were often demonstrable. The temperature was slightly raised (up to 38° C.), the blood picture showed a moderate leucocytosis with a marked shift to the left, and the blood sedimentation rate was increased. Surgical and post-mortem examination showed a diffuse sloughing enteritis of the jejunum, ileum, and colon, beginning in the terminal part of the duodenum or the first loop of the jejunum. The most severe lesions were tube-like areas of necrosed mucosa a foot or more in