is any less liable to infection than a series of mustard burns of similar intensity and distribution. Nor has it been shown that under these circumstances the thermal burns would heal any more quickly than the mustard burns. Until such an investigation has been made it is begging the question to hold that the mustard burn is specially prone to sepsis and slow to heal.

Summary

An account is given of some observations made on the treatment of a series of skin burns caused by mustard gas under tropical conditions. It was not possible to make any accurately controlled investigation, but it appeared that no local treatment achieved better results than could be expected in untreated burns.

On the basis of factors such as comfort, convenience, expense, etc., zinc cream (*A.P.F.*) and vaselined gauze were the best of the treatments investigated for breaches of skin continuity. Triple dye, used as a paint, was valuable for treating small superficial lesions.

Suspensory bandages were of great value in the treatment of scrotal burns.

There was a comparatively low incidence of sepsis, in spite of the fact that the burns were treated under an approximation to field conditions.

There is no reason to suppose that mustard burns behave any differently from thermal burns in regard to their healing or liability to infection.

The kindness of the Australian Chemical Defence Board in permitting publication of the work carried out under its direction is gratefully acknowledged. Permission to publish has also been received from the chief scientist, Ministry of Supply.

I should like also to record my indebtedness to Mr. J. W. Legge and to the numerous colleagues who assisted in the application, supervision, and assessment of the treatments investigated.

REFERENCES

- Chiesman, W. E. (1944). *British Medical Journal*, 2, 109.
Cullumbine, H., and Sinclair, D. C. (1942). Unpublished work.
Davis, M. I. J. (1944). *J. Amer. med. Ass.*, 126, 209.
Koonitz, A. R. (1944). *Arch. Surg.*, 48, 284.
Mackintosh, J. M. (1920). "The Clinical Aspects of Mustard Poisoning." Glasgow University M.D. thesis.
Ministry of Health (1942). E.M.S. Instructions, Part I: Medical Treatment and Special Centres. Supplement 43. H.M.S.O., London.
Sinclair, D. C. (1944). Unpublished work.
— (1948a). *British Medical Journal*, 2, 290.
— (1948b). *Brit. J. Derm. Syph.* In press.
Soltau, A. B., and Elliott, T. R. (1912). *Official History of the Great War: Medical Services: Diseases of the War*, vol. II. H.M.S.O., London.
Stewart, R. L. (1937). *British Medical Journal*, 1, 380.
War Office (1944). *Army Medical Department Bulletin*, No. 31. London.

INCIDENCE OF DIABETES MELLITUS IN CHILDREN AND NEED FOR HOSTELS

BY

P. HENDERSON, M.D., D.P.H.

Medical Officer, Ministry of Education

In an annotation in the *British Medical Journal* of Feb. 7, 1948 (p. 261), it was stated that "the incidence of diabetes is unknown, although some idea of its prevalence can be obtained from the mortality returns." Lawrence (1944) wrote: "Diabetes has been recorded in infants of a few months, but is very rare under 6 years of age." In America several surveys have been made. Joslin (1946) quoted the findings of the National Health Survey of 2,500,000 persons, carried out under the auspices of the United States Public Health Service in 1935-6, when it was found that in the age group 0-14 years one child in every 2,500 was a diabetic: the incidence was the same for boys and girls, and

for all age groups was 0.37% of the total population (adults and children). Wilkerson and Krall (1947) reported the results of a thorough investigation into the incidence of the disease in Oxford, Mass. Of the 4,983 inhabitants 3,516 had their blood and urine tested. After allowing for the 29.4% of the town's citizens who had not been tested it was estimated that 1.7% of the population had diabetes.

Joslin (1946) also mentioned that the mortality rate from diabetes in the U.S.A. in 1943 was 27.1 per 100,000 of the population; in England and Wales in 1944 it was 9.6 per 100,000. The much lower diabetes death rate in England and Wales supports the view that the incidence of the disease is considerably less in these islands than it is in the U.S.A.

Present Investigation

In the absence of up-to-date information about the extent of diabetes in children in this country it was decided to make a survey, during the period February-June, 1948, in the following areas: Bristol, Birmingham, Birkenhead, Wallasey, Liverpool, Bootle, Salford, Manchester, Bradford, Leeds, Newcastle-upon-Tyne, Sheffield, Norwich, Ipswich, Great Yarmouth, Cambridge, Norfolk, Isle of Ely, Cambridgeshire, and East and West Suffolk.

All the school medical officers (who were, with one exception, also medical officers of health), the professors of child health, and all the hospitals in these areas likely to have child patients with diabetes were approached for information. The physicians of 32 hospitals co-operated: only one hospital ignored the requests made to it.

All lists of children supplied by the hospitals were cross-checked against the lists held by the school medical officers and the medical officers of infant welfare clinics to avoid duplication of names and to ensure that children not living in these areas were excluded, since many of the hospitals had patients from a wide countryside and from districts not included in this survey.

Size of Population Sampled.—The 16 cities and towns and the five counties selected for this inquiry had a total child population under 16 years of age of 1,307,000; there were about 750,000 school-children, being between one-sixth and one-seventh of the total school population of England and Wales: the rate was broadly similar for urban and rural children. Table I gives the number and age distribution of the children.

TABLE I.—Age Distribution

Age	Urban	Mainly Rural	Total
0-4 ..	470,000	70,000	540,000
5-9 ..	386,000	50,000	436,000
10-15 ..	290,000	41,000	331,000
Total	1,146,000	161,000	1,307,000

TABLE II.—Incidence of Diabetes in Sample Population

Age	No. of Children with Diabetes		
	Boys	Girls	Total
0-4 ..	1	2	3 (1 in 180,000)
5-9 ..	28	28	56 (1 in 8,000)
10-15 ..	70	54	124 (1 in 3,000)
Total ..	99	84	183 (1 in 7,000)

Of the 180 school-children (Table II) 145 (1 in 5,000) lived in urban areas, and 35 (1 in 3,000) lived in mainly rural areas. If the sample can be taken as representative of the child population then the estimated number of diabetic school-children in England and Wales is 1,200 (1 in every 4,000-5,000).

Number of Diabetic Children in Need of Hostel Treatment

Several of the physicians who provided information expressed the opinion that diabetic children should remain at home and not be sent to hostels. One expressed the view of several when he wrote: "All diabetic children should be brought up to lead normal lives with other children, and I find that this is possible with nearly all of them." Another wrote: "I do feel that on the whole children suffer more by being moved from their home environment than they gain by going to live in a hostel." Others, however, considered that hostels were essential for a small minority of diabetic children; one gave the history of a girl aged 12 who spent most of her life in two hospitals, having been admitted to each alternately in diabetic coma. Another child, who was not receiving the treatment prescribed, was found to have a mother who did not understand the instructions given to her as "she was deaf and otherwise peculiar."

An interesting finding was the greater demand for hostels for children from rural areas. Table III gives the number and age distribution of the children in the sample population who were recommended for admission to a hostel.

TABLE III.—*Children Recommended for Admission to Hostel*

Age	Boys	Girls	Total
0-4	1	—	1 (1 in 500,000)
5-9	4	3	7 (1 in 60,000)
10-15	7	6	13 (1 in 25,000)
Total	12	9	21 (1 in 60,000)

Of these 21 children seven were already in a hostel or hospital. Thus in this series one diabetic child in every nine was recommended for a hostel. On these figures there is need for hostels for at least 130 diabetic children in England and Wales. The survey also showed that in mainly rural areas one school-child in every 13,000, and in urban areas one in every 50,000, should be in a hostel for diabetics. The main reason for the greater need for hostels for rural children is the lack of specialist treatment facilities within easy reach of their homes. The difference in the incidence of diabetes between the urban and rural population was not sufficient to explain the great disparity in the proportion of school-children recommended for a hostel in these two groups.

Since September, 1939, the London Education Authority has maintained a diabetic unit for 60 boys and girls at its large residential school at Hutton, Essex. On the average, only about 16 of the 60 are London children, giving a rate of one child per 20,000 of the London school population. In view of the effects of the war on London homes it was to be expected that more London diabetic children would require hostel accommodation than children in provincial cities and towns. Of the 16 London children living in Hutton during the period of this inquiry eight were fatherless, motherless, or had parents who were separated.

Allowing for the higher rate of hostel provision required for London and rural children, rather more than 130 places are needed. On the basis of the present investigation hostels for 130-150 diabetic children should be provided for a school population of about 5,000,000. In addition to the 60 places at Hutton a second hostel for 30 diabetic boys and girls was opened at Deal by the Church of England Children's Society in January, 1949. There are prospects of a third hostel for about 30 children being opened in 1949.

Anyone having doubts about the value of a hostel for a minority of diabetic children should visit Hutton. There the boys and girls are taught the diabetic way of life; they test their own urine and, under supervision, inject their own

insulin. They are supervised by a trained nursing staff, and a specialist physician visits fortnightly. They lead as free and as normal a life as any child in a public boarding school. Their out-of-school life is full of variety. They take part in any physical activity or game that they choose and they all learn to swim in the school swimming-bath. Most are allowed home for the usual school holidays. In early summer they all go to camp for a fortnight at the seaside. They grow up into self-reliant men and women, fully capable of leading happy and useful lives. Unfortunately, some parents are so foolish that they prematurely withdraw their children from the hostel. Whittaker (1948) reported the deaths of five children who had been removed from Hutton against advice. And in this country in 1945, according to the Registrar-General (1945), 51 children under 15 years of age died from diabetes. These melancholy figures represent a yearly wastage of young lives that could be reduced considerably if adequate hostel facilities were available.

Summary

Of 1,307,000 children under 16 years of age (being from one-sixth to one-seventh of the total child population under 16 years of age in England and Wales) living in 16 English cities and towns and five counties, 183 (1 in 7,000) were known to have diabetes. Of the 180 school-children, 145 (1 in 5,000) lived in urban areas and 35 (1 in 3,000) in mainly rural areas. An incidence rate of 1 in 7,000 is only about one-third of that (1 in 2,500) found in 1935-6 in the survey sponsored by the U.S. Public Health Service.

The incidence rose sharply from 1 in 180,000 for children under 5 years to 1 in 8,000 for children 5-9 years and to 1 in 3,000 for children aged 10-15 years.

About one diabetic child in nine was recommended for a hostel. In the mainly rural areas one school-child in every 13,000, and in the cities and towns one in every 50,000, was considered suitable for a hostel; in London the rate was 1 in 20,000. The higher London figure was at least partly due to the effects of the last war, since half the total number of the London children were fatherless, motherless, or had parents who were separated. The high rural rate was mainly due to the absence of specialist treatment facilities within easy reach of the children's homes.

Allowing for the greater need for hostels for diabetic children in London and the rural areas, this survey revealed the need for hostels for 130-150 children under 16 years of age in England and Wales with a school population of about 5,000,000. At the time of this investigation there was hostel accommodation for only 60 diabetic children in the London County Council's School at Hutton. A second hostel for 30 children was opened in Deal in January, 1949, by the Church of England Children's Society, and there are prospects of a third hostel being opened during 1949.

I gratefully acknowledge the help given to me by the school medical officers of the areas mentioned in this article and by the professors of child health, physicians, almoners, and secretaries of the hospitals in these areas who so willingly supplied me with information.

REFERENCES

- Joslin, E. P. (1946). *The Treatment of Diabetes Mellitus*, p. 33. London.
 Lawrence, R. D. (1944). *The Diabetic Life*, p. 127. London.
 Registrar-General (1945). *Statistical Review of E. & W.*, Part 1, Medical, p. 119. London.
 Whittaker, H. (1948). *Brit. med. Students' J.*, 2, 14.
 Wilkerson, H. L. C., and Krall, L. P. (1947). *J. Amer. med. Ass.*, 135, 209.

The Minister of Health and Local Government of Northern Ireland stated recently that plans are on foot to provide this year 200 additional beds for tuberculosis sufferers in Northern Ireland. The Minister added that he would further consider a suggestion that the Government should make representations to the proper authorities towards making arrangements for the treatment in Swiss hospitals of patients suffering from tuberculosis in view of the present shortage of beds in Northern Ireland.