Discussion

Induced epilepsy is likely to be of value only in the case of simulated inability to remember identity. The usual procedures should be employed in true conditions of psychogenic amnesia. Induced epilepsy is valueless in restoring truth in any other kind of lie (except perhaps hysterical mutism, deafness, or blindness) because it uses a neural phenomenon which makes the patient able to identify himself correctly before he is capable of remembering external events and sorting them out.

Summary

A case is described in which induced epilepsy was employed to establish identity in a patient who feigned amnesia for his name and the total previous period of his life. This type of amnesia and the neuro-physiological rationale of the treatment are discussed.

REFERENCES

Hemphill, R. E., and Walter, W. Grey (1941). *J. ment. Sci.*, **87**, 256.
Parfitt, D. N. (1944). *Ibid.*, **90**, 511.

AN UNUSUAL CASE OF CHRONIC MYELOID LEUKAEMIA

BY

R. A. RUSSELL TAYLOR, F.R.C.S.

Medical Superintendent and Surgeon, Pinderfields General Hospital, Wakefield

The interesting feature of the following case is the response to treatment devised for a mixture of cirrhosis of the liver and myeloid leukaemia.

Case History

The patient, a man aged 61, was admitted to this hospital on March 13, 1946, with chronic myeloid leukaemia, marked splenomegaly, and ascites. A careful history revealed that he had been treated for pernicious anaemia since 1931, but the report of a blood count taken on Dec. 8, 1931, failed to substantiate this diagnosis. It was as follows: haemoglobin, 76%; red cells, 3,820,000; colour index, 1; white cells, 9,600 (polymorphs 75%, lymphocytes 15%, eosinophils 1%, basophils 2%, large hyaline cells 2%, transitional cells 5%).

The next report to which we had access was dated March 11, 1941, and the blood count was as follows: haemoglobin, 98%; red cells, 4,300,000; colour index, 1.14; mean diameter of red cells, 7μ ; nucleated red cells, 1,000 per c.mm.; poikilocytes and polychromatic cells; white cells, 26,000 (polymorphs 56%, eosinophils 1%, basophils 2%, lymphocytes 9%, monocytes

6%, neutrophil myelocytes 20%, eosinophil myelocytes 1%, basophil myelocytes 3%, myeloblasts 2%). The report stated: "The white cell picture tends towards that of chronic myelogenous leukaemia."

On June 14, 1941, the white blood count had increased to 36,000, but the differential percentages remained constant.

No further reports were available until Nov. 3, 1944, when the blood picture was: haemoglobin, 69%; red cells, 3,420,000; colour index, 0.99; nucleated red cells, 1,270; reticulocytes, 2.2%; white cells, 35,700 (polymorphs 44%, eosinophils 1%, lymphocytes 25%, monocytes 12%, myeloblasts 4%, neutrophil myelocytes 11%, eosinophil myelocytes 1%, basophil myelocytes 2%). Blood counts were then taken at fairly regular intervals; these are recorded in Chart 1. In the differential counts the percentage of myelocytes varied between 6 and 30%, while that of the myeloblasts varied between 1 and 23%.

The most characteristic blood count was that taken on Aug. 24, 1948, and is therefore given in detail: red cells, 3,100,000; haemoglobin, 61%; colour index, 0.99; white cells, 17,130. Differential count was as follows: neutrophil polymorphs, 9,350 (55%); basophils, 510 (3%); neutrophil "staffs," 850 (5%); neutrophil metamyelocytes, 1,800 (10.5%); neutrophil myelocytes, 3,150 (18.5%); myeloblasts, 950 (5%); lymphocytes, 340 (2%); monocytes, 0; normoblasts, 90 (0.5%); erythroblasts, 90 (0.5%).

On March 14, 1946, paracentesis abdominis was performed and 12 pints (6.8 litres) of ascitic fluid were removed. The very marked and extensive oedema of both lower limbs was not affected in any way even after massage. On March 15 a blood transfusion of 2 pints (1.14 litres) of blood was given, and this was repeated on March 29, April 4, and May 24. The patient also received massive doses of iron daily.

Paracentesis abdominis was repeated on April 16, and 14 pints (7.95 litres) of ascitic fluid were removed. This process was repeated at rapidly diminishing intervals during the next three months, until as much as 24 pints (13.6 litres) were being removed every five days. The exact dates and amounts are

CHART 2.—Amounts of ascitic fluid withdrawn.

given in Chart 2. The protein content of the ascitic fluid varied between 2 and 2.5 g. %.

By this time the patient was markedly emaciated and the abdomen was grossly distended by ascitic fluid and a large spleen, which reached right down into the pelvis. His liver, however, was only slightly enlarged.

On July 2 an estimation of the plasma protein gave a reading of 5.1 g. %, and a hippuric-acid test showed that the amount of hippuric acid passed in the four-hour period was only 2.7922 g., with the maximum excretion in the second hour. It was therefore decided to give 1 g. of choline chloride and 1 g. of cystine twice daily. This was done on the assumption that choline and cystine can help to prevent hepatic cirrhosis due to a very low protein intake.

The hippuric-acid tests were repeated, with the following results :

Date	Amount of Hippuric Acid Passed in 4-hourly Period	Maximum Excretion
12/7/46	2.7504 g.	3rd hour
19/7/46	3.1794 g.	3rd "
30/7/46	3.0220 g.	2nd "
6/8/46	3.0468 g.	2nd "
20/8/46	3.3484 g.	2nd "
29/8/46	3.5882 g.	2nd "

It was therefore concluded that there had been a great improvement in the liver function during this period.

Attempts were also made to raise the low plasma protein level by repeated plasma transfusions, high protein diet, and the addition of "casydrol" by mouth. Plasma protein estimations were as follows : July 2, 1946, 5.1 g. % ; July 12, 5.6 g. % ; July 15, 5.4 g. % ; July 19, 5.5 g. % ; July 29, 6.2 g. % ; Aug. 6, 6.2 g. % ; Aug. 20, 6.8 g. % ; Sept. 2, 6.85 g. % ; Oct. 1, 1947, 7.8 g. % ; Jan. 1, 1948, 8.8 g. %—i.e., 5.2 g. of albumin per 100 ml. of serum and 3.6 g. of globulin per 100 ml. of serum (albumin-globulin ratio = 1:1.5) ; March 9, 1948, 6.25 g. %—i.e., 4.5 g. of albumin per 100 ml. of serum and 1.75 g. of globulin per 100 ml. of serum ; albumin-globulin ratio = 2.6 : 1. It was, however, noticed that although the plasma protein level was 6.8 g. by Aug. 20, 1946, the ascites persisted.

The patient felt very much better, and it was then decided to try to keep the white blood cells within reasonable limits by the administration of urethane by mouth. He was given 1 g. daily for four weeks and the white blood count was recorded daily. After that he was placed on a maintenance dose of 1 g. thrice weekly and weekly blood counts were recorded.

The remarkable fact was that the ascites after tapping took much longer to recur, and when he was tapped on Nov. 12, 1946, 24 pints (13.6 litres) of ascitic fluid were removed. Since that date he has been receiving two "fersolate" tablets three times daily, 1 g. of choline chloride daily, 1 g. of cystine daily, and 1 g. of urethane thrice weekly, with occasional rest periods as determined by blood counts.

He was seen again on March 31, 1948, when he was admitted to hospital for another paracentesis—i.e., after an interval of 16 months. Fourteen pints (7.95 litres) were removed on this occasion, and he remained free from signs and symptoms of ascites until Sept. 4, when he was last seen. His liver was just palpable at the costal margin, whilst his spleen extended only a hand's breadth below the left costal margin. His weight has remained fairly stationary in spite of his excellent appetite, and he feels very fit though his haemoglobin is low.

Since May, 1948, I have administered, in addition to the other medicines, 4 ml. of "anahaemin" intramuscularly at weekly intervals in an attempt to raise the haemoglobin and red blood cells, but so far without success. It has also been necessary to administer the urethane in doses of 1 g. daily to keep the white blood count within reasonable limits.

Discussion

This case is most interesting because when the patient was first seen in April, 1946, it was thought that he would live only a few weeks. The remarkable change which took place in his general condition after the administration of the

cystine and choline chloride was astonishing. His plasma protein rose fairly rapidly, the ascites became less pronounced and required less frequent tapping, his liver became normal in size, and he rapidly put on weight. After the addition of urethane his white blood count was kept within reasonable limits and his spleen rapidly diminished in size.

It is impossible to draw any conclusions from only one case, but I consider that there is a very fruitful field for the treatment of such cases with cystine and choline chloride in combination with urethane.

As pointed out by Cheney (1934), the presence of ascites in cases of chronic myeloid leukaemia seems to be rare, and very few cases have been recorded in the literature. He gives a detailed description of two such cases, and mentions the close resemblance of this condition to Banti's disease. The importance of making a correct diagnosis is stressed, as also is the institution of the appropriate treatment. He considers that the only probable causes of the ascites in these cases are portal thrombosis and low colloidal osmotic pressure of the blood plasma.

I would like to express my deep appreciation to Dr. F. Young for her invaluable help in the pathological investigations in connexion with this case.

REFERENCE

Cheney, G. (1934). *Acta med. scand.*, **81**, 14.

RUPTURE OF THE LIVER ASSOCIATED WITH PARTURITION

BY

JEAN R. C. BURTON-BROWN, M.D., M.R.C.O.G.
F.R.C.S.

Assistant, Nuffield Department of Obstetrics and Gynaecology,
Oxford

AND

JOHN A. SHEPHERD, M.D., F.R.C.S.Ed.

Lecturer in Surgery and First Assistant, Department of Surgery,
University of Liverpool

A case is described in which an unusual complication developed during or after labour, giving rise to an acute abdominal condition. A rupture of the liver associated with haemoperitoneum was found at operation.

Case History

The patient, a 6-para aged 32, was admitted on Sept. 11, 1948, 17 hours after parturition, with severe upper abdominal pain. The onset was sudden, occurring one hour after expulsion of the placenta and after the midwife had left the house. The patient became collapsed and vomited several times. She described the pain as sharp, radiating to the back and right shoulder, and increasing in severity on movement and respiration. All six previous pregnancies and labours were normal. During the last month of the present pregnancy she had suffered from a persistent epigastric pain.

On examination the patient was pallid and weak and was prevented by the pain from moving in bed or sitting up. The temperature was 97° F. (36.1° C.), pulse 112, and blood pressure 110/85. There was generalized abdominal tenderness and guarding, and shifting dullness was detected. The liver was palpable and tender, projecting 1½ in. (3.75 cm.) below the right costal margin. Peristaltic sounds were active. The pelvic organs were normal, although considerable peritoneal tenderness was elicited by palpation of the lateral fornices. The haemoglobin was 51%. A catheter specimen of urine contained 100 mg. of protein per 100 ml.

Since the precise diagnosis was undetermined, it was decided to transfuse the patient with 4 pints (2.27 litres) of blood and keep her under close observation. Heroin in small doses was given periodically.