of drug). The initial smaller dose was to minimize any tendency there might be to a sharp reactive rise of temperature on beginning treatment—a tendency reported in the series of Spink *et al.*

After starting the drug the pyrexia, which had been broadly of an intermittent pattern swinging between limits of 97 and 102° F. (36.1 and 38.9° C.), became in about 24 hours more continuous and reached over 103° F. (39.4° C.); in a further 36 hours it began to fall rapidly and reached normal in about another 24 hours, where it remained. It was still steadily normal 10 days later, when the patient's wish to return home was acceded to (April 22), and on interviewing him on May 12 he reported continued freedom from his former febrile sensations, and a single check on his temperature gave a normal level.

The drug was easily administered and caused no apparent toxic reactions.

I am indebted to Dr. R. E. Jones, of the Public Health Laboratory, Coventry, for serological data and other assistance, and to Dr. T. H. W. Ritchie, of Bedworth, Warwickshire, for clinical information.

J. F. GALPINE, M.D., M.R.C.P.Ed., D.P.H., D.C.H.

Medical Superintendent, Isolation Hospital, Coventry.

REFERENCE

Spink, W. W., Braude, A. I., Castaneda, M. R., and Goytia, R. S. (1948). *J. Amer. med. Ass.*, **138**, 1145.

Arteriovenous Fistula of the Lung

[WITH PHOTOGRAVURE PLATE]

Until recently arteriovenous fistula of the lung has only rarely been reported. It was thought worth while, therefore, to record the following case, with details of the investigations made to establish the diagnosis. The question of the advisability of surgical treatment in this case is discussed.

CASE REPORT

The patient, a man aged 36, first came under medical care in August, 1947, complaining of a dull ache in the right upper chest which he attributed to strenuous and unaccustomed exercise taken some two weeks before. There was a history of slight morning cough for some years, with an occasional trace of mucoid sputum, in which he had noted streaks of blood once only, on the day before he was first seen. This streaking has since been observed on another occasion. No dyspnoea or other symptoms have been present. His previous history was not relevant, and on close questioning epistaxis or other evidence of a bleeding diathesis was not admitted. Goldman (1948), in a review of the literature, has stressed the hereditary factors in the aetiology of the condition, but in this case the family history, so far as it could be obtained, was also negative.

Physical examination did not reveal any cyanosis or finger-clubbing, and no abnormal physical signs were found in the respiratory system. The pulse was regular at 80 per minute and was not collapsing in character. There was no evidence of enlargement of the heart or of cardiac murmurs. The blood pressure was 135/75 mm. Hg in both arms. A soft murmur, heard only in systole, was present over a roughly circular area about 6 in. (15 cm.) in diameter in the lower part of the right axilla. The murmur was faint but could be well heard with the patient lying on his left side and extended over a pillow. It was not appreciably affected by respiration. Careful examination of the skin and mucous membranes revealed no evidence of other haemangiomas or telangiectases. Other systems were normal.

The diagnosis was chiefly established by radiography. The postero-anterior film (Plate, Fig. 1) showed a mottled opacity in the lower zone of the right lung which on the lateral film overlay the lower part of the heart shadow. Between the opacity and the vascular markings of the right hilum was a longitudinal shadow which on tomography (Fig. 2) could be clearly identified as consisting of two main parts representing

branches of the pulmonary artery and vein. On screening, opinion was divided on whether the shadow pulsated, but it was noted in a bronchogram (Fig. 3) that, whereas nearly all the bronchial shadows, including those in juxtaposition to the heart, were present and clearly defined, those close to the main opacity were blurred.

Mr. R. C. Brock kindly arranged for angiography at Guy's Hospital, and the resulting radiograph (Fig. 4) leaves no doubt regarding the vascular nature of the lesion. Mr. Brock also reported a normal bronchial tree on bronchoscopy.

Blood examination revealed: red cells, 6,100,000 per c.mm.; haemoglobin (Haldane), 110%; white cells, 6,000 per c.mm., with a normal differential count. Oxygen saturation was found to be 89% in a specimen of arterial blood. The sedimentation rate was 2 mm. in one hour (Westergren). Culture of sputum revealed a mixed growth of normal pulmonary flora. The vital capacity was 4,500 c.cm. The electrocardiogram (three standard leads) was normal in all respects.

The case has now been observed over a period of seven months without any change in the clinical or radiological picture.

DISCUSSION

On the data available the lesion must be regarded as a vascular abnormality or tumour in the substance of the lung, with a minimal pulmonary arteriovenous shunt but no evidence of similar lesions or telangiectasia elsewhere, though the presence of such lesions cannot be excluded *ante mortem*. In view of the minimal shunt in this case the possibility of the systemic (bronchial) circulation being implicated, as suggested by Maier *et al.* (1948), must be considered. The main practical problem is whether surgical removal should be advised. On the one hand we are dealing with a virtually symptomless lesion with very little disturbance of the normal physiology. On the other there is the danger of haemorrhage and the possible extension of the lesion with an increase in the shunt or even malignant change. At the moment there would appear to be no general contraindications to surgery, but from the few reported cases it would seem that operative removal is not without risk, particularly where there is a large pulmonary arteriovenous shunt. There is also no certainty that other lesions are not present.

Surgical removal has been advised but so far refused, so that pathological data on the nature of the lesion are not available.

I would like to thank Dr. F. H. Young, physician to the Brompton Hospital, for permission to publish this case and for his helpful criticism; Dr. F. Prime for performing the oxygen saturation test; and Dr. L. G. Blair for the skiagrams.

F. D. BEDDARD, M.B., B.S.,
late A.R.M.O., Brompton Hospital.

REFERENCES

Goldman, A. (1948). *Amer. Rev. Tuberc.*, **57**, 266.
Maier, H. C., *et al.* (1948). *J. thorac. Surg.*, **17**, 13.

At a meeting of the Professional and Technical Council B of the Whitley Councils for the Health Services on April 20 the following agreement was reached on the salaries of medical laboratory technicians employed in the National Health Service in England, Wales, and Scotland. *Student and Junior Technician*: Aged 16, £110 per annum; 17, £125; 18, £145; 19, £165; 20, £195; 21, £225; 22, £240 10s.; 23, £260; 24, £279 10s.; 25 and over, £299. On passing the Intermediate Examination of the Institute of Medical Laboratory Technology (or equivalent qualification), or the Inter. B.Sc., the salary is increased by £13 per annum at any point on the scale. If no examinations are passed, the salary is increased to £312 per annum at age 30. *Technician*: £370×£15=£435. On gaining the Fellowship of the Institute of Medical Laboratory Technology (or equivalent qualification) the salary is increased by £15 at any point on the scale. *Senior Technician*: £450×£20=£530. *Chief Technician*: £530×£20=£650. London weighting: In the Metropolitan Police Area all scales are subject to London weighting on the following basis: Age 16-20, £10; 21-25, £20; 26 and over, £30. The new scales operate from April 1, 1949.

G. RIDDELL ROYSTON: ACUTE PNEUMONITIS IN A BERYLLIUM-WORKER

FIG. 1.—Soft mottling. (June 27, 1947.)

FIG. 2.—Apparently typical miliary tuberculosis. (July 11, 1947.)

FIG. 3.—Considerable resolution. (Aug. 7, 1947.)

F. D. BEDDARD: ARTERIOVENOUS FISTULA OF THE LUNG

FIG. 1.—Postero-anterior view.

FIG. 2.—Tomogram.

FIG. 3.—Bronchogram.

FIG. 4.—Angiogram.

K. DAMODARAN: INFECTIVE HEPATITIS AND PORTAL CIRRHOSIS

FIG. 1.—Liver $\times 300$. Marked cellular infiltration. Second week.

FIG. 2.—Glomerulus $\times 400$. Showing pigment granules. Second week.

FIG. 3.—Liver $\times 250$. Post-arsenical hepatitis. Cirrhosis and round-cell infiltration.

FIG. 4.—Liver $\times 240$. Portal cirrhosis, showing round-cell infiltration.