intravenous injection of soluble antigens (horse serum) into animals, while somatic antigens (whole bacteria) produce their effect mainly in the lungs, liver, and spleen (Ehrlich *et al.*, 1949; Gregory and Rich, 1946). Most of these are regions commonly involved in Still's disease, and, as has been shown, the lungs, the heart, and the liver may not escape.

Conclusions

Study of the most acute cases of Still's disease emphasizes the widespread nature of the lesions. Although the majority of cases proceed to progressive arthritis, some patients may recover completely. Characteristic types of pyrexia can be recognized at different stages, and significant effects on the blood picture have also been observed. In rare instances the clinical picture may change from Still's disease to leukaemia. The close resemblance of Still's disease, particularly at the onset, with certain rare syndromes of an allergic nature suggests that rheumatoid arthritis may have a similar aetiological basis. Should this be so, some form of primary bacterial invasion seems to be likely in the course of events.

I am indebted to several of my medical colleagues for allowing me to study their cases, and to Dr. Bodian and Mr. Martin for the histological sections and microphotographs reproduced in the photogravure plate.

REFERENCES

- Bland, E. F., and Jones, T. Duckett (1935). *J. clin. Invest.*, **14**, 633.
 Blechmann, G., and Carcanagues, P. (1940). *J. Méd. Paris*, **60**, 254.
 Cohen, H. (1948). *Textbook of the Rheumatic Diseases*, ed. by W. S. C. Copeman, p. 182. Livingstone, Edinburgh.
 Cohen, S. Solis (1913). *Trans. Ass. Amer. Phys.*, **28**, 739.
 Collins, D. H. (1937). *Rep. chron. rheum. Dis.*, **3**, 49.
 Ehrlich, W. E., Seifter, J., and Foreman, C. (1949). *J. exp. Med.*, **89**, 23.
 Ellman, P., and Ball, R. E. (1948). *British Medical Journal*, **2**, 816.
 Felty, A. R. (1924). *Bull. Johns Hopk. Hosp.*, **35**, 16.
 Garrod, A. E. (1910). *Quart. J. Med.*, **3**, 207.
 Gregory, J. E., and Rich, A. R. (1946). *Bull. Johns Hopk. Hosp.*, **78**, 1.
 Hench, P. S. (1949). *Proc. Mayo Clin.*, **24**, 167.
 — and Rosenberg, E. F. (1944). *Arch. intern. Med.*, **73**, 293.
 Humphreys, E. M. (1948). *Ann. intern. Med.*, **28**, 12.
 Imrie, A. H., and Aitkenhead, A. C. (1939). *Lancet*, **2**, 421.
 Kahlmeter, G. (1939). *Acta med. scand.*, **102**, 432.
 Klemperer, P. (1948). *Ann. intern. Med.*, **28**, 1.
 Schlesinger, B., and MacCarthy, D. (1949). *Diseases of Children*, ed. by D. Paterson and A. Moncrieff, **2**, p. 686. Arnold, London.
 Schlesinger, H. (1899). *Mitt. Grenzgeb. Med. Chir.*, **5**, 441.
 Spühler, O. (1941). *Schweiz. med. Wschr.*, **71**, 25.
 Türk, E. (1938). *Arch. Kinderheilk.*, **114**, 65.
 Weber, F. Parkes (1946). *Lancet*, **2**, 931.

The current issue of *Planning* is the three-hundredth in the series which PEP began some 17 years ago. It looks back on the early history of PEP between its foundation in 1931 and the outbreak of war. A brief account of its origin mentions the men who were most concerned with the development of the group. In the early days PEP was given some special grants by the Rockefeller Foundation, but over the period 1933–9, 12 major and two minor reports and more than 150 broadsheets were published on a budget of no more than £56,000 for these seven years. There is a brief description of the *ad hoc* working groups set up by PEP, and an attempt is made to assess its contribution to contemporary knowledge and ideas. The first major report to be prepared by the Civic Division of PEP was that on *The British Social Services*, and this was followed in December, 1937, by the *Report on the British Health Services*. Three groups which did important work during the war were the Social Security Group, the reconstituted Health Group, and the Population Policies Group. The report on *Population Policy in Great Britain*, however, was not published until April, 1948.

PSYCHOGENIC RHEUMATISM

BY

WILLIAM TEGNER, M.R.C.P.

DESMOND O'NEILL, M.D., M.R.C.P., D.P.M.

AND

ANNA KALDEGG, Ph.D., B.A.

(From London and Guy's Hospitals)

In a large hospital to which patients are referred suffering from "rheumatism" considerable diagnostic experience is needed in order to classify the various conditions included in this omnibus term. There is no agreement at present on the scope of the term; it is customarily taken to indicate a group of painful disorders of the locomotor system. It is obvious that a large number of syndromes come under this heading. While the arthritides, neuralgias, and other somatic conditions as a rule present straightforward signs and symptoms, patients are continually encountered whose experiences do not seem to fit in with any recognized syndrome and in whom physical signs are absent or minimal. In such cases one may be tempted to regard the patient as suffering from pain which is primarily of psychogenic rather than organic origin. If this hypothesis is correct, it is plain that rheumatic disorders of this type call for particular care in diagnosis and for treatment more far-reaching than analgesics and physiotherapy.

In recent years there has been a growing disinclination to accept all rheumatic complaints as somatic in origin. Even before the war one of us had coined the term "polyalgia syndrome" to describe the symptoms of which some patients complained. In England, Ellman *et al.* (1942) studied 50 patients with "fibrositis": 23 had no physical signs, although they had been diagnosed as suffering from somatic disease; 25 showed hysterical tendencies. An interpretation of the rheumatic complaint in psychological terms was given by these workers. An important contribution was made by Flind and Barber (1945); many of the subjects investigated by them were found to be suffering from psychiatric rather than organic states. Hench and Boland (1946) gave a full description of psychogenic rheumatism in a paper on the incidence of rheumatic disease in the United States Army; this condition was the third most common diagnosis made in their series of 800 patients.

If psychogenic rheumatism is to be accepted as an entity both by those who diagnose and treat rheumatic disease and by psychiatrists, our concept of it needs clarification and acceptable diagnostic criteria must be established. This demands close collaboration between clinician and psychiatrist: it is the purpose of this paper to show the results of such an endeavour.

Fifteen patients regarded by the clinician as suffering from psychogenic rheumatism were referred for detailed psychiatric examination. In each case the patient was first examined by routine clinical methods. The haemoglobin and sedimentation rate were estimated. The group contained 14 women and one man; the distribution by age was: 15–29, four; 30–39, three; 40–49, four; 50 and over, four. The preponderance of women in this group was in accord with the assumption that psychogenic rheumatism in clinical practice is considerably commoner in women. A control group of patients suffering from a painful disorder of which the cause was known was investigated by the same means. This group consisted of subjects of the same age and sex distribution, and broadly of the same social class. The diagnoses were: osteoarthritis (3),

brachial neuralgia (3), fracture of a joint (3), cervical rib (1), peri-arthritis (1), lesions of an intervertebral disk (2), rheumatoid arthritis (1), alveolar abscess (1). The psychiatric examination was conducted along three main lines—clinical interview, psychometric examination by two projective methods (Rorschach test and Thematic Apperception test—T.A.T.), and appraisal of information given by a relative, employer, or friend.

Results

A. Form of Rheumatic Complaint

Expression of Complaint.—Brief and consistent terms ("sharp," "sore," "aching"): controls, 10; patients, 3. Compound or unusual terms ("twisting," "heaviness," "burning," "stinging and stabbing," "shooting and numbness," "terrible," "hard to describe"): controls, 5; patients, 12. The patients often seemed to be trying to put into words an indescribable sensation. The impression to the observer is frequently that of a mystical or ecstatic experience, with an undercurrent of satisfaction, which recalls Helene Deutsch's (1947) elaboration of masochism as a primary feminine trait. Terms may be used which convey the description of a mental state rather than of a physical stimulus—e.g., "irritation." The psychological precipitants are at times stated quite plainly—e.g., "boredom causes irritation."

Site of Discomfort.—Localized: controls, 11; patients, 3. Widespread: controls, 4; patients, 12. In the majority of the control group discomfort was confined to the area of the lesion or to an area related to it. The distribution of discomfort in the test group was commonly very extensive—e.g., neck, spine, knees, shoulders, calves, hands. The phrase "pain all over" was employed by several patients.

Factors Influencing Severity of Discomfort.—In the controls the most common aggravating factor was touch or movement of the affected part: 11 cases. Among the patients the following were mentioned: work, 4; relaxation, 4; mood, 2; noise, 1; heat or cold, 2; doubtful, 2. Relief of distress in the control group followed rest in 10 cases and treatment in four. Of the patients, seven did not admit relief with any activity or form of treatment; in the remainder some relief occurred with rest or exercise, a hot bath, or quiet. Analgesic drugs produced improvement in six out of seven control subjects in which they were tried; out of eight patients, none noticed any effect from these drugs. Two control subjects considered that the complaint was influenced by their state of feeling, as compared with six patients. Variations in the severity of discomfort throughout the 24 hours, or from day to day, showed no great differences between the groups. The effect of climatic conditions—cold and damp—was similar in both groups. Degrees of disability from slight to total were noted in six control subjects; only one patient suffered any disability, and this was slight.

Associated Symptoms.—Depression, tension, irritability, poor sleep, or phobias were present in 14 of the patients and in five of the controls. In two of the latter the nervous symptoms were considered to be secondary to the somatic disorder.

B. Circumstances of Onset

Two of the controls and 13 patients became aware of the onset of the disorder in circumstances associated with emotional tension. In the remaining 13 controls the onset of illness appeared to be fortuitous. In the majority of the test group the rheumatic condition could be shown to have a comprehensible meaning in relation to the patient and her life situation.

C. Psychiatric Status

Clinical Examination.—In 10 of the control group no significant psychiatric abnormality was found. In the other five the physical condition was accompanied by nervous symptoms:

The first of these subjects complained of worries, fears, tension, and fatigue. It was considered that her conditions of housing were producing anxiety and frustration, which increased the disability due to her organic state (osteoarthritis). In the second, depression, lack of energy, and poor sleep appeared about one month after the onset of brachial neuralgia and seemed to be a consequence of pain and limitation of activity.

In the third, a right-sided cervical rib had begun to cause symptoms a few days after a fall on the right hand. On examination the patient also admitted to shakiness, depression, disturbing dreams, and fear of the dark and of open spaces. These were discovered to accompany sexual maladjustment with dyspareunia and fear of pregnancy. They appeared unrelated to the rheumatic complaint.

In the fourth subject pain from peri-arthritis of the right shoulder was accompanied by fatigue, headache, and insomnia. The shoulder condition followed a fall a few months before the time of examination; the other symptoms had been present over many years and were the expression of a lifelong neurotic state.

The fifth subject, with a twelve-months history of rheumatoid arthritis, complained of lassitude, depression, and disturbed sleep. She gave a history of previous nervous illness. In view of the common observation that rheumatoid arthritis may become manifest after emotional trauma, the circumstances of onset in this patient were carefully reviewed. No significant precipitating event was found, and the nervous symptoms were considered to be secondary to the arthritis.

In only one patient of the test group were there no associated nervous symptoms. With two possible exceptions the rheumatic condition in every patient in this group formed one component of a reaction pattern which was also expressed in disturbance of affect and behaviour. The pathogenesis of the condition in each patient is considered in more detail below.

Psychometric Findings.—(a) *Rorschach test*: This is a sensitive indicator of personality disturbance. Interpretation of records is a skilled task and cannot be reduced to a quantitative scoring method. For the purposes of this investigation the accompanying table of incidence of certain Rorschach signs of maladjustment was drawn up; it shows a greater degree of disturbance in the test group than in the control group taken as a whole. If the two control subjects with neurotic features on clinical examina-

Results of Rorschach Test

	Responses < 15 or > 70	No Colour Response	Colour Response C Only	Rejections Other Than IX	Last 3 Cards			Content Categories Under 5	P Responses Under 4
					Under 30%	Above 45%	Total		
Controls	4	4	—	5	6	—	6	6	7
Controls, without neuro- tic subjects	2	2	—	4	6	—	6	4	5
Patients	11	6	2	8	8	2	10	10	11

tion are excluded the difference becomes more pronounced. More patients than controls showed evidence of difficulties in the emotional field: domination by emotional impulses or a morbid fear of and withdrawal from situations likely to arouse them. "Blocking"—complete failure to handle adequately such situations—was more often found. Awareness of alien forces within the personality of a sinister and threatening nature was commoner in the patient group.

Signs of sexual maladjustment occurred more often. The various effects of anxiety on the test performance were shown to a considerably greater extent in the controls than in the patients' records.

(b) *Thematic Apperception Test* (selected series of cards): The patients showed in general a greater preoccupation with themes of violence, injury, crime and punishment, murder, and suicide. For example, one at least of the cards was interpreted as showing a murder by six of the patients, as against three controls; as showing a suicide by four patients, as against one control; and as showing some form of violence by five patients, as against one control. Themes of guilt after wrongdoing were also more frequent in the patients' records.

The material from the Rorschach and T.A.T. records will be used at a later date to build up a formulation for the pathogenesis of illness in the test group.

D. Physical Status

The control group showed the physical signs appropriate to the physical disorder from which they were suffering. The complaint was in general consistent in its expression and the discomfort was localized to one, usually anatomical, body area.

The test group complained in the main of diffuse and severe discomfort, inconstant in site, in degree, and in nature. The terminology used to express the complaint was often bizarre. The distribution of the discomfort did not correspond to that of a recognized physical disorder. Physical signs of a skeletal abnormality were absent. To say, however, that no physical signs can be elicited is not usually correct. A typical reaction to examination was found. Tenderness was often diffuse and due to skin hyperaesthesia. The patients found relaxation difficult, and muscular tension had often to be overcome before full joint movement could be demonstrated. Deep reflexes were exaggerated. In several cases the patient took the examiner's hand and applied it to the painful portions of her body.

X-ray examination and sedimentation rate estimation in the test group showed no abnormality.

Pathogenesis

The absence of physical signs of organic disease and the clinical characters of the complaint, especially the distribution of the discomfort and the significance of the complaint in the patient's total reaction, place this type of rheumatism in a class apart from the hitherto recognized rheumatic disorders. Its formulation as a "psychosomatic" condition cannot, however, be considered as adequate unless the morbid physiology underlying the physical symptoms can be articulated with the psychodynamics of the patient's emotional state.

In the test group a tentative formulation of the pathogenesis could be arrived at in 13 of the 15 cases. A fuller account embodying the psychometric material and further physiological data will be published later; a summary of each case, necessarily simplified, is here given.

Miss A., aged 24, is described as "fussy, upright, proud; holds herself very upright." Mother: rheumatism; nervy, quick-tempered. Mother died in 1943; in 1945 father began to carry on a liaison with another woman, who drank. Patient objected to this, and there were quarrels, after which she felt very disturbed and tense. Her feelings "seem to bottle up inside." Onset of generalized pains, mainly neck and spine, in 1945. Pathogenesis: somatic expression, perhaps in muscular tension, of strong moral resentment of father's behaviour.

Mrs. B., aged 50, was nervous and timid in childhood; illness diagnosed as chorea. Nail-biting and enuresis until 11 years. Home unhappy. Marriage unsatisfactory; husband has "bad nerves" and accuses her constantly of infidelity. "Prolapse of womb" in 1946; after this she was convinced that "something was wrong inside" which had not been discovered. Pain in the right side and back began at this time, accompanied by hysterical outbursts. Pathogenesis: nosophobia.

Mrs. C., aged 45. Father: violent, unstable. Mother: nervy. As a child patient was timid and nervous and afraid of father. Marriage happy; wanted large family but had only one child. Operation at 30 for sterility; at 42, uterus and one ovary removed. Very distressed by this; felt inferior to other women. Backache began after the operation. Rorschach: gross disturbance by a sinister fantasy life, with aggressive and sadistic trends. Pathogenesis: somatic expression of self-punitive trend at her failure to bear children.

Mrs. D., aged 39, is an unstable, dull, shallow person who nevertheless has considerable determination and can usually get her way with people. Married during the war and left her husband next morning as she could not tolerate sex relations. Felt strongly the need for home and security. Onset of pain in back and perineum while living alone in London. This resisted all treatment until she found a congenial job with friendly people. Pathogenesis: somatic expression of intense self-concern and need for affection.

Miss E., aged 27. Father: rheumatism; bad temper. Mother: rheumatism; irritable. Patient a diligent, conscientious, active person. Is compelled by circumstance to do a good deal of housework as well as her job, and resents it. Both parents are difficult and obstinate, and there are continual family quarrels. Onset of generalized pains after an increase in household duties on her sister's marriage. Pathogenesis: somatic expression of suppressed resentment of her parents' attitude and demands.

Mrs. F., aged 42, was nervous and timid as a child; enuresis till 12 years. Regular dysmenorrhoea; after marriage, disinclination for sex. Onset of pain in rectum while on evacuation from London during the war; this was relieved by injection for haemorrhoids, but she was told that the "neck of the womb was out of order." Became convinced that she had some disease of the uterus, probably cancer; pain in rectum returned, with backache. Pathogenesis: nosophobia.

Mrs. G., aged 53. Father: rheumatism "all over"; irritable. Mother: rheumatism. Patient nervous and worrying as a child. Family unhappy. All her life has had to look after invalids; her husband died of tuberculosis after a long illness. Had one child which lived only two weeks; pain in the back at this time. Had to take care of a sister with Parkinsonism who did not appreciate what was done for her; exacerbation of rheumatism at this time. Depressed; considers her life is empty. Pathogenesis: somatic expression of anxiety with guilt over her aggressive impulses towards sister.

Mr. H., aged 23, was wayward and delinquent as a boy; preferred sport to work. Served in R.A.M.C. during the war, much of the time abroad. Found this "nerve-racking"; longed for home. Onset of pain in left chest during a football match, probably from a minor injury, in Singapore after return from home leave. After demobilization, dissatisfied with civilian job and living conditions. Pathogenesis: prolongation of physical symptom due to frustration and discontent.

Mrs. I., aged 42. Mother: rheumatism; very concerned over children's health. Patient is married and has four children, who are a constant worry to her lest they fall ill or are injured. Never lets them out of her sight, but is impatient if they resist her—"would like to take a knife to them." Onset of backache during a depressive illness. T.A.T.: preoccupation with death and suicide. Pathogenesis: somatic expression of tension from markedly ambivalent feelings towards children.

Mrs. J., aged 33. Father: backache, nervous. Mother: "curved spine." Patient always wished to do well in business and was happiest when managing a public house. Has always been irritable, tense, and unable to relax. Marriage fairly satisfactory, but she had dysmenorrhoea, distaste for sex relations, delayed conception, and vomiting in pregnancy. She sleeps badly, and while half awake sees vivid pictures of death

and injury. T.A.T.: preoccupation with and fear of sadistic impulses. Onset of pain in back, legs, and hands during last pregnancy. Pathogenesis: somatic expression of discontent from thwarted masculine aims and ambitions.

Mrs. K., aged 32. Father: rheumatism in head, arms, shoulders, knee; "always rubbing himself"; "nerves in the head." Patient "takes after father very much." Onset of aching pain in back, arms, and legs while living away from London during the war. No immediate source of emotional tension found. Patient very preoccupied with her complaint and sure she is going to be crippled with rheumatism, as her father and aunt were. Probable pathogenesis: nosophobia.

Miss L., aged 19. Father died when patient was 11. Brought up by mother, with whom she has always got on badly. Patient described as "hysterical; flares up easily." Both she and her mother have violent tempers. Patient says she wants to leave home, yet is clearly very dependent on mother. Has one boy friend, with whom she quarrels continually. They became engaged, hoping that this would ease things; the quarrels ceased, but the patient developed pain in shoulders and arms, with odd sensations in the limbs. Pathogenesis: somatic expression of anxiety from two main conflicts—over dependence, and over relation to fiancé; with utilization of illness as a means of avoiding marriage.

Mrs. M., aged 52. Father: ill-tempered and unstable; left home. Patient strong and active; reared in a convent. Marriage difficult; husband a chronic invalid who died, leaving her with five young children. She had to work to support the family. Onset of pain in back and limbs after a confinement in her early thirties. Repeated mild attacks of depression, ascribed to worry of looking after children. T.A.T.: marked preoccupation with theme of injury and reparation as between mother and child. Probable pathogenesis: somatic expression of guilt over care of children.

Miss N., aged 59. Father: nervous, temperamental. Mother: rheumatism. Patient has energetic and forceful disposition, with strong intellectual and social ambitions, which have never been realized. At about 24 she began to suffer from insomnia with "visions," in which she could see herself as a crowned queen; with this, she felt irritation under the skin. At about 50 pain began in knees, ankles, and fingers, worst during the night, when she cannot sleep. T.A.T.: preoccupation with and fear of sadistic impulses. Pathogenesis: somatic expression of suppressed resentment from thwarted ambition, precipitated by menopause.

Miss O., aged 43. Mother: fainting attacks; chronic fatigue; dominating and strict with her children. Patient active, vigorous, ambitious, and successful; reached her goal—to be a head mistress. Onset of backache during the war, at a time of especial stress while she was concerned over danger to her mother from the air raids. Had to give up her career to look after mother because of illness. Pathogenesis: somatic expression of anxiety from conflict over relation to mother.

Diagnosis and Treatment

The principal diagnostic features of psychogenic rheumatism appear to be in (1) expression of complaint; (2) area of discomfort; (3) failure to conform to an established organic syndrome; (4) setting of the illness in relation to the patient's life situation; and (5) lack of response to physical treatments.

The treatment and disposal of patients regarded as suffering from psychogenic rheumatism has always provided a real problem to the clinician. It is part of the syndrome that they do not respond to treatment which is limited to physical measures. It seems clear that to these patients their suffering is something very personal and valuable; it makes them important, interesting, and worthy of sympathy, and is a protection against an unkind world. To treat them with passive physiotherapy is to acquiesce in their own diagnosis of disease and to allow it to take a deeper root. Such treatment usually gives temporary relief but does not cure; after a time it may come to be

indispensable to the patient. Physiotherapy which is painful may remove the symptom; another symptom will then take its place. Injections of a local analgesic may be tried; they generally have a dramatic effect. Even as little as a few drops of 1% procaine may lead to fainting and hysterical manifestations.

One of us has attempted to treat the patients in groups by general ultra-violet irradiation followed by exercises; this method has had some success. Group psychotherapy of selected patients has now been started and has already given encouraging results. It is probable that an approach along these lines will prove to be the most profitable measure.

Summary

An investigation by a physician, a psychiatrist, and a psychologist of a group of patients suffering from a rheumatic disorder without signs of organic disease is described; a comparison with a control group is made.

In the former group the disorder was found to have certain special characters; in most cases the onset and course of the condition could be correlated with emotional disturbance. The condition seems to merit recognition as a clinical entity: it may be described by the term "psychogenic rheumatism."

We are indebted to Dr. T. A. Munro for his co-operation and advice; to Dr. K. W. N. Palmer, of the Department of Physical Medicine at Guy's, for his assistance with the control patients; and to Professor Aubrey Lewis and Dr. J. L. F. Maine for helpful suggestions.

REFERENCES

- Deutsch, Helene (1947). *Psychology of Women*. London.
Ellman, P., Savage, O. A., Witkower, E., and Rodger, T. F. (1942). *Ann. rheum. Dis.*, 3, 56.
Flind, J., and Barber, H. S. (1945). *Quart. J. Med.*, n.s. 14, 57.
Hench, P. S., and Bolland, E. W. (1946). *Ann. intern. Med.*, 24, 808.

A CASE OF RHEUMATOID ARTHRITIS SIMULATING PYOGENIC INFECTION

BY

D. S. SHORT, M.D., M.R.C.P.

Medical Registrar, Southmead Hospital, Bristol

The rarity of high fever in rheumatoid arthritis may be judged from the fact that Bannatyne (1896) on one occasion recorded a temperature of 105° F. (40.6° C.). This observation was later quoted by Garrod (1907) and by Jones (1909) as still unrivalled. Modern authors confine themselves to the statement that the temperature may rise to 102–103° F. (38.9–39.4° C.) and that a slight pyrexia may persist for several weeks.

The highest leucocytosis recorded in an uncomplicated case appears to be the figure of 15,800 mentioned by Erving (1903). Most modern writers omit all reference to the possible occurrence of a marked leucocytosis, and Gibson (1948) declares that "an increase of over 12,000 leucocytes per c.mm. is rare."

The following case is more reminiscent of the disease as seen in children, when fever and general symptoms may be present for weeks or months before definite articular manifestations make their appearance and fever and leucocytosis may be so marked as to simulate a pyogenic infection or leukaemia.

Case Report

The patient, a clerk aged 26, had previously been healthy save for a succession of sore throats.