BRITISH MEDICAL JOURNAL

LONDON SATURDAY JULY 30 1949

PROBLEMS OF AGEING AND CHRONIC SICKNESS*

BY

A. P. THOMSON, M.C., M.D., F.R.C.P.

Professor of Therapeutics, University of Birmingham; Physician, United Hospitals, Birmingham

The Problem

In the short span of 150 years the progress of medicine has been chiefly responsible for two distinct problems of population in Western Europe. In Great Britain, for instance, during the first part of that period the reduction of infant mortality and the control of infectious disease led to a rapid and embarrassing increase in the number of youthful inhabitants—from about nine millions in 1801 to forty millions at the close of the nineteenth century. Decline of the birth rate and emigration checked the increase, but the astonishing advance of curative medicine in the past 50 years has so prolonged the average duration of life that the total population is to-day larger by ten millions, and of the fifty millions in this country now no fewer than seven millions are over 60 years of age. As it is inevitable that in the immediate future the proportion of the aged will increase, and that the survival of individuals with chronic maladies will be greater, the urgency of the problem of the proper provision for the chronic sick and aged needs no emphasis. It has not indeed been neglected. Within the past two years this College (1948) and the British Medical Association (1947, 1948) have published reports from special committees on the care and treatment of the elderly and infirm, and in addition there have been a number of papers from individuals whose work lies mainly in this field. The relevant literature is interesting and abounds with suggestions for reform. There is a general demand for better classification of the patients and for their segregation, like with like, in appropriate institutions; the need for special housing for the old is emphasized, and a plea is made for the establishment of yet another medical specialty in the shape of geriatrics.

When I sought, however, for some factual and statistical analysis of the patients now in the institutions and for precise information about how they got there and why they stayed so long I often sought in vain, though Affleck (1947) in his psychiatric approach to the problem included some valuable figures, to which I will refer later. Lack of dispassionate statistical analysis is a weakness in these reports, and as there is nothing in the literature from the hospitals for the aged and chronic sick that compares, for instance, with Sheldon's (1948) account of the condition of the old people living in ordinary homes in Wolverhampton I

decided to undertake an independent investigation; in addition it seemed probable that conditions would vary in different parts of the country.

The Position in the Midlands

In the Birmingham Hospital Region, which includes five counties with a population of 4.2 millions, there are 50 institutions concerned with the reception of the aged and chronic sick. Their variety is remarkable; most of them are small and understaffed, with old and dingy buildings and poor equipment. Some are parts of acute general hospitals; none have been well designed for their present use, and the concern expressed by the Nuffield Surveyors for the fate of their patients was amply justified. Most of these institutions have other functions, of which the most important is the accommodation of poor and homeless people who, though not sick, are in general unable to fend for themselves.

The number of beds available (apart from those for the insane and simply destitute) in these 50 institutions is 5,780, and of these no fewer than 2,400 are contained in five large establishments in densely populated areas. For comparison the total available for general medical and surgical use in the acute hospitals is only 5,960, but in addition there are about 1,000 beds for the same purpose in the teaching hospital. Briefly, of every 100 beds for general medicine and surgery in the region at least 45 are reserved for the chronic sick. This large proportion was to me surprising.

Dr. Richard Padley, of the University Department of Medical Statistics and Statistical Officer to the Regional Board, to whom I am indebted for these figures, also made an estimate of the staff employed in caring for the chronic sick (excluding medical and laboratory services), and found at the end of 1948 that 3,650 persons were so engaged. In other words 100 chronic beds required on average a whole-time staff of 61, made up as follows: trained nurses, 4; assistant or other nurses, 21; domestic workers, 22; laundry, 4; maintenance, 5; administration, 5. These are serious figures; with an ageing population it is obvious that the staffs necessary for institutions of this kind may soon become a serious drain on the national reserve of man-power.

The field to be explored was so extensive and the problems were to me so novel and perplexing that it seemed

*The first of two Lumleian Lectures delivered before the Royal College of Physicians of London on April 5 and 7.

impossible to investigate all the institutions closely, and I decided to concentrate attention on the largest—the Western Road Infirmary in Birmingham—and to be content with occasional visits to others.

First Impressions of Western Road

My most vivid impression of the chronic wards of this hospital is of an atmosphere of profound apathy. The patients seldom spoke to each other; they rarely moved, and although there was a library service and newspapers were provided it was exceptional to see one reading. At first we thought that this unnatural calm was a consequence of discipline imposed in preparation for our visits, but we were assured it was not so and that there was nothing unusual in it. Even when we were hours together in a ward and examined in turn every patient in it the prepotent sense of still foreboding and quiet resignation endured.

Contact with the patients was not difficult; histories of their social background and infirmities were mostly accurate and readily given, but what made them remarkable was the conspicuous absence of emotional overtones as they recounted them. There was no bewailing of their fate, no clamant sense of grievance, and they spoke of life's adventure with the detachment of a disembodied observer rather than with the passion of a participant. One old lady told us a strange story of many years of colourful life breeding and breaking ponies in the Welsh Marches, of visits to country fairs to sell them, of strange adventures in wild country, and of independent freedom rarely to be attained in England now. An accident crippled her and she had been long in bed. Her face was immobile as she spoke; the glory and the zest had departed; she was interested mainly in her tea, which was a little late. Food, indeed, was their major interest; it was astonishing how much they ate. Nearly all of them were well nourished, and most of them were stout. The wards came momentarily to life at meal-times.

Apathy prevailed also in the large rooms for those who were able to dress and move about: they sat listlessly and in silence; the men smoked as long as their tobacco lasted. In these rooms there was a striking difference between the sexes: the women were much more careful of their personal appearance; they were cleaner, their clothes were neater, and they were more readily responsive when addressed. Nearly all the women had capacious bags in which they collected a strange and pathetic assortment of trifles of little value—fragments of food, bits of cloth, and empty boxes. Both rooms emptied several minutes before a meal, and long queues were formed at the doors of the dining-hall, in which there was strict segregation of the sexes; even the married couples did not sit together, and they rarely expressed any wish to do so.

We were warned that the prevailing calm was sometimes broken by outbursts of petulant misbehaviour which were commonly the result of childish jealousy, resentment, or misdemeanour. Pilfering was common and led to trouble. The manner of the theft was often ingenious and astute. One man coveted the pipe that belonged to another who also had half a crown. The pilferer, although almost incapable of movement, got up one night and stole the half-crown. A little later he bought the pipe with the money he had stolen from its owner. The men who were up and about had another interest besides food—they gambled. Nearly every morning there was an hour of feverish activity in which they studied the racing information and arranged their bets for the day. The results they

heard on the radio. Football pools were not neglected. Officially their financial resources were limited to five shillings weekly allowed to them out of their old age pensions, but a few earned small sums in addition by doing odd jobs. They had considerable freedom, and when fit enough and in the humour they went into the city on social occasions; not rarely they were drunk on their return. Some of the more adroit deposited their old age pension books with friends outside the institution and managed to retain the whole sum by drawing it when they went visiting. One old woman had defeated the regulations of authority in this manner for over two years; she had glorious "blinds" on the proceeds and was a source of considerable trouble. In the acute wards the atmosphere was like that of an ordinary general hospital. Most of the patients had been admitted from their homes but a short time before; they were interested, and still hopeful of recovery sufficient to allow them to gather up the threads of their customary way of life. It was here alone that we saw any evidence of anxiety; in the chronic wards there was the apathetic indifference I have described and in others there was frank psychosis, but in none of these was anxiety prominent.

The records of the patients in the wards for the deteriorated showed that many of them had passed from the acute block to the chronic beds, where later they had developed incontinence or other unpleasant antisocial habits which led to their transfer to this final refuge. Whether the process of decay is inevitable or whether it is caused or accelerated by the apathy surrounding the chronic beds is a matter worthy of consideration.

Medical Treatment

The medical treatment in Western Road Hospital was surprisingly good, although there were only five whole-time medical officers, including the medical superintendent, to care for 1,000 bedridden patients, and they had in addition many other responsibilities—a large venereal department, 150 patients who were up and about but unfit to return to their homes, and 300 inmates, including provision for maternity, under the control of the Social Welfare Committee set up by the National Assistance Act. My registrars made frequent visits, took histories, and made clinical examinations of the standard usual in a teaching hospital, and each week the staff of the infirmary, my registrars, and I spent a long afternoon together in the wards comparing the result of our separate observations. It was rarely necessary to change a diagnosis or possible to modify treatment. A considerable amount of laboratory and radiological investigation had been done in the departments of the Dudley Road Hospital close by.

The individual case records were, it is true, brief and to the uninitiated almost incomprehensible, and they were, unfortunately, unsuitable for exact analysis.

In this connexion I have used the words "medical treatment" in their narrowest sense to signify the therapeutic use of rest and drugs, because in the infirmaries no other form of treatment is generally possible. In Western Road no masseuse had worked for years; there was a room reserved for physiotherapy, but on the door of it was the notice "Do not enter," and within was a heap of electrical apparatus apparently in good order. Rehabilitation could not be attempted, because there was no staff and all the day rooms were full of beds; there was no occupational or diversional therapy, but a few women patients knitted and did needlework. I never saw a man engage in any spontaneous activity in the wards.

Nursing

Of the nursing in Western Road Infirmary I find it difficult to speak with due restraint. When we began our survey the patients were roughly classified thus:

	Males	Females	Total
Group 1: Bedridden	393	550	943
Group 2: Allowed up but unfit for transfer to homes without nursing care ..	67	73	140
	460	623	1,083
The number of incontinent patients among those confined to bed	162	295	457

In addition there was a small maternity ward and a considerable venereal-disease department.

The number of full-time nurses was 181; there were besides 90 half-time nurses, and if they be reckoned as the equivalent of 45 the total available was 226. Only 14 of these were State-registered; the remainder were either assistant nurses or in training.

These figures were given to us before we went into the wards, and I was prepared for an unpleasant experience: it seemed certain that we would find bedsores, dirty beds with the inevitable foul smell, and general untidiness and lack of cleanliness.

The truth was the exact reverse. The wards were clean, the beds were impeccably made (indeed, as we found later when studying consequent deformities of the feet, they were too well made and too firmly tucked in), there was no smell or untidiness. We were quite ruthless and perhaps a little rude in our first inspection; we pulled down bedclothes here and there looking for bedsores and dirty beds and found none. There were only nine bedsores in the whole institution, and these were in cases of gross organic nervous disease in the last stages.

Two nurses were on duty in wards containing 70 patients, of whom 30 were incontinent; they never stopped working—they did not come to speak to the doctor of the ward or to the matron unless they were sent for; quietly and efficiently they went on with their job. In the incontinent cases the backs “were done” every four hours day and night.

The equipment of the wards was lamentable; basins with running water were few and far between, the sluice-rooms were mainly archaic, though here and there a bed-pan washer had been installed; there were no pulleys over the beds; the baths were of the ordinary domestic type—the bedridden were wheeled to them once a week and laboriously lifted in and still more laboriously lifted out. Many of the women patients could not do their own hair, and yet it was tidy and obviously tended regularly by the staff. At night one assistant nurse was in charge of the 70 beds; to turn a helpless patient she had to fetch a nurse from another ward and render her the same service in return. The relations between the staff and the patients were good; in our later clinical examinations we heard frequent spontaneous tributes to the kindness of the nurses, and we were surprised at the extent of their knowledge of the social background and personal predilections of the patients.

Conditions such as these are by no means exceptional. In many institutions in the Birmingham Region they are much worse; in few are they as good. For instance, at Exhall Hospital, in Coventry, there were only two resident nurses for 274 patients of the same sort; at night there were often not enough to allow one to each ward of 50 beds,

and “runners” were employed to call nurses when and where they were most needed. Yet the work was done; we found the wards clean and the patients well tended and without bedsores. In the smaller hospitals classification of the patients was impossible, and we saw babies and young children in beds side by side with senile demented.

In their quiet endurance and their efficiency, as in their triumph over discouraging circumstances and lack of proper equipment, the nurses recalled the virtues of their fathers in the rank and file of the county regiments who held the trenches in Flanders in the campaign of 1914-18 and saved Europe. Certainly by their work they have delivered the hospitals of this country from chaos, for if the infirmaries fail the work of the general hospitals will be speedily stifled by the accumulation of cases of chronic sickness in their wards.

Rivers of ink have already flowed into the sterile sands of controversy about the nursing problem and I do not wish to add to them. I confine myself to the statement that the work of nursing in the hospitals of this country is most inequitably distributed; that a disproportionately heavy burden is borne by the staffs of the institutions for the chronic sick; that, despite neglect and poor equipment, the most heavily burdened have done their work efficiently and humanely, and their triumph owes little to theoretical instruction, academic distinction, or flowers and shampoo-rooms in the nurses' homes.

These simple facts should be considered by those whose duty it is to shape the future of nursing. Kindness, courage, practical resource, and patient endurance of the routine of daily work are qualities more valuable in nurses than a superficial knowledge of the intricacies of anatomy and physiology. It may be that the pursuit of a high average of academic attainment for State-registered nurses is a mistake. It might be better to select a small number for intensive training for positions of responsibility as sisters, matrons, and administrators and to rely more on character and practical work in the wards as the test for registration for the great majority. Unless some change is made it is certain that for many years defects of primary education will exclude a large number of excellent and intelligent nurses from the ranks of the registered. In any event the condition of the sick in the infirmaries demonstrates the foolishness of the contention that the measure of the efficiency of the nursing staff is the average duration of the stay of the patients in hospital.

The recently published Report of the Working Party on the Recruitment and Training of Nurses is not a distinguished document, but paragraph 247 deserves notice for its almost incredible stupidity. It runs: “It is clear that for some time to come, owing to the shortage of trained staff, it will be essential to continue to use the services at least of those assistant nurses now recognized and employed. Such evidence as we have been able to obtain points to the fact that useful service can be rendered in the nursing of mental and tubercular patients and probably also of fever patients and the chronic sick by persons who are not fully trained nurses.” For many years four out of five nurses responsible for the care of the chronic sick have not been fully trained. I have paid my tribute to their skilful and devoted work and hope they will not be distressed by this slighting reference in the report.

Tentative Conclusions Based on Pilot Survey

Our preliminary survey was undertaken to discover whether improvement in the staffing and equipment and the application of the standards of investigation and

RELATIVE AGE DISTRIBUTIONS OF PATIENTS IN WESTERN ROAD INFIRMARY

CHART 1

treatment of a good general hospital would be likely to ameliorate the condition of the chronic patients to such a degree that they would be able to return to their own homes. We were soon convinced that improvement of that extent is quite impossible for the great majority, although a number might be made fit enough for transfer to institutions in which a more active and congenial life is possible than in a hospital for the chronic sick.

The reasons for that discouraging conclusion are three. First, nearly all the patients are the victims of pathological conditions which are either degenerative or genetic in their origin and beyond the scope of therapeutics in its present stage of development. Secondly, long confinement to bed without occupational or diversional therapy, without massage and passive movement in hospitals so grossly understaffed that independent activity of any kind has been discouraged for fear of accident, has led to rapid deterioration in the shape of contractures, deformities of the feet, and the development of incontinence. Thirdly, and probably the most important, is the subtle disintegration of human personality that is so common in these patients. Many literally do not wish to leave the beds in which they have lain so long; several refused admission to my wards in the General Hospital for the reason that if they were cured they might be compelled to leave the infirmary and resume the struggle of life alone. Like T. S. Eliot's women in the Cathedral at Canterbury, they did not wish anything to happen: they wished to be left to perish quietly. We had to comfort others by telling them that we had no intention or desire to discharge them before they would consent to give a history or co-operate in examination. Whether the establishment of homes in which a certain amount of nursing and domestic help is provided would encourage many of these patients to leave the infirmaries willingly is a matter to be decided by trial. I am not very hopeful of the result of such an experiment. I feel that most of the inmates in the chronic wards must be regarded as a sad inheritance from a long period of neglect—like many of

the deteriorated patients in the mental hospitals—and that the opportunity for effective treatment has long since been lost.

At this stage it was clearly necessary to check the accuracy of our impressions by reference to statistical control. My colleague, Professor T. McKeown, of the Department of Social Medicine, with one of his senior lecturers, Dr. C. R. Lowe, came to our help and devised record forms suitable for mechanical sorting and analysis. The students of the University of Birmingham made us a grant of £1,000 to pay the cost of the investigation. At a later date Birmingham Regional Hospital Board, at the suggestion of the Hospital Management Committee at Stoke-on-Trent, sanctioned a similar survey in the Potteries, and soon we hope to arrange another in the rural peace of Herefordshire. The final figures will be published independently from the Department of Social Medicine, but by

the kindness of Dr. Lowe I am able to present the results of a pilot survey, based on a preliminary analysis of the records of 714 patients—318 male and 396 female.

Age, Mobility, and Mental Condition

The mean age of the males is 74 years and of the females 76; 92% of the males and 95% of the females are 60 years and over; 70% of the males and 71% of the females are over 70 years; and 27% of the males and 37% of the females are 80 or over (Chart 1). The largest ten-yearly group for men is from 70 to 79 years and for women from 75 to 84; both these groups are 43% of the respective totals. Clearly the outstanding characteristic of the patients in this infirmary is advanced age.

Of the 714 patients 394 were completely bedridden (49% of the males and 60% of the females); of the remaining 320 only 56 men and 13 women were fit enough to venture out of doors.

At the time of the survey we judged that all but 5% of the bedridden were beyond hope of rehabilitation, but unfortunately it was impossible to determine accurately what their condition had been at the time of admission. We believe that their complete disablement has been in some measure due to continued rest in bed and to a large degree has been aggravated by it, because later we found suggestive evidence that a considerable proportion of the patients had been admitted from their homes simply as a consequence of some domestic upheaval and not on account of deterioration of their condition.

Sixty-three per cent. of the men and 52% of the women were reported to be "occupied" in the sense that they would occasionally read, write, knit, or help in the ward. My own observation, already recorded, suggests that this estimate is excessive. In Affleck's survey at Leeds he included 788 patients in five separate and much smaller hospitals; their age distribution and rough clinical classification corresponded well with ours, but the proportion of bedridden was much lower (in Leeds 33.9% of the males

and 41.7% of the females). Affleck does not give any figures of the nursing staff available, and it may be that the difference between the two groups was due to greater activity in Leeds than was possible in the wards at Birmingham.

Mentally we classified 32% of the males and 39% of the females as abnormal and in so doing ignored senile forgetfulness, irritability, and other minor manifestations of elderly decrepitude; we considered 25% of all the patients certifiably insane. These figures are in exact agreement with Affleck's estimate of 37.3%, and give further support to the explanation suggested by Lewis (1946) of the difference between the first admissions to English and American mental hospitals of patients over 65 years of age. The difference is not in the incidence of psychosis but in the method of dealing with it. In England a large proportion of these cases are cared for in the hospitals for the chronic sick, and in America they are in the mental hospitals. In England many patients thus escape the statistical analysis of the Board of Control.

Affleck believes that the vast majority of the mental disorders among the chronic sick are organic in basis, although he admits the possibility of the importance of functional elements derived from hopelessness and lack of interest. The reason for his belief is the undoubted fact that in nearly all these cases it is easy to elicit evidence of somatic disease such as arteriosclerosis, cerebral thrombosis, and cardiac insufficiency. On the other hand, in ordinary clinical practice cases of that type are common with no evidence whatever of mental failure.

I confess that I am left with the uneasy suspicion that much of the psychiatric disorder in the institutions for the chronic sick may be the result of their apathy and routine; no dogmatic statement, and indeed no opinion, is justified at the present time. What is needed is a systematic study of the mental and physical condition of the patients at the time of their admission to the infirmaries and of their subsequent progress. Such study has hitherto been impossible because the institutions for the aged and chronic sick have been neglected by both physicians and psychiatrists and because their staffs have been too small and too overburdened with routine work to undertake it.

Marital Status

Of the men 58% are widowers, and of the women no less than 73% are widows. Both these figures are significantly higher than those for similar age groups in the general population of Birmingham. The proportion of the unmarried in the infirmary is also higher than outside. The result is conveniently illustrated in Chart 2, which shows that, if a man or woman is married and the mate survives, the risk of ending his or her days in an infirmary is considerably reduced—a conclusion that may bring comfort to those who worry about the social institution of marriage. In Leeds only 34.5% of the men were widowers and 59.3% of the women widows.

Duration of Stay

The mean duration of stay for males was 34 months and for females 37 months (the range in both sexes from 0 to 18½ years); 26% of the men and 28% of the women had been in the infirmary for more than three years; 71% of the women and 62% of the men had been there for a year or more. These figures do not represent the work of the infirmary accurately; they leave an impression of a slow drift to inevitable death over a period of many months, and this is far from the truth. During 1948, for instance,

MARITAL STATUS (Patients 65 years and over) compared with 1939 National Registration, Birmingham.

CHART 2

there were 958 deaths in the institution; of these 656 (68%) occurred within 100 days of admission and no fewer than 548 (57%) within 50 days; only 12% of those that died had been in Western Road for two years or more. Clearly an important part of the work was the treatment of acute illness in old people, and we decided to investigate this aspect of the problem more closely by reviewing the subsequent history of every patient admitted during one year. Mr. Broad, the clerk in charge of the records at the infirmary, who had already supplied us with the figures relating to the deaths in 1948, undertook the formidable task of preparing a summary of the case histories of 2,478 patients admitted between Jan. 1 and Dec. 31, 1946 (Tables I and II and Chart 3), and Dr. Padley has provided me with the following analysis:

TABLE I.—Admissions, Western Road Infirmary, 1946

Age	Males	Females	Total
Under 50	260 (17.7%)	124 (12.3%)	384 (15.5%)
50-59	199 (13.5%)	75 (7.4%)	274 (11.1%)
60-69	357 (24.3%)	218 (21.6%)	575 (23.2%)
70-79	493 (33.5%)	384 (38.1%)	877 (35.4%)
80+	161 (11%)	207 (20.5%)	368 (14.9%)
Total	1,470	1,008	2,478

TABLE II.—Fate of Patients Admitted in 1946. Period of Follow-up 26 Months

	Males	Females	Total
Died:			
In less than 4 months	437 (29.7%)	355 (35.2%)	792 (32%)
In 4 to 26 months ..	115 (7.8%)	82 (8.1%)	197 (7.9%)
Discharged:			
In less than 4 months	784 (53.3%)	385 (38.1%)	1,169 (47.1%)
In 4 to 26 months ..	93 (6.3%)	103 (10.2%)	196 (7.9%)
Survived	41 (2.8%)	83 (8.2%)	124 (5%)
Total	1,470	1,008	2,478

Months are lunar months of 28 days.

The fallacies attending conclusions drawn from a survey of patients within a short period of time are at once apparent, because large numbers of chronic patients will

accumulate over years and distort the picture of the work done. More detailed analysis yielded some interesting results: (1) Men are admitted to the infirmary at an earlier age than women; only 44% of male admissions are 70 years or over, whereas 58% of the females are of that age group. (2) No less than 32% of all admissions died within four lunar months and in the same period 47.1% had been discharged. (3) In 26 months a further 15.8% of the patients either died or were discharged (7.9% in each group). (5) Only 5% of the total admissions survived 26 months or more, and in this group the proportional survival of women was three times greater than that of the men (8.2% as against 2.8%). To adopt a simile suggested by Lancelot Hogben—the patients entering an infirmary are like a crowd of old people passing slowly through a gate into a long avenue. Within four months more than three-quarters have left the avenue either by death or by return to their homes. In two years 19 out of 20 have gone and only a few remain passing through the shade: these stragglers may long survive, and their accumulated numbers at a given moment of time give a false impression of the true condition of life within the grove.

A hospital for the chronic sick is a bad description of Western Road Infirmary, and I suspect for many others. Most of the patients admitted are acutely ill, and their treatment is undertaken in an institution without a laboratory or radiological department and with a staff so small that it is remarkable that the treatment and the nursing were as good as we found them to be. It is true that the clinical and laboratory staff at Dudley Road Hospital, close by, had given all the help that lay in their power, but it would not be candid to pretend that the arrangements were satisfactory. The taking of a radiograph involved a journey of half a mile, and many of the patients were unfit for such adventure; in any event the radiological department was too small for the needs of the acute hospital and desperately overworked. The difficulties of laboratory investigation by messenger and transport of specimens need no elaboration.

I have compared Dr. Nagley's results at Western Road with Mr. Cosin's (1948) at Orsett Lodge Hospital, in Essex. At Western Road the number of admissions was greater than at Orsett and the proportion of older patients was also larger (see Table III). Cosin does not deal with the fate of the patients under 60, and a comparison of those over that age gave the result shown in Table IV. It seems likely that the greater proportion of deaths at Western Road was due to the greater age of the patients. What is surprising is that the ratio of discharges to admis-

TABLE III

	Western Road, Birmingham	Orsett Lodge, Essex
Total admissions in one year	2,478	1,430
Aged 60 years and over ..	1,820 (74%)	780 (55%)
Under 60 years ..	658 (26%)	650 (45%)

TABLE IV

	Western Road, Birmingham	Orsett Lodge, Essex
Total admissions ..	1,820	780
Died ..	902 (49%)	296 (38%)
Discharged ..	807 (44%)	284 (37%)

sions in this age group was also larger in Birmingham, because at Orsett there was an active department of physiotherapy and there was none at Western Road. The inference is that the discharge of an elderly patient from an infirmary depends more on the condition of his home than it does on the attention of a masseuse.

WESTERN ROAD INFIRMARY: ADMISSIONS 1946

RESULTS OF 2-YEAR FOLLOW-UP

□ Died ▨ Discharged ■ Remaining

CHART 3

Prelude
to Admission

No less than 77% of all the patients were admitted direct from their own homes; not more than 18% had passed through a general hospital, and of this number one-sixth had passed through the general hospital in a technical sense only—that is, they had been sent on to the infirmary from casualty and out-

patient departments and had not been in the wards. In brief, six out of seven patients had been admitted to this understaffed and ill-equipped institution without adequate investigation or treatment. To the credit of the general practitioners of Birmingham it should be recorded that in more than one-half of the patients admitted to the infirmary under the age of 60 years some form of laboratory or other investigation had first been undertaken in the difficult circumstances of their own homes. The figures are in striking contrast with those from Leeds, where it is stated nearly all had been thoroughly examined in hospital.

A study was made of the medical and nursing attention required by individual patients at the date of admission, and it was found that in 33% all that was needed was a visit from a doctor once a month; this group had no skilled nursing of any kind at home, and it was clear that the only reason for their entrance to the infirmary was some sort of domestic catastrophe such as the death of a spouse or friend or illness of the person who had previously cared for them. These cases are entirely a social problem; for the most part they needed only help in their houses or untrained attention to their physical disability. The estimate of 33% is a conservative one; when there was any doubt about a particular case it was excluded from the group. It was clear that many of these patients had

deteriorated in the infirmary—a consequence, we believe, of the prevailing apathy and of prolonged rest in bed. That so large a proportion as one-third of all the patients had been admitted without disability sufficient to require admission on medical grounds lends support to the suggestion of Brooke (1949) that social services organized by the hospital to care for the elderly in their homes might save many of them from the last dreary journey through an infirmary to a squalid tomb.

Investigations at Highbury, a home for 270 old people in Birmingham under the administration of the local authority, warranted the same conclusion. In this institution the mean age of the patients was 78, higher than it was in the infirmary, and they were living a much more active and pleasant life at a weekly cost of less than half that of an infirmary bed. Had this home not existed we were satisfied that a very large proportion of its inmates would have required accommodation in an infirmary. There is some dispute about the proper size of such homes, and the National Corporation for the Care of Old People favours small ones for 25 to 30 residents. The master and matron of Highbury were convinced, on the other hand, that in large cities homes for old people should ideally have about 400 beds, and that most of these should be in small separate rooms, some with two beds for married couples.

When I asked whether the provision of separate rooms would not make the cost of domestic service prohibitive they replied that, on the contrary, it would solve the problem, because nearly all the old people would keep their own room spotlessly clean without help, but many of them resented cleaning part of a dormitory. Their main arguments in support of their preference for the large home were that the old people of the cities liked the atmosphere and bustle of a crowd, that communal entertainment in the shape of films, discussions, and stage shows was much more easily and better organized than in the small homes, and that the feeding would be better and more economically done.

Attitude to Leaving the Infirmary

One-third of all the patients were unwilling to leave the infirmary in any circumstances; as I have already said, the mere suggestion that they might be made well enough to do so distressed them seriously, and we found that this frame of mind was directly related to the duration of their stay in the institution. Thus, within the first three months of their admission 65% of the men and 71% of the women are eager to return home, but after four years the figures fall to 45% of the men and 28% of the women, and I am sure that the fall would be even greater if the patients were actually confronted with the opportunity of leaving. In part this reluctance to leave is due to the disintegration of their homes, for we found that more than half (57%) of all patients had no home of any kind to which they could go if they were discharged. Of the 270 patients with a home available to them less than a half (44%) could expect any form of help in the house, and more than one-half (56%) would either be compelled to live entirely alone or in a house deserted during working hours.

It is significant that during the first three months of their stay 71% of the males and 51% of the females have a home to which they would be welcome on their return; thereafter the home and family background progressively recedes and after four years only 24% of the males and 13% of the females have any sort of home available to them. Manifestly the cohesion of the homes of the elderly is fragile: absence for more than three months is sufficient to damage it; in four years more than three out

of four of their homes have vanished. Two problems emerge clearly from this analysis: the first is concerned with the maintenance of the elderly home, and the second is that of provision of adequate treatment and social help for the old when they fall sick.

Factors Affecting the Cohesion of the Home

Psychologists and social anthropologists agree that the forces which promote cohesion in a group are derived from fundamental instincts modified by habit and custom and, to a very slight degree, by rational considerations. The instincts are of two kinds—the self-regarding and the group-maintaining: the first are the instincts of self-protection, sustenance, and assertion, and the second those of mating, nursing, and the herd. At the most primitive level these instincts were satisfied by the organization of society in family groups, but the passion and pugnacity inseparable from mating led to the institution of exogamy to avoid the internecine strife that ensued from sexual selection within the family. Exogamy established the tribe as the social unit; conquest and enslavement of the conquered made the nation. The discipline of law with the habits and customs that it implies maintains the cohesion of a people so long as the makers of the law are wise enough to see that its system does not conflict unduly with the fundamental instincts that lead to the development of society or with its interests.

It is the opinion of many with long experience of the work in institutions for the chronic sick and aged that the sense of family unity and responsibility is weakening, and that people in this country are less interested than they were in the fate of their elderly and disabled relatives. We are also told that parents to-day are not as willing to sacrifice personal comfort and pleasure in the interests of their children, and that many children at the age of puberty or soon after lose all sense of responsibility and affection for their family.

Accurate estimation of the value of evidence of this kind is difficult, because it emanates from those whose work constantly brings them face to face with the tragedy of the broken home and their judgments may be unduly melancholy. Sheldon's (1948) report on the old people of Wolverhampton shows how ardently the flame of family affection may burn despite discouraging circumstances, and suggests that the evidence should be accepted with caution. It would be unwise, however, to neglect it, for it comes from many sources and its volume is increasing.

The loosening of the ties of English family life and the rapid increase of divorce attracted serious attention in 1918 at the end of the first world war and have become more pronounced during the past 30 years, especially in the period of the second campaign.

Disruption of a family by conscription and long terms of separation inevitably diminished cohesion; so did unemployment and the application of the means tests in the uneasy period of truce. During the industrial depression many a father found his dignity and authority as the main support of a family replaced by a sense of ignominious dependence on the earnings of children, who sometimes resented the burden of the maintenance of their parents' home.

The deplorable condition of houses and their overcrowding made the position of sick and disabled members of a family increasingly difficult, and the burden of their care in restricted accommodation seemed intolerable. Many social workers regard this factor as the predominant one, and it is often pressed in argument by those who seek a bed in an institution for a dependent relative. It cannot,

however, be alone responsible for the increased demands on the infirmaries, because, first, the overcrowding of houses is less than it was 50 years ago, when the average size of families was larger, and, secondly, applications for institutional provision often come from those who cannot reasonably claim that their housing accommodation is unsatisfactory. It is difficult, too, to induce families to take back their relatives when new houses are available to them. Clearly there is now less reluctance to seek the aid of public assistance in solving the problems of family life. The administration of the infirmaries has become more sympathetic and humane, the horror of the Victorian union is no longer so deterrent, but it is also possible that slackening of the moral fibre of the family and a demand for material comfort and amenity outweigh the claims of natural affection.

A shorter working week for men has encouraged the employment of women in industry; homes therefore have less attention at a time when they are difficult to run. For the same reason domestic service is almost unobtainable and local authorities are unable to recruit "home helps" to care for the sick and aged in their houses.

It may well be, therefore, that the weakening of family cohesion which so many observers claim to have detected is in part a result of the trend of recent legislation. As a contribution to the study of this social problem we investigated the frequency of visiting at the infirmary, with the following result. Eighty-two per cent. of the men and 79% of the women are visited at least once a week for the first three months of their stay; thereafter the frequency of visits declines steadily, but it is not until four years have elapsed that the total of weekly visits falls to 40% for the men and 46% for the women.

Irrespective of the duration of stay of the whole sample of 714 patients, no less than 52% of the males and 68% of the females are visited by friends or relatives once a week; only 48 men (15%) and 28 women (7%) are never visited. The remainder were visited at intervals greater than a week.

In view of the depressing environment of the infirmary, of the unpleasant nature of many of the disabilities in the patients, and of the difficulties of travel, these figures show that even if natural affection has declined, as some believe, it is still a potent influence in the social life and conduct of the English. Curran *et al.* (1946) reached the same conclusion in Glasgow, and it must never be forgotten that at least 95% of the aged never go into an institution: with the help of their families and friends, or more rarely alone, they struggle on to the end.

Indisputably necessary as it was to do something to improve the material standard of life of the people, it must always be remembered that it is possible to do too much. Freedom from tyranny, from poverty and want, from the fear of unemployment, and from the hopeless despair of unattended and persistent sickness is no doubt desirable, but whether it is good to achieve it without effort on the part of the individual or his family—to have it, as it were, "laid on" by the State—is questionable, and the matter is more commonly decided by an appeal to sentiment or self-interest than by reason.

The instincts of self-protection, sustenance, and assertion imply struggle to satisfy them, and if society makes the struggle unnecessary the vitality and survival value of its individual members will diminish. Similarly, the power of the group-maintaining instincts will suffer if the provision of a home, the training of children, and the care of its disabled members are no longer the ambition of a family but the duty of a local or central authority.

In these debatable matters a decision requiring high qualities of statesmanship is necessary before new plans are made to deal with the problems of the chronic sick and infirm. The problem is simple. Shall they become the responsibility of society in large institutions or should we encourage their friends and their families to care for them whenever possible?

In the interest of social cohesion psychology and anthropology favour the second course, and so does ethics, and, while it has been the constant aim of religion and morals to extend the affections of the individual beyond the range of his personal and group relations, both agree that such extension must be based on a lively sense of responsibility for the weaker members of the family and on service to them. Aesthetics in days of realism is often confused with sentiment, but there is a beauty and a value in personal service to a crippled friend or relative that no hospital can rival. Claims of expediency and economy may be vulgar, but they are insistent, and the fact that care of the sick and disabled in their homes is cheaper than in institutions is worth emphasis.

For all these reasons, and to avert the distressing disintegration of human personality that is so common in patients who lie for years in an infirmary, it would be wiser to endeavour to preserve the sense of family responsibility and to encourage people to tend their own sick and guard the aged.

That does not mean that families should be expected to bear the whole burden. Society must help them, and the problem really hinges on the type and extent of assistance to be provided. Institutions will always be necessary, but the aim should be to reduce the demand for them, and the measures to be adopted must not conflict with the claims of human instinct or with the principles of ethics, morals, and religion.

(A list of references will appear at the end of the second lecture, which will be published in our next issue.)

A memorandum prepared by the Minister of Health's Standing Advisory Committee on Cancer and Radiotherapy and endorsed by the Central Health Services Council has just been issued for the general guidance of regional boards and boards of governors. The memorandum suggests that no recognition should be accorded to "cancer" as a specialty in itself. To secure full efficiency a cancer service should provide a single organization to serve a population of the order of two million to four million. The area served should normally, but need not necessarily, be coterminous with that of a regional hospital board. Each cancer service should have a co-ordinating committee, appointed jointly by the regional hospital board and board of governors, concerned with the allocation of work to hospitals, records, follow-up, transfer of patients, provision of equipment, and other matters. It should include representatives of the university and medical school, and should have the advice of local health authorities and of a professional and scientific subcommittee including representatives of the universities and the appropriate medical school, medical officers of health, and general practitioners. Facilities for preliminary consultation should be established at an adequate number of recognized peripheral hospitals, which should be visited regularly by specialists for consultation with the general practitioner and local staff to assist in preliminary diagnosis and for follow-up. Similar facilities should exist in all units of the organization. The co-ordinating committee should advise on the selection from among the physicians, surgeons, radiotherapists, and other specialists on the staff of hospitals in the area of those who are to form the headquarters staff of the cancer service. Treatment by radiotherapy should be under the control of specialists engaged solely in that work.