Chart showing rise in latent period at successive freezings.

As stated in the preceding paper, sex is immaterial provided

added. It was sealed in ampoules containing 0.5 ml. each. These were frozen in a solution of CO₂ snow in "cellosolve" (ethylene glycol monoethyl ether, B.D.H.) to -79° C. for varying periods (from 15 minutes to 182 days). Each ampoule was then thawed, sometimes by immersion in water at 2° C., sometimes by holding in the hand. (No difference was noticed between the two methods.) If it was to be tested it was opened, the contents drawn up into a syringe, and 0.05–0.1 ml. (as in Cramer's original experiment) injected into the right

the injection is made in the nipple line. If the mince was to be refrozen it was dropped (after an interval of approximately five to six minutes) again into the cellosolve. Freezings other than the first were for 15 minutes to one hour. Tables A and B show the variations in the experiments and the results.

It is interesting to note that the appearance of those tumours which show infectivity after two or even three thawings is delayed with each successive thawing. Table B also demonstrates the slight heightening of activity when the same tumour is frozen for some days instead of for 15 minutes. The Chart demonstrates graphically the rise in the latent period for each tumour at successive freezings. It is therefore obvious that some activity is lost each time the tissue is thawed.

Conclusions

Repeated freezing and thawing of mouse mammary carcinoma tissue reduces its infective activity.

The apparent difference between the behaviour of sarcoma and carcinoma on repeated freezing and thawing is a quantitative and not a qualitative one.

PROPAGATION OF MOUSE CARCINOMA BY DRIED TUMOUR TISSUE

BY

IDA MANN, D.Sc., F.R.C.S.

AND

WILLIAM J. DUNN

(From the Laboratories of the Imperial Cancer Research Fund, London, N.W.7)

The propagation of mouse sarcoma, both chemically induced and sporadic, by means of tumour tissue frozen to -79° C., thawed, and then dried *in vacuo* at -15° C. has been demonstrated by Professor Gye (1949). That investigation revealed the presence of an agent, other than a living sarcoma cell, which has the properties of a virus. Since the spontaneous mammary cancers of the Strong A and R III strains of mice are known to be caused by the Bittner virus, it appeared to us desirable to try to propagate mammary cancer by the same method. The difficulties are, however, theoretically greater in the case of carcinoma than in sarcoma.

In the first place, although we know that the virus is liberated in its active form by prolonged freezing of the minced tumour, we also know that it is tissue-specific and must therefore be brought into intimate contact with mammary tissue. We know that in its active form it can infect male mammary tissue as easily as female if injected in the nipple line. But this tissue specificity in itself leads to a

lower percentage of tumours in carcinoma than in sarcoma, since in the case of the sarcoma mutant of the virus any and every subcutaneous connective-tissue cell is susceptible to infection, and connective tissues are, moreover, everywhere easily accessible to an active sarcoma virus.

In the second place, experiments with serial freezing and thawing have shown that the Bittner virus is extremely thermolabile, and that many tumours will not withstand a second freezing and thawing. Since the method of drying sarcoma tissue described by Gye, Begg, Mann, and Craigie (1949) involves freezing the minced tumour in an ampoule, thawing it, transferring it to the drying-flask, and refreezing it at the beginning of the drying process, it is likely that only a few carcinomas (such as M103 and M107, described in the preceding paper at page 253) will yield active dried material by this method. We know certainly that such tumours as "63" and M91 (in the previous paper) will not withstand drying by this method. We also ascertained this

FIG. 1.—Section through the R III sporadic mammary tumour used in Experiment 111 (see Table).

FIG. 2.—Section through the mince of the tumour shown in Fig. 1, after drying. No cellular structure is seen.

experimentally by drying "63," two C3H, and two Strong A sporadic tumours by this method. These gave uniformly negative results.

A method was therefore required which would obviate the need for thawing the material before transferring to the drying-flask. This has been done in two ways; first, by grinding up the frozen material to powder, in a mortar embedded in CO₂ snow, dropping the powder into the drying-flask, also cooled below freezing-point, and sticking it to the bottom of the flask with a small amount of ice-cold 5% glucose solution, which immediately solidifies. This is necessary, since the material to be dried must be in close contact with the bottom of the drying-flask. The second and easier way is to freeze the tumour mince in the flask in which it is to be dried. For this method a tubular container was used in place of the Erlenmeyer type of flask usually employed. This made storage at -79° easier, and all that was required was to take the tube from the "deep freeze" and attach it at once to the drying apparatus. This method has the slight disadvantage that material cannot be removed from the flask for test for activity after freezing but before drying. This is easy when the first method is used.

Either method, however, gives successful results. We have succeeded in six consecutive experiments, using six different sporadic mammary cancers, in obtaining tumours from dried material in both male and female mice. The tumours appear in the nipple line at the site of injection in from four to 10 weeks. All the mice were young and well under the cancer age. Photomicrographs of sections

of R III sporadic mammary tumour used and of the tumour produced by inoculation of the dried material are reproduced in Figs. 1, 2, and 3. The Table shows the results.

Technical Details

1. *Preparing the Tumours for Freezing.*—The mouse is killed with coal gas. The tumour is dissected out aseptically, a small piece fixed in Bouin's solution, the rest minced in a sterile Craigie mincer and diluted with an equal volume of 5.3% dextrose solution. If it is to be stored in ampoules, 0.5–1 ml. is placed in each by means of a syringe with a wide-bore needle. The ampoules are then frozen rapidly to -79° C. and stored in solid CO₂ for the desired time. If storage in the drying-tube is desired 1 ml. of the mince is run into the tube and by manipulation over crushed CO₂ "dry ice" is made to freeze in a thin layer spread over the sides and bottom of the tube. The tube is then stored as above.

2. *Drying.*—The frozen ampoule is removed from the storage cabinet, cracked open, and the solid mass extracted into a sterile mortar embedded in CO₂ snow. All the instruments used are sterile and chilled by keeping in test-tubes embedded in CO₂ snow. The solid mass is ground in the mortar and most of it tipped into the cold drying-flask, which stands in a vacuum jar of "cellosolve" at -15° C. Ice-cold 5% glucose solution is dropped with a pipette on to the powdered mass in the flask. This solidifies and sticks it to the bottom of the flask. The drying is then started (see Craigie (1949) for details of the apparatus). The residual moiety of the frozen material left in the mortar is now

FIG. 3.—Section through the tumour produced by inoculation of the dried material into the breast tissue of a 6-weeks-old R III mouse.

Production of Mammary Cancer with Tissue Frozen and Dried without Thawing

No. of Expt.	Tumour	Time Frozen Before Drying	No. Inoculated	Tumours Obtained	Sex	Time of Appearance	Control: Frozen Only	
							Inoculated	Tumours
82	Mixture of 3 sporadic strong A tumours	3 weeks	6	3	1 F., 2 M.	5 weeks	3	1
82	"	4 "	6	3	2 F., 1 M.	4 "	3	2
97	"	56 days	6	2	2 F.	7 "	3	3
99	R III tumour from tumour which had been frozen 138 days and then inoculated	131 "	6	2	1 F., 1 M.	10 "	3	3
106	Strong A tumour from tumour frozen 14 days	24 "	6	2	2 M.	8 "	3	3
111	R III sporadic	5 "	6	1	1 F.	5 "	Not done	

thawed and inoculated into three mice of the appropriate strain as a control for the activity of the frozen tissue.

3. *Inoculation of Dried Material.*—When the drying is complete a small portion of the dry material is dropped into xylol and embedded direct for microscopical examination (no dehydration is of course necessary). The rest is wetted by running into the flask 2–3 ml. of a 1 in 500–1,000 neutral solution of cysteine. This produces an even emulsion, 0.05 to 0.1 ml. of which is then injected in the nipple line of each of six mice (three male and three female) of the appropriate strain. Attempts were made to improve the results by adding 33% rabbit serum saline to the cysteine and by the use of embryo extract. The latter is definitely harmful and the former of no greater value than cysteine alone, which is therefore recommended.

It must be repeated that avoidance of thawing at any time before drying is complete is essential. Experiment 82 was repeated without taking this precaution and using material known to be active, and no tumours were obtained from the dried material.

Conclusions

The active form of the Bittner virus obtained from mouse mammary cancer by freezing can be successfully dried *in vacuo* provided no thawing occurs before the material is dry.

Active dried mouse mammary cancer produces tumours in both male and female mice when injected into breast tissue.

REFERENCES

- Craigie, J. (1949). *Brit. J. Cancer*, 3, in press.
 Gye, W. E. (1949). *British Medical Journal*, 1, 511.
 ———, Begg, A. M., Mann, I., and Craigie, J. (1949). *Brit. J. Cancer*, 3, in press.

The annual report of the National Council of Social Service (1947–8) clearly shows how independent organizations can still make a helpful contribution to public welfare by working in collaboration with the developing statutory services and, even more important, by undertaking research work in new fields. The British delegation played a notable part at the International Conference of Social Work at Atlantic City in 1948. In the same year the National Council organized with great success an International Youth Conference in London, and delegates from 25 countries took part. The activities of the Council's associated groups have also continued to expand. The National Old People's Welfare Committee, for instance, has established 17 county committees and 235 local committees. The National Assistance Act has increased the need for co-operation between voluntary agencies and Government departments. In the words of the report, "The problem is so large and increases so rapidly that it requires the combined efforts of both statutory and voluntary agencies to meet the needs. . . . The stage is set therefore for a concerted effort over the whole country for the welfare of the older members of the community." A revised and enlarged edition of the handbook *Old People's Welfare* is to be published this year.

URETHRITIS, GONOCOCCAL AND NON-SPECIFIC, TREATED BY AUREOMYCIN

BY

R. R. WILLCOX, M.B., B.S.

Physician in charge of Venereal Diseases Clinic,
 King Edward VII Hospital, Windsor; Senior
 Assistant Medical Officer, Department of
 Venereal Diseases, St. Mary's Hospital

AND

G. M. FINDLAY, C.B.E., M.D., Sc.D., F.R.C.P.

The treatment of non-specific urethritis has up to the present been unsatisfactory. Sulphonamides combined with urethral irrigation have in some cases been effective, but penicillin has proved of little value. Streptomycin has been shown to be successful in certain cases (Pulaski, 1947; Kane and Foley, 1947; Willcox, 1949); it is also known to be active against pleuropneumonia-like organisms both in animals and in man—a point of some interest, since there is evidence to suggest that this group of organisms may sometimes be associated with the distressing and by no means uncommon complaint of non-specific urethritis.

In the present communication the effects of "aureomycin" in non-specific urethritis are described. Pleuropneumonia-like organisms were obtained from the urethra in certain of these cases.

Aureomycin hydrochloride, dispensed in the form of capsules each containing 250 mg., can readily be given by mouth (Duggar, 1948), and oral administration is in fact probably more satisfactory than injection. Aureomycin is known to be effective against various rickettsial infections, brucellosis (Spink *et al.*, 1948; Galpine, 1949), staphylococcal and pneumococcal infections, and atypical pneumonia. Certain organisms infecting the urinary tract, such as *Aerogenes aerobacter* and *Streptococcus faecalis*, are susceptible, but *Proteus vulgaris* and *Pseudomonas pyocyanea* are resistant. Collins *et al.* (1948) and Finland *et al.* (1948) found that gonorrhoea was influenced: of 66 cases given an average of 1.3 g. over a period of 1.2 days 49 were cured, 11 were considered doubtful successes, and six were failures. Two cases of non-specific urethritis, both of which had resisted repeated courses of penicillin, sulphonamides, and streptomycin, were also successfully treated: in one, organisms of the pleuropneumonia-like group were present in the urethra before treatment. This patient was given 9 g. of aureomycin by mouth in six days: he was symptomatically improved within three days, and cultures were negative. The other patient, of whom no bacteriological details are given, received 11 g. in five days. He rapidly became asymptomatic.