BRITISH MEDICAL JOURNAL

LONDON SATURDAY JULY 2 1949

British Medical Association

ONE HUNDRED AND SEVENTEENTH ANNUAL MEETING, HARROGATE, 1949

President's Address

MAN AND THE MACHINE

BY

C. W. CURTIS BAIN, M.C., D.M., F.R.C.P.

Senior Physician, Harrogate General Hospital

During the course of his address last year Sir Lionel Whitby dealt with the changes in the practice of medicine which might follow the introduction of the National Health Service. I propose to consider a trend which has developed in recent years, and which might in some circumstances be accelerated by the new Service. I shall illustrate this trend by a study of the progress of events in cardiology, because that is the branch of medicine with which I am most familiar.

During the many centuries that preceded the discovery of the circulation by Harvey in 1628 the views of the profession upon this subject can only be described as queer. Following the teaching of Galen, they believed that the blood was manufactured in the liver from the food; thence it circulated along the veins with an ebb-and-flow motion to all parts of the body. Some passed into the chest and reached a blind diverticulum—the right ventricle—and thence seeped to the lungs where it was rid of some of its impurities. The air, or *pneuma*, taken into the lungs was conveyed thence by the pulmonary veins to the left ventricle. Here it met a little blood which had trickled through invisible pores in the interventricular septum, and the highly purified mixture of air and blood was distributed along the arteries to the different organs. There was no connexion between the two circulations, nor did the heart act as a pump. Hence, although the pulse was examined from the early days of ancient Egypt, it is difficult for us to gather what was learnt from it, and the beginnings of the scientific approach to circulatory problems should be attributed to Sir John Floyer, who wrote in 1710. He had a watch constructed which ran for a minute only, and with its aid made accurate observations on the rate of the pulse in regard to age, sex, emotion, diet, and climatic conditions. He also noted alterations in frequency due to different diseases and in response to various drugs. The slowest pulse he recorded was 55, the fastest 132. It is interesting to note in passing that Floyer also wrote a book on the merits of bathing, which reintroduced the practice of hydrotherapy into this country and gave an impetus to the development of spas.

During the course of the eighteenth century Vieussens described the water-hammer pulse of aortic incompetence, as well as the back-pressure phenomena of mitral stenosis; all the valvular lesions were recorded by pathologists such as Morgagni; Heberden produced his classical description

of the symptoms of angina pectoris; and Withering in 1785 wrote his *Account of the Foxglove*. But the next great advance in technique did not come until 1819, when Laennec introduced the stethoscope. This caused attention to be focused upon cardiac murmurs, so that, although notable advances were made in other directions, including James Hope's conception of cardiac failure, and the recognition of Adams-Stokes disease, the nineteenth century is chiefly distinguished by its concentration upon the differential diagnosis of the valvular lesions. Curiously, although disease of the coronary arteries was recognized, what we now know to be cardiac infarcts were ascribed to chronic myocarditis.

Developments in Cardiology

At the turn of the century four developments took place which revolutionized the practice of cardiology. The first was the impact of Mackenzie. Although his instrument, the ink polygraph, is defunct, his work, which was largely carried out while in general practice at Burnley, fixed the attention of all upon the paramount importance of the heart muscle, and away from the obsession of the previous era with the murmurs of valvular deficiencies. This fundamental conception has formed the basis of nearly all the advances of this century. The second event was the discovery of the x rays by Roentgen in 1895, and the application of the new method to cardiac problems through orthodiagrams by Moritz in 1902. The third, and perhaps outstanding cardiological achievement, was the construction of the electrocardiograph by Einthoven in 1903. As so often happens in medicine, this was the culmination of the work of several men, of whom Burdon-Sanderson, of Oxford, with his capillary electrometer, and Waller should be mentioned.

The fourth advance is one upon which I might be permitted to enlarge because of its local interest. In 1896 Riva Rocci designed a mercurial sphygmomanometer which did not differ much from those in use to-day. About that time George Oliver was devoting his summers to his practice in Harrogate and his winters to research. Oliver had an inventive turn of mind, and in 1898 described a haemodynamometer to the Physiological Society, in which the blood pressure was recorded by means of a calibrated spring. Three years later he published records of blood-pressure readings showing the effects of exercise, of temperature, of age and weight on the same lines as Floyer

had done with the pulse two hundred years before. Sir Clifford Allbutt was then at Leeds and was interested in the problems of hyperpyesia, and so it came about that, after the introduction of the sphygmomanometer, blood-pressure readings were being taken regularly in the West Riding, and especially perhaps in Harrogate, before the rest of the country had become accustomed to them. Consequently the method was regarded with due suspicion for some time. Oliver was also an early believer in the bedside experimental method, and so came to inject his son with an extract of suprarenal tissue obtained from a local butcher. This sent up the boy's blood pressure and was directly responsible for the isolation of adrenaline. The name of Oliver is almost forgotten now, but I still treasure a letter from Wenckebach, written shortly before his death, in which he asks if I remembered his friend Oliver, for "he was a splendid man."

The Electrocardiograph

The electrocardiograph was first used to distinguish and classify the cardiac arrhythmias, most of which concern the auricles, and with this the name of Lewis will always be associated. Lewis bought an electrocardiograph for himself and was allowed to install it in a cellar at University College Hospital. With this instrument he was able to recognize auricular fibrillation in 1909, and in 1920 showed that the cause of the arrhythmia was a circus movement in the auricles. In his book, *The Mechanism and Registration of the Heart Beat*, he described many other forms of cardiac arrhythmia. But Lewis did much more than this. He founded a school of cardiology which attracted students from other lands, who, returning home, developed new lines of research in many directions. Thus from the period after the first world war the tempo of advance has been much quickened, and it will be easier henceforward to deal with each technique separately than to take them in chronological order. At this time, too, a group of physicians who were examining pensioners with heart disease decided to meet annually in order to discuss subjects of common interest. Thus was born the Cardiac Club, which in 1936 was expanded to the present Cardiac Society. It must be admitted with regret, however, that since the era of Lewis no great advance in cardiological technique has been initiated in this country.

A more extended use of the electrocardiograph was foreshadowed in 1920, when Pardee described specific changes in the curve following cardiac infarction. The addition of an apical chest lead in 1932 added considerably to the number of anterior infarcts which could be diagnosed. In 1934 Wilson began his work on the "V" unipolar precordial and limb leads, which reduce distortion from the remote electrode to a minimum. With this technique six chest leads are taken on a line extending from the right of the sternum to the mid-axilla. In selected cases another series may be taken higher up the chest in the third intercostal space. Sometimes oesophageal leads, which come into close contact with the back of the heart, may be added. By this method of electrical exploration of the heart few cases of infarction escape detection, and the site of the infarct can be determined with fair accuracy. Furthermore, from the presence or absence of Q waves and of R-ST displacements it is usually possible to say if the whole thickness of the ventricular wall has been affected or if the infarct is only subendocardial, while the number of precordial leads involved provides a clue to the size of the infarct. Ventricular hypertrophy causes changes in the chest leads; alterations in the position of the heart can be distinguished by the unipolar limb leads. But perhaps

the main advantage of Wilson's technique is that it has placed the electrocardiographic abnormalities upon a firm basis of electrophysical fact. No longer is it necessary to learn the different types of curve by rote; the reason for the altered deflections can now be understood.

The Vector Cardiogram.—An adaptation of the electrocardiogram, of which more may be heard in the future, is the vector cardiogram, which was described by Schellong in 1937. A vector is a term denoting electromotive force, and it can be represented as a line, the length and direction of which show the magnitude and direction of the force. In the ventricles the vector at any given moment represents the electrical axis, and the vector cardiogram can be constructed from the electrocardiogram if the three standard leads are taken simultaneously. When the termini of the serial positions of the vectors are joined together they will be seen to form a loop. By means of a cathode-ray tube it is possible to record this loop on a fluorescent screen. In pathological conditions the loop takes abnormal forms, and it may be that the vector cardiogram will be found eventually to be a more convenient way of recording myocardial lesions, such as infarcts, than the electrocardiogram.

X-ray Examination of the Heart

X-ray investigation may be made by cardioscopy or by examination of the antero-posterior and two oblique films. Kymography is also used in some centres.

Cardioscopy is as important a part of the examination of the heart as electrocardiography. Not only can a general enlargement of the heart be made out with far greater accuracy than by palpation or percussion, but changes in the shape of the organ enable enlargement of the different chambers to be recognized. The type of pulsations is also important, and those in the left ventricle may vary from the wide excursions of aortic incompetence to the localized absence, or even systolic expansion, found in anterior infarction. In a moving object like the heart much more can be learnt from cardioscopy than from films, which are mainly useful for purposes of record. It is true that measurements made from films enable the transverse cardiac diameter to be calculated, and the relation of this to the internal diameter of the chest can be ascertained; but in doubtful cases tables based upon the weight and surface area of the patient must be consulted, and for ordinary purposes the opinion of an experienced observer during cardioscopy is just as reliable.

Kymography.—In 1912 Gott and Rosenthal described a method of recording the pulsations of the borders of the heart by means of a multiple slit grid, and in 1931 the method was developed by Stumpf. A kymogram preserves as a record the variations in pulsation that can be seen on cardioscopy, and is used in many centres both in the localization of infarcts and in such conditions as constrictive pericarditis, in which all pulsation is minimal.

The Vital Capacity ; Circulation Rate

The vital capacity test was described first by John Hutchinson in 1846 for use in pulmonary diseases, but in 1921 Peabody applied the method to the problems of cardiac dyspnoea. The test is quick and very easy to do: the patient merely breathes out into the spirometer as deeply as he can after the biggest possible inspiration. A young athlete should exceed 5,000 c.cm.: the patient with cardiac failure can barely manage 1,000 c.cm. Obesity, thyrotoxicosis, and psychasthenia all lower the vital capacity. The results can be corrected in accordance with a scale for height and

weight. But much more can be gleaned by watching the test being performed. A normal subject of feeble type merely puffs into the tube, and has to be pressed into producing 1,500 c.cm. or so. The young athlete blows the spirometer up to 5,000 c.cm. or more at once—an all-out effort with a quick ending. The former athlete, now developing emphysema, reaches nearly the same level by squeezing out the last few hundred c.cm. with the utmost determination. Indeed, as Hope Gosse pointed out in a personal communication, this prolonged ending is as good a functional test of emphysema as we have, and in the lower ranges is seen also in patients with heart disease complicated by emphysema. A Hutchinson spirometer is a valuable addition to the armamentarium of the cardiologist as well as of the chest physician.

In 1926 Blumgart and his associates injected a radioactive substance into the brachial vein and were able to detect its arrival at the brachial artery of the other arm. Thus they were able to measure the time taken by the blood to flow up the vein to the heart, through the pulmonary circuit and back to the left side of the heart, and finally out to the arteries of the arm. Many other more simple, but nevertheless less accurate, tests have since been devised for the same purpose, the end-point being obtained either by a taste as in the decholin test, or by a flush as in the histamine test. The ether test measures the time taken from the brachial vein to the lungs; the amyl nitrite test the time from the lungs to the face. Tests of the speed of the blood-flow are helpful only within limits. Apprehension, thyrotoxicosis, and severe anaemia all increase the speed. If in heart failure the speed is slow, and the time lengthened, the end-point may be difficult to make out owing to dilution of the injected substance. Also, the end-point in many of them, such as the decholin and saccharin tests, is subjective, depending upon the appreciation of a taste, and the reaction time of different patients is apt to vary. But the tests take little time and require no apparatus, and within their limits they can provide useful information.

Phonocardiograms ; Venous Pressure ; Catheterization

In 1894 Einthoven devised a method of recording the heart sounds. In 1912 Lewis, by means of sound-tracings, elucidated the nature of the additional heart sound that causes gallop rhythm. Interest in phonocardiograms then lapsed till a few years ago, but now several types of instrument are on the market. Phonocardiograms register the vibrations which produce the heart sounds and murmurs, and they enable these to be timed. Successive records may show the progress of events in one patient. It is not certain yet if the tracings are reliable enough to be submitted as satisfactory evidence of a valvular lesion in cases where oscillations appear and yet no murmur is present. If not, they would not seem to offer any great advantage over auscultation.

It has long been customary to gauge the venous pressure by observing the fullness of the veins in the neck or under the tongue, but direct measurement of the venous pressure dates from 1910, when Moritz and Tabora introduced a large-bore needle into the vein at the elbow and attached it to a saline manometer. Since then various modifications have been devised, of which the most recent has the advantage that a fine needle can be used. The venous pressure provides a measure of the filling pressure of the right side of the heart. In health it varies from 4 to 10 mm. Hg: in heart failure it may rise to 30 mm.

In 1929 Forssman passed a ureteric catheter through his own median basilic vein into his right auricle. He then

walked to the x-ray department and had it photographed *in situ*. In 1932 radio-opaque material was passed through the catheter into the heart by Monis and Carvalho in Spain, and pictures were obtained of the pulmonary vessels. In 1944 Cournand and Ranges left the catheter in position for an hour without detriment to the patient. This new method has led to important advances in three directions: it has enabled the pressures in the right auricle and ventricle to be measured directly; accurate estimations of the cardiac output can now be made; the catheter may be used to introduce radio-opaque material for angiocardiology.

Right Auricular and Right Ventricular Pressures.—In 1946 Cournand attached a manometer to the catheter and was able to measure the pressure in the right auricle and right ventricle. In health the pressure in the right auricle is about zero; in right ventricular failure it may rise to 30 mm. Hg, at which level it is as high as the pressure in the veins. In the right ventricle the systolic pressure is normally about 25 mm. Hg, while the diastolic pressure is about zero. In right ventricular failure the systolic pressure may rise to above 100 mm.

Cardiac Output

To measure the cardiac output by the principle of Fick it is necessary to know the difference in the oxygen saturation of the arterial and venous blood. Since different streams of venous blood returning to the heart vary in their degree of oxygen unsaturation it is an advantage to be able to take samples from the right auricle. To ensure thoroughly mixing, samples are now taken from the right ventricle or pulmonary artery. A sample of arterial blood must also be obtained, and for this an in-dwelling needle has to be inserted into the femoral artery. When the samples have been secured a rather complicated gas analysis is needed in order to calculate the arteriovenous oxygen difference. The test therefore requires a team, each member of which must be skilled in the part he has to play. Estimations of the cardiac output are accurate only if the patient is relaxed, and relaxation cannot always be easy to procure with a catheter in the heart and a needle in the femoral artery. Using this method, McMichael and Sharpey-Schafer have been able to show that while in most forms of heart failure the output is rather low, in conditions such as anaemia, thyrotoxicosis, or emphysema a high output is needed, and congested heart failure may be present while the output is still above the normal level.

The ballistocardiograph, designed by Starr in 1941, provides another means of obtaining the cardiac output. The blood ejected from the heart in systole, as it moves towards the head, makes the body recoil towards the feet. When the column of blood strikes the arch of the aorta the body recoils upwards. The forces of these two movements can be measured. The patient lies upon a table, which is suspended so that all but the longitudinal movements are prevented. The movements of the body on recoil are resisted by a spring, and photographed. The method is not claimed to be as accurate as estimations made by the principle of Fick, but arterial and venous punctures are avoided and so it is more feasible for the patient to relax.

Angiocardiology

In 1937 Castellanos injected "diodone" intravenously into children with congenital heart disease and succeeded in visualizing the different chambers of the heart. Since then the method has been developed by Chávez in Mexico, where the injection is made into a cardiac catheter inserted into the jugular vein, and in the United States.

Angiocardiography provides the same information for the heart that a barium meal does in the stomach. By means of a series of rapid exposures taken at one-second intervals, the opaque material can be seen to fill in sequence the right auricle, the right ventricle, the pulmonary vessels, the left ventricle, and the aorta. If a defect is present in the auricular or ventricular septa the left ventricle and aorta will become opaque earlier than usual. A dextroposed aorta is clearly seen; so is a patent ductus in which the shadow of the pulmonary artery persists owing to the flow of blood into it from the aorta through the ductus. But angiocardiography presents certain difficulties.

The procedure entails the introduction of a large-bore needle or cannula into the vein at the elbow. Unless a general anaesthetic is given, the concentrated solution of diodone causes an unpleasant burning sensation in the head. In adults with thick chest walls it is difficult to get enough penetration to make the outline clear. It is not possible to say how general an application the test may come to have in cardiac diagnosis. So far it has been used in this country mainly in children for whom an operation for the relief of pulmonary stenosis is being considered. It seems likely, however, that improvements in technique will lessen some of the present difficulties and increase the scope of this valuable diagnostic aid.

Volume of the Heart

A method of investigating the volume of the heart was devised by Nylin in Stockholm in 1943. Two simultaneous skiagrams are taken at right angles to each other, and then both the transverse and antero-posterior diameters of the heart can be measured. From these the volume can be calculated. The results obtained compared well with direct measurements of the volume made later at necropsy. In some very large hearts the volume was almost two litres, which is about half the total volume of the blood found in normal subjects. With a cardiac output of about four litres a minute this would mean that the blood would take nearly fifteen seconds to pass through each ventricle, which is not far short of the total time taken by the blood to travel from the vein at the elbow through the right ventricle, the pulmonary circuit, and the left ventricle to the tongue in normal subjects.

Radiocardiography, introduced by Prinzmetal in 1949, allows this emptying time of the heart to be measured. Radioactive sodium is injected intravenously, and the concentration of the particles is obtained by a Geiger-Müller counter and recorded on a revolving drum. In normal subjects the radioactive material passes rapidly into the right ventricle and a large deflection is inscribed. It is then diluted in the pulmonary vessels so that the left ventricular deflection which follows is smaller and more prolonged. When either ventricle is dilated it empties more slowly, and the descending limb is correspondingly lengthened. If the filling pressure is increased the ascending limb will be more oblique. When the left ventricle is at fault, the two deflections may be fused. A delay in the emptying of the right ventricle will cause the blood to pass more slowly into the pulmonary system, and this of itself will lead to a slow upstroke of the left ventricular deflection. Otherwise, the test measures the emptying time and indicates the filling pressure of each ventricle separately, and may be expected to add to our knowledge of the haemodynamics of cardiac failure.

Volume of the Blood ; Capillary Pressure

It might be thought that an estimation of the total volume of the blood was hardly necessary for the solution of

cardiac problems, but this is far from the case. An increase in the blood volume may be the earliest change in the circulation to follow the weakening of the strength of the heart beat. A slight diminution in the flow of blood to the kidneys may lead to retention of salt and thus to an increase in the volume of the blood in order to maintain the osmotic pressure. The first clinical test was introduced in 1937 by Evans, who injected a known quantity of a blue dye. Blood samples, taken when equilibrium had been reached, enabled the dilution of the dose to be estimated, and so the volume of the blood could be calculated. The method is not very accurate and has the disadvantage of turning the patient blue for some time. Radioactive red cells have recently been injected and samples taken from the brachial artery of the other arm. Greater accuracy is claimed, but no results have yet been published in this country. The blood volume, which normally is about four litres, is always increased when clinical oedema is present.

By means of very fine capillary pipettes, Landis in 1930 was able to measure the pressure at the venous and arterial ends of the capillary loop. Normally the positive pressure in the capillaries, which would tend to allow fluid to escape into the tissues, is balanced by the osmotic pressure of the plasma proteins, which holds it back in the vessels. In cardiac failure the raised venous pressure destroys this balance and oedema forms. Complicating factors occur in polycythaemia, when the plasma proteins are increased, and oedema may not be present even though the venous pressure is high. The reverse is seen when the plasma proteins are low, as in nutritional oedema, in which the circulation may be normal.

Procedure

The complete investigation of a patient with heart disease may therefore be a complicated affair: it comprises an electrocardiogram, with standard, unipolar precordial, and limb leads, oesophageal leads, and a vector cardiogram; x-ray examination, including cardioscopy, skiagrams in three positions, and a kymogram; measurement of the vital capacity; phonocardiograms to register the abnormal vibrations due to valvular disease; recording of the arterial, venous, and capillary pressures and of the right auricular and right ventricular pressures by means of a cardiac catheter; cardiac catheterization to obtain the cardiac output if this is not done by means of a ballistocardiogram; measurement of the circulation time; estimation of the volume of the heart with skiagrams, that of the blood by means of the blue dye or with tracers—tracers will also enable the emptying time of each ventricle to be ascertained by radiocardiography—angiocardiograms, in all cases in which a shunt is suspected, to interpret doubtful or unusual shadows seen on cardioscopy or in a skiagram. To these must be added blood tests such as the erythrocyte sedimentation rate in cardiac infarction; the cytology of the blood when anaemia plays a part in the causation of heart failure; a blood culture in infective endocarditis; and the Wassermann reaction. In suspected thyrotoxicosis a basal metabolic rate may be required.

With all this multiplicity of tests one wonders what in the future may be the position of the clinician. He cannot hope to do all the tests on each patient himself. Should he delegate the work of investigation to others? If so, which tests and to whom? The blood examinations fall naturally into the province of the pathologist, but most of the remainder, including cardiac catheterization, seem clearly to come within the scope of the cardiologist. If they were to pass into the province of another department, such as

thoracic surgery, that department would have to assume responsibility for the final diagnosis.

On the other hand, if a team of investigators performs all these tests on every patient with heart disease, is the clinician as we know him really necessary? Reliance upon tests engenders a certain habit of mind in which more confidence is felt in a figure than upon an opinion based on clinical judgment. Indeed, it has been rumoured that in certain more advanced stations the clinician has already been found to be redundant, and his place has been taken by an integrator who, by assessing the results of the investigations statistically, has been able to come to a diagnosis and decide on treatment without seeing the patient at all.

Valuation of Tests

All the masters of medicine have insisted on the importance of an accurate history. Unless the patient is carefully questioned regarding his symptoms diagnosis will suffer. We may therefore dismiss at once the notion that the clinical examination can be superseded. Should we then go to the other extreme and discard all tests, or at least all tests which cannot be performed by the clinician at the time of his examination, as being without value? The answer surely is "No"; but we should discriminate, and by using each test in its proper place avoid making them our masters.

No one should practise cardiology who has not had a thorough grounding in electrocardiography. The electrocardiograph is an instrument of precision. It has great uses but also limitations. In inexpert hands it can be quite dangerous. Hospital cardiographic departments should be entrusted only to those who have been trained in a recognized cardiac department. The same applies to the x-ray examination of the heart. The art of cardioscopy is not to be learnt in a day. Yet a correct appreciation of the size of the different chambers of the heart is almost, if not quite, as important an adjunct of the examination of cardiac patients as is electrocardiography. To these two tests I would add another—namely, the vital capacity—because so much can be gleaned from it and it takes so little time. These three should form part of the routine examination of patients with heart disease, as much as the recording of the blood pressure and the auscultation of the heart sounds.

Mackenzie had nothing but scorn for those who would "use an assistant to take an x-ray photograph of the heart and an electrocardiogram, and even blood pressure readings, and then sit down to study the report." Not much, in fact, is gained by taking an electrocardiogram, or the blood pressure oneself, but the remainder should not be left to subordinates. If one would not accept the findings of another on auscultation, it is difficult to see why his findings on cardioscopy should not equally require verification.

Tests of the speed of the circulation require no apparatus and take little time. They are available to anyone who cares to use them. But the information they give is often disappointing, and it is doubtful if they are worth including in a routine examination.

Accurate measurements of the venous pressure are helpful in many ways both in the prognosis of cardiac failure and in assessing the results of treatment. When the apparatus with a fine needle is generally available, venous pressure measurements will probably be included in the examination of patients with heart failure.

The exploratory work on angiocardiology has so far been done mainly by cardiologists, but the problems con-

cerned are almost entirely radiological. A series of rapid exposures have to be made with sufficient penetration to visualize the bolus of diiodone as it passes through the heart. Probably, as time goes on, the technique will fall more and more into the field of the radiologist, and cardiologists will ask for the additional information in appropriate cases. Angiocardiography will then have to be made available in all hospitals where a cardiac clinic is established.

It is a regrettable fact that when surgeons invade a province hitherto considered to be purely medical, diagnosis almost at once becomes more exact. Not so long ago the diagnosis of congenital morbus cordis was considered quite adequate by paediatricians. It is true that the development of cardiac radiology directed the attention of cardiologists to the congenital lesions years before any operative intervention was considered feasible, and main types, such as the tetrad of Fallot, valvular pulmonary stenosis, atrial septal defect, coarctation, and the patent ductus, were recognized clinically. But the advent of thoracic surgery, and especially of the Blalock-Taussig operation, has opened a new field, and now pulmonary atresia, tricuspid atresia, and major and minor degrees of transposition are distinguished. For this, angiocardiology and cardiac catheterization are often required.

Cardiac catheterization is difficult and laborious. To obtain a thorough mixing of the venous blood the catheter may need to be pushed as far as the pulmonary artery. This takes time, and there are many pitfalls, as the tip of the catheter may double back and appear in unexpected places. If the catheter passes from the right auricle to the left or from the right to the left ventricle the pressures will alter and the samples will change from venous to arterial blood. Hence the importance of the procedure in establishing the presence of shunts in congenital heart disease. Otherwise it is useful chiefly owing to the possibility of observing directly the changes in pressure of the different chambers of the heart, and changes in the cardiac output, caused by heart failure or induced by drugs. It would seem probable that cardiac catheterization will be performed only in hospitals large enough to have a thoracic surgical unit or at research centres.

Most of these tests were devised for purposes of research. They were developed at single centres, and were later adopted generally because they were found to be useful in diagnosis. Other methods are still in the stage of development. They may or may not come to have a general application. Such are the ballistocardiogram, the radiocardiogram, and the measurement of the volume of the heart.

The ballistocardiogram enables the cardiac output to be obtained without the unpleasantness of a cardiac catheter and an arterial puncture. But estimations of the output do not help much in the diagnosis and treatment of patients with heart disease. Those with "high-output failure" have also fast circulation rates, and the circulation time is an easy test to do. In ordinary congestive heart failure the output falls only a little, and estimations of the cardiac output are unlikely ever to form part of the routine investigation of such patients.

The radiocardiogram and the estimation of the volume of the heart have opened up a new approach to the problems of heart failure. They have shown that the dilated heart empties itself slowly. With the radiocardiogram the emptying time of each ventricle can be measured separately. The test is very recent, but it is certainly important, and we may expect to hear more of it in the future.

A Suggestion

Almost all of these various techniques have been devised either in Germany or, more recently, in America. Since the war we have been engaged in making up for the lost years and bringing ourselves up to date. A major difficulty has been the necessity for importing all the apparatus required, most of which is expensive and has to be paid for in dollars. The result is that only a little is available in a few centres. We would not wish this state of affairs to continue, but should prefer to regain the position we held during the time of Lewis, developing our own lines of research and constructing for ourselves the apparatus we need. The problem is how this may best be done, having regard to the probable financial state of the country during the next few years.

The Medical Research Council finances some specific investigations and supports research establishments at certain centres, such as at University College Hospital, where Lewis worked. But this does not seem to be enough. We must encourage research at all levels, in small hospitals as well as large. Some men work better in a team, others when left to pursue their own lines of thought. We should not forget that Mackenzie revolutionized the ideas of his time while engaged in general practice, and was not looked upon with any favour by the authorities whilst doing so. Even Lewis had to buy his own electrocardiograph. If research is confined to a few centres, only those projects that commend themselves to the directors of the centres may receive attention.

A possible solution might be for the specialist organizations to have some share in the control of the funds to be expended, or else they might form advisory committees of the Medical Research Council. For research in cardiology the Editorial Board of the *British Heart Journal* might be appropriate. A man who wished to investigate some point could submit his plans, together with his needs in regard to apparatus; and, if the project was approved, arrangements could be made for the apparatus to be constructed for him. Promising lines of research would thus be scrutinized by those who later would be responsible for accepting the work for publication. If this method were combined with the normal research carried out at approved centres, a man with an original mind should find sufficient opportunity to try out his ideas.

New techniques are designed primarily in order to advance knowledge. As such they are to be commended. No country that is not active in research can be considered to be in the forefront of the battle against disease. Tests, too, lead to greater precision in diagnosis and treatment. Some, such as electrocardiography and cardioscopy, are in daily use; others, such as angiocardiology and radiocardiography, may well in the future be required as a routine in selected cases. Cardiac catheterization offers a ready means of assessing the efficacy and mode of action of cardiac drugs. The danger of tests lies in the risk that undue reliance may be placed upon them at the expense of the clinical examination, leading eventually to the stage when an electrocardiogram is required before a normal heart can be diagnosed. If ever the National Health Service were controlled by a bureaucracy these tests might be demanded as part of the opinion by a specialist. It is to be hoped, therefore, that advice will continue to be accepted from medical committees, whose members represent the best interests in the profession, so that tests may not be allowed to exceed their proper function as the tools of clinical medicine.

In the historical survey free reference to *A History of the Heart and Circulation* by Willius and Dry (1948) has been made. Other references are given below.

REFERENCES

- Blumgart, H. L., and Weiss, S. (1927). *J. clin. Invest.*, **4**, 15.
 Castellanos, Pereira, and Gacia (1937). *Arch. Soc. Estud. clin. Habana*.
 Chávez, I., Dorbecker, N., and Celis, A. (1947). *Amer. Heart J.*, **33**, 560.
 Courmand, A., and Ranges, H. A. (1941). *Proc. Soc. exp. Biol., N.Y.*, **46**, 462.
 Einthoven, W. (1903). *Pflug. Arch. ges. Physiol.*, **99**, 472.
 — and Geluk, M. A. J. (1894). *Ibid.*, **57**, 617.
 Floyer, J. (1710). *The Physician's Pulse Watch*. London.
 Forssman, N. W. (1929). *Klin. Wschr.*, **8**, 2085.
 Gibson, J. G., and Evans, W. A. (1937). *J. clin. Invest.*, **16**, 301.
 Gött, T., and Rosenthal, J. (1912). *Münch. med. Wschr.*, **59**, 2033.
 Harvey, W. (1628). *Exercitatio Anatomica de Motu Cordis et Sanguinis in Animalibus*. Frankfurt.
 Heberden, W. (1772). *Med. Trans. Coll. Phys. Lond.*, **2**, 59.
 Hutchinson, J. (1846). *Med.-chir. Trans.*, **29**, 137.
 Laennec, R. T. H. (1819). *De l'auscultation médiate*. Paris.
 Landis, E. M. (1930). *Heart*, **15**, 209.
 Lewis, T. (1911). *Mechanism of the Heart Beat*. London.
 Mackenzie, J. (1902). *The Study of the Pulse*. Edinburgh.
 McMichael, J., and Sharpey-Schafer, E. P. (1944). *Brit. Heart J.*, **6**, 33.
 Monis, E., Carvalho, L. de, and Saldanha, A. (1932). *J. Radiol. Electrol.*, **16**, 469.
 Morgagni, J. B. (1761). *De Sedibus et Causis Morborum per Anatomen Indagates Libri Quinque*. Venetus.
 Moritz (1902). *Münch. med. Wschr.*, **49**, 1.
 Nylin, G. (1943). *Amer. Heart J.*, **25**, 598.
 Oliver, G. (1901). *Blood and Blood Pressure*. London.
 Pardee, H. E. B. (1920). *Arch. intern. Med.*, **26**, 244.
 Peabody, F. W. (1921). *Med. clin. N. Amer.*, **4**, 1655.
 Prinzmetal, M., Corday, E., Spritzler, R. J., and Flieg, W. (1949). *J. Amer. med. Ass.*, **139**, 617.
 Riva Rocci, S. (1896). *Gaz. med. Torino*, **47**, 981.
 Schellong, F., Heller, S., and Schwingel, E. (1937). *Z. Kreisforsch.*, **29**, 497.
 Starr, I. (1941). *Amer. J. med. Sci.*, **202**, 469.
 Stumpff, P. (1934). *Fortschr. Röntgenstr.*, **49**, 211.
 Vieussens, R. (1715). *Traité Nouveau de la Structure et des Causes du Mouvements Naturel du Cœur*. Toulouse.
 Wilson, F. N., Johnston, F. D., MacLeod, A. G., and Barker, P. S. (1934). *Amer. Heart J.*, **9**, 447.
 Withering, W. (1785). *An Account of the Foxglove*. Birmingham.

TUBERCULOUS EMPYEMA

BY

D. A. ANDERSEN, M.D., F.R.C.S.

The purpose of this paper is to present the results of a "follow-up" of a series of 74 cases of tuberculous empyema treated in a period of seven years, from 1941 to 1947, in Norwegian hospitals. With six exceptions the patients were all admitted to the Third Department of the Ulleval Municipal Hospital, Oslo, under the care of Dr. Carl Semb. They came from other hospitals or sanatoria, or were referred by chest physicians in various parts of Norway.

Details of Follow-up Method

Every case of tuberculous empyema discharged from the department between Jan. 1, 1941, and Dec. 31, 1947, is included in the series. In 19 the condition of the patients did not allow of any major surgical treatment; in 55 a plastic operation on the chest wall was carried out.

According to information obtained up to May, 1948, 25 have died. Of the 49 survivors 43 have been followed up fully and their condition ascertained; in six cases the follow-up was incomplete. Where possible the patients attended Ulleval Hospital and 16 were examined by me. The local and general condition, the presence or absence of a fistula, the vital capacity, the sedimentation rate, and the presence or absence of tubercle bacilli in the sputum were determined, together with a straight radiograph in each case and a planogram series when indicated. The