REFRESHER COURSE FOR GENERAL PRACTITIONERS

THE ACUTE THROAT

BY

T. ANDERSON, M.D., F.R.F.P.S., F.R.C.P.Ed.

Lecturer in Infectious Diseases, Glasgow University

The complaint of "a sore throat" must be one of the commonest made to the doctor, and as the condition sometimes calls for urgent treatment the correct diagnosis is of some importance. Although the initial complaint focuses attention upon the throat it is necessary to remember two fundamental points: first, a throat infection may exist without sufficient pain to draw notice to its presence, which bears out the old clinical rule that examination of a patient—particularly a child—is not complete until the throat has been inspected; secondly, it should never be forgotten that the throat inflammation may be the local manifestation of some general disease.

Examination of the Throat

The examination of the throat is such a simple procedure that a description of the necessary manipulations is often taken for granted. Yet to obtain a proper inspection in a writhing youngster calls for tolerance and persistence on the part of the doctor. With patience and persuasion the co-operation of the child will often be fully attained, and where this course is possible it is much to be preferred. No attempt should be made to use force. The following points may be of assistance in difficult circumstances.

(1) *Holding the Child.*—When there is complete refusal to co-operate it will be necessary to pinion the child. The best method is to bring both arms across the chest and for an assistant to hold them there by catching both wrists in one hand. The assistant's other hand should hold the head firmly on the bed by placing the open palm across the forehead, the thumb to one side and the fingers to the other. In such a way the child may be held firmly and painlessly.

(2) *Opening the Mouth.*—Sometimes an unwilling child will hold the jaw tightly clenched. The spatula should then be inserted along the left-hand side of the mouth between the teeth and the buccal mucosa. Firm pressure inwards as far back as possible on the lateral aspect of the alveolar margin will usually result in the mouth being opened. The spatula can then be rapidly turned flat and passed on to the tongue.

(3) *Position of the Spatula.*—The spatula should be passed far enough into the mouth for the distal end to cover the posterior third of the tongue. Pressure should be downwards and forwards to enable a full view to be obtained of the fauces without forcing the tongue back into the throat, which not only blocks the airway but causes the child to "gag."

Diagnostic Points

When examining the throat the doctor must ask himself the following questions:

(1) *Is the Throat Acutely Inflamed?*—The soft palate, faucial pillars, tonsils, and pharyngeal walls should be carefully inspected for redness and oedema. The lesion may vary from a slight reddening of the palatal fringe, uvula, and anterior pillars to oedema so gross that the

swollen uvula, palate, and tonsils almost close the faucial airway.

(2) *Is there an Inflammatory Exudate?*—This may be unilateral or bilateral, and three stages in its formation may be recognized. First comes the appearance of numerous discrete follicular flecks; secondly, there is a confluent, soft, loosely attached and rather pultaceous exudate; and, thirdly, one sees a confluent, firm, leathery membrane which is more or less firmly attached to the subjacent mucosa. Note that exudate is usually confined to the tonsillar surfaces, but membrane may extend on to the fauces, soft palate, or pharyngeal wall.

(3) *Is the Lesion Raised?*—When exudate or membrane is present its level relevant to the adjacent mucosa must be noted. The doctor must satisfy himself whether it is superimposed upon the mucosa or level with the adjacent surface, as if forming the base of a shallow ulcer.

(4) *Is the Membrane Discrete?*—It may be one continuous piece, or made up of numerous solitary areas with intervening normal mucosa.

The causes of a sore throat may be grouped under three broad headings—local virus infections, local bacterial infections, and general diseases.

Virus and Bacterial Infections

Influenza may start with a complaint of sore throat. In such cases the throat often shows little abnormality apart from some enlargement of the solitary follicles on the pharyngeal wall. There seems little doubt, however, that an upper respiratory virus infection can produce a generally inflamed throat with or without tonsillar exudate exactly similar in appearance to that of an acute bacterial infection.

Coccal Infections.—It is impossible to differentiate clinically between the various acute coccal infections. In general *Str. pyogenes* produces a severely inflamed and oedematous throat usually of sudden onset and associated with such great pain that swallowing is exquisitely sore. A tonsillar exudate is common with streptococcus, pneumococcus, or staphylococcus infections. This exudate may become confluent and closely simulate a false membrane, but it is more friable and white or yellow in colour. The associated lymph glands are enlarged and tender but remain discrete. In such cases the ear drums should always be examined. The drum-head may show a diffuse pinkness and slight lack of lustre that suggests a coccal tonsillo-pharyngitis.

Scarlet Fever.—A sore throat precedes the onset of scarlet fever, and the initial rash is often seen on the soft-palate, giving it a speckled appearance. The skin rash will usually confirm the diagnosis within 18–24 hours.

Pharyngeal Abscesses.—A peritonsillar abscess may develop on top of a pre-existing acute tonsillitis. The unilateral, tense swelling of the fauces and soft palate, pushing tonsil and uvula to the opposite side, is easily recognized, and the intense pain is a characteristic feature. Retropharyngeal abscess can be difficult to diagnose,

particularly in young children, because of the filling of the pharynx with mucus as a result of the difficulty in swallowing. The swelling produces a curious choking stridor which may be mistaken for laryngeal obstruction, but the latter produces a much "drier" stridulous with harsh cough. The fauces may show little inflammation in such cases, and should the character of the respiration have aroused suspicion there is only one course of action—persistence in obtaining a good view of the pharyngeal wall.

Diphtheria

A few years ago diphtheria would have taken first place in any consideration of the differential diagnosis of a sore throat. As a result of a reduction in incidence, largely due to active immunization, the clinician need not to-day regard it as occupying so high a place on the list. But it is well to keep two considerations in mind. First, in face of an appearance which suggests diphtheria a history of immunization should never encourage the practitioner to consider the diagnosis impossible. Secondly, it must be remembered that in the "immunized" subject a diphtheritic infection may produce an appearance indistinguishable from that of follicular tonsillitis. Typically, infection of a susceptible person with *C. diphtheriae* has an insidious onset and produces a mildly inflamed throat; the greyish membrane is "all of a piece" and often shows a slightly curled edge which may spread on to soft palate, faucial pillars, or pharyngeal wall when diagnosis is delayed. In general, the inflammatory reaction is limited to a zone surrounding the membrane and the diffuse bright redness characteristic of a streptococcal infection is seen less commonly. Pain is not a feature—indeed, it is often surprising to find how little pain may be associated with an easily recognized, severe diphtheria. Adenitis of the tonsillar glands, so often referred to in textbooks, is the mark of the severe or late case and is of little assistance in making an early diagnosis.

The most confusing appearance is produced by a severe *gravis* diphtheritic infection. The inflammation is intense and the soft palate and anterior aspect of the faucial pillars may be very oedematous and covered with a thin glairy mucus which is not unlike white of egg. This is, in fact, early membrane. The typical membrane may be slight or absent and the condition carelessly regarded as a quinsy. When diphtheria is epidemic the clinician should never hastily use a knife on what seems an acute peritonsillar abscess, for if the lesion is in reality diphtheritic incision greatly worsens the prognosis.

The diagnosis of diphtheria is never easy and even the expert can define no absolute diagnostic criteria. It should be more widely appreciated that modern diphtheria antitoxin is a highly refined and concentrated product, 8,000 units of which can be obtained in a 2 ml. syringe. When the clinician is in any doubt a dose of this magnitude is a safe procedure and will be adequate for most early cases. Serum reactions are now so rare as to be negligible.

Vincent's Angina.—This condition, like diphtheria, has an insidious onset, and as a rule produces little rise in temperature. The throat may simulate diphtheria in that the slough covering the ulcers suggests membrane formation. It will be observed, however, that the membrane is on a level with the normal mucosa and that it makes up several discrete, linear lesions separated by more or less normal mucosa. Although ulceration is seen in rare cases of diphtheria the lesion is continuous and does not consist of individual ulcers. In Vincent's angina the whole mouth is often unhealthy and the gums are inflamed and swollen—a further aid to diagnosis.

Sore Throats in Schools

In closed communities such as boarding schools, outbreaks of acute tonsillitis are most frequently streptococcal in origin. The cultures from the initial cases will usually decide which organism is responsible, and as a rule the swabs of throat and nose produce a heavy, almost pure, culture. Early diagnosis and isolation of those infected are the measures most likely to limit spread, but the high carrier rate which often accompanies such an outbreak may entail the isolation of a large proportion of the individuals. To begin with, all the residents must be subjected to a clinical examination at which particular attention is paid to ears, throat, and nose for the presence of otorrhoea, enlargement or irregularity of tonsils, or rhinitis. Those who show such abnormalities should have cultures taken from nose and throat and should be separated from the others; for it is this group which usually contains the carriers. The remainder should be inspected daily for a period of seven days and any individual showing signs of upper respiratory tract infection isolated at once. When cases continue to crop up among those who showed no abnormality of throat, nose, or ear two possibilities should be considered: first, that the epidemic is food- or milk-borne, and, secondly, that there is an unsuspected carrier remaining among the apparently healthy residents. In connexion with the latter possibility it is important to remember that the nasal carrier is the more dangerous.

When tracing the source of an epidemic which may involve a considerable number of persons the doctor should make early contact with the local medical officer of health and the nearest unit of the Public Health Laboratory Service. Precise knowledge of the organism involved is essential for any useful epidemiological work. For example, by determination of the serological type in streptococcal outbreaks it is possible to make more accurate decisions regarding the importance of a carrier or to assess the significance of milk or food in the spread of infections. Early consultation with the bacteriologist will ensure not only that he receives the right kind of specimens but also that he is able to proceed with the further definition of the pathogen involved.

Specific prophylaxis is only practicable in outbreaks of diphtheria, when combined active and passive immunization should be started. This consists of the injection of 500–1,000 units of diphtheria antitoxin into one arm and 0.5 ml. of A.P.T. (alum precipitated toxoid) into the other. Six to eight weeks later a second injection of 0.5 ml. of A.P.T. is given. Prophylaxis of streptococcal infections by the administration of sulphonamides is not advisable as a routine measure.

General Infections

Glandular Fever.—The anginose form of infectious mononucleosis produces a throat which may be indistinguishable from an acute coccal infection—indeed, the secondary invader is often a *Str. pyogenes*. Glandular fever is a systemic virus infection. Generalized lymphoplasia and splenomegaly will probably be present, but the diagnosis may be suggested by the sore throat suggestive of streptococcal tonsillitis appearing as a secondary manifestation after a few days of general malaise. In the usual streptococcal infection the sore throat is the first manifestation of an acute illness. There is little doubt that infectious mononucleosis now occurs more frequently than it did in the past. Accurate diagnosis depends upon the differential white cell count, which shows a mononuclear

leucocytosis and the characteristic atypical cells, and the demonstration of heterophile antibody in the patient's serum by the Paul-Bunnell reaction. This reaction is variable in the time of its occurrence and often becomes positive for the first time late in the illness. A titre of 1/64 is significant. Should the patient have received horse serum in the form of an antitoxin the bacteriologist must be informed, so that he may take steps to eliminate the resulting false-positive reaction.

Agranulocytic Angina.—The throat may show ulceration and membrane formation. There is, however, a characteristic lack of inflammatory reaction in contrast to the apparent acuteness of the lesion, and the soft palate, uvula, and fauces look pallid, soft, and jelly-like. The lesion is more frequently seen in adults; and a history will often be obtained of the habitual use of a drug containing the benzene ring. The amidopyrine group and, rather infrequently, the sulphonamides constitute the commonest aetiological agents. Diagnosis depends on lack of polymorphs seen on a carefully taken blood film. The film should preferably be made on cover slips to secure an even distribution of the white cells. If it is taken on a slide the most careful scrutiny should be made along the edges and at the end of the smear, for it is here that the polymorphs will usually be found.

Toxic Drugs.—Whether agranulocytosis is in fact a form of acute sensitization or not is still uncertain, but there is no doubt that an ulcerative stomatitis and pharyngitis without agranulocytosis can occur as part of a drug sensitization. The whole mouth is inflamed in this condition and as a rule there is some confirmatory evidence such as a rash. It is important to appreciate that the individual who has been sensitized to one drug is the more easily sensitized to others, and care should be exercised in the use of forms of therapy involving sensitizing drugs, such as aspirin and the sulphonamides.

Other Conditions.—Acute leukaemia sometimes presents for the first time as an acute ulcerative tonsillitis. The child—for it is more frequently seen in childhood—usually appears ill and anaemic. There may be petechial skin haemorrhages, the spleen is often enlarged, and sternal tenderness is usual. Diagnosis can only be made after a careful examination of the blood.

Finally, in the adult, secondary syphilis should always be borne in mind as a cause of ulcerative tonsillo-pharyngitis. The presence of a typical rash may give a pointer to the correct diagnosis, but failure to respond to any form of treatment is often the first feature to make the clinician think twice about the diagnosis. Toxaemia usually appears mild in contrast to the severity of the local lesion. The Wassermann reaction will confirm the diagnosis.

Bacteriological Procedures

In the past, the focus of attention upon diphtheria resulted in the clinician tending to seek bacteriological assistance only when this diagnosis was under consideration. A carefully taken swab from the tonsillar surfaces, however, is of value in all throat infections and is particularly desirable in severe cases because of the guidance it may give in planning treatment. The swab must be taken before any form of chemotherapy is used. Most bacteriologists include a direct film in their examination of the throat swab, as well as culture on blood agar, Loeffler's slope, and a special medium to differentiate the organisms of diphtheria. The report may therefore supply valuable information in regard not only to diagnosis but also to prognosis.

Treatment

Antitoxin.—Although in many cases treatment will be guided by the results of the bacteriological examination, it is often necessary (and in a suspected case of diphtheria essential) to start treatment before the results of such examination are known. The dose of diphtheria antitoxin ranges from 5,000 to 100,000 units. Where the disease is suspected an intramuscular dose of 8,000 units will suffice. For the late, severe case when hospital admission cannot be immediately secured a dose of the order of 50,000 units will be required, and in such an event intravenous administration is particularly desirable. Such cases are best treated in hospital, however, and the practitioner will find that most fever hospitals are anxious to admit them even when the diagnosis is unconfirmed or in doubt.

Penicillin.—For the coccal infections and for Vincent's angina penicillin is the treatment of choice. It should be realized that the need is to secure an adequate local concentration of the antibiotic. The condition must not be thought of as a mere surface infection. Local therapy by lozenges and so forth is therefore completely inadequate and systemic administration in high dosage is indicated. If the aqueous solution is used a four-hourly dose of 100,000 units should be injected intramuscularly: procaine penicillin may be employed to advantage here, and two or three injections of 500,000 units during the first 24 hours may be followed by a lower dosage for the remainder of the time. For all throat infections due to susceptible organisms treatment must be continued for seven days to secure the best results.

Although there is no specific treatment for infectious mononucleosis the throat infection is usually due to either *Str. pyogenes* or Vincent's infection, and penicillin will prove effective. In agranulocytic angina, also, the administration of penicillin has greatly improved the outlook. Pentonucleotide should be injected intramuscularly. During the first 24 hours three or four injections of 5–10 ml. may be given, and thereafter the dosage will depend on clinical response. The administration of the antihistamine drugs might be considered, but the results have not been encouraging. In severe cases repeated small blood transfusions are valuable. The diet should be plain and contain no flavourings or adulterants, and care should be taken to avoid the use of any chemical which might have a sensitizing action.

Sulphonamides.—The administration of sulphonamides has been intentionally omitted from the treatment of the acute throat. The efficacy of these drugs has never been convincingly demonstrated in such conditions. Since they are undoubtedly toxic materials, carrying appreciable risks of sensitization, their administration is inadvisable, and in certain conditions may even be dangerous.

Finally, it must be pointed out that many mild throat infections clear rapidly and effectively without specific treatment. Gargling is so frequently indulged in by the patient before seeking advice that its continuation may be advised for its effect in cleansing the mouth. For those adults who can gargle, warm water with glycerin of thymol may also prove soothing to the throat. When pain is more severe the application of poultices to the neck, inhalations of friar's balsam, and, of particular value, the administration of aspirin powder, which is allowed to melt slowly in the mouth before swallowing—such measures are often adequate to secure rapid relief. The hasty rush for penicillin as a panacea for all cases is to be deprecated. It may do the patient little harm, but it can induce in the physician an easy optimism which may eventually cloud his whole judgment.