REFRESHER COURSE FOR GENERAL PRACTITIONERS

FAILED FORCEPS

BY

F. J. BROWNE, F.R.C.S., F.R.C.O.G.

Emeritus Professor of Obstetrics and Gynaecology, University of London

The term "failed forceps" is applied to the type of obstetric case in which an attempt has been made to deliver with forceps but without success. The causes of failure have been fully discussed in papers in this *Journal* and elsewhere by Phillips, Farquhar Murray, Miller, Fletcher Shaw, Stacey, and Claye, and are chiefly as follows.

Contracted Pelvis and Disproportion

In one-third of Miller's 558 cases of failed forceps this was the cause, either alone or combined with a posterior position of the occiput. The difficulty arises mostly from failure to recognize insuperable disproportion between the foetal head and the pelvis, or because in cases of moderate disproportion the forceps is applied too soon and before sufficient time has been given for the head to mould and to descend far enough into the pelvic cavity. Incredible as it may seem, in over 50% of Miller's cases the head was still free above the brim on admission to hospital after attempts at forceps delivery outside had failed. One of the elementary rules regarding the application of forceps had thus been broken.

The most frequent cause of disproportion is contracted pelvis. Severe degrees of contracted pelvis due to rickets are now rare, but minor degrees of contraction, especially of simple flat and generally contracted pelvis, are common and apt to be overlooked before labour sets in if certain simple rules are neglected.

(1) In a primigravida the foetal head should engage deeply in the pelvis in the last two weeks of pregnancy. If it does not do so a reason must be sought. A common cause is contracted pelvis, and this condition should always be excluded. This may be done by simple clinical methods—measuring the patient's height (under 5 feet (1.5 m.) is always suspect), looking for rickets, and measuring the diagonal conjugate—but the present tendency is to use x-ray pelvimetry, which alone can give accurate measurements of the diameters, including the true conjugate, and the interspinous diameter of the outlet. Disproportion, which is rare in the absence of pelvic contraction, can be assessed by Munro Kerr's modification of Muller's method, with or without anaesthesia.

(2) In multigravidae the history of past labours may give warning of possible trouble. For example, in one of the cases of failed forceps recorded by Fletcher Shaw the woman had had four previous confinements, two of which ended in craniotomy and two in caesarean section.

Unrecognized Unrotated Posterior Positions of the Occiput

This occurs mostly in primiparae. In all the series referred to above it was the second most frequent cause of failure. It is necessary to stress once more the importance of a thorough examination of all patients before applying the forceps, first by abdominal palpation and then per vaginam, if necessary with the whole hand in the vagina under deep anaesthesia. It is necessary to make sure of the following points: (1) that the os is fully dilated; (2) that the occiput is anterior; (3) that the head has

descended at least to the level of the ischial spines. These can be easily felt on each side and mark the level of the pelvic floor. Application of forceps to a head above the spines is a "mid forceps" application, below this a "low forceps."

Abdominal examination will reveal much. It will show how far the head has descended into the pelvis. If any part of the head is felt on abdominal palpation delivery is likely to be difficult and may even be dangerous for the child. Occipito-posterior positions cannot usually be diagnosed abdominally but only by introduction of the whole hand into the vagina under deep anaesthesia.

There are two ways of diagnosing persistent occipito-posterior cases. The first and most commonly used is to feel the ear—its free border is directed towards the occiput. It should be remembered that the ear may be folded forward, and this may give rise to doubt. Another plan is to feel for the nape of the neck. When this is found—a smooth flat surface—the forefinger can be passed a little farther on to the anterior shoulder and the head and body rotated by the internal hand alone in one single movement, if possible without dislodging the head out of the pelvic cavity. There is only one fallacy that I have found in this method of diagnosis: if the head is much moulded the forehead may be so elongated that it is mistaken for the neck. This error can be avoided by keeping the possibility in mind and pushing the exploring hand a little farther on. If it is the forehead the finger will come to the face instead of the flat back and shoulder. It should be noted that slipping of the forceps is almost always due to its application to an unrotated occipito-posterior. If this happens the blades should be removed and the position reassessed.

Application through an Undilated Cervix

In 50% of Claye's cases this was the cause of failure. As every student is taught that full dilatation is necessary before application of forceps the mistake can only arise through difficulty in diagnosing full dilatation. Claye has a useful word on this. "If," he says, "the cervix is fully dilated the head has come so far through it that its rim is no longer palpable. If the cervix is palpable it is not fully dilated." To this the writer would add that "full dilatation" means full dilatation between pains. Too often the medical attendant examining the patient (as is correct) during a contraction and finding the head descend and no rim of cervix palpable concludes that the os is fully dilated and decides to apply forceps. When the patient is anaesthetized, however, he may find that the os is only three-quarters or less dilated. By that time all preparations have been made for forceps, and if inexperienced the doctor may not have the moral courage to desist and confess his error. It is always wise, therefore, in assessing dilatation of the cervix before application of the forceps, to keep the examining finger in the vagina after the pain has passed off.

Application of forceps through an imperfectly dilated cervix most commonly takes place in primiparae in whom the first stage of labour has been prolonged, and especially

if the membranes have ruptured prematurely. These are always anxious cases. There may be disproportion or an occipito-posterior position, in both of which cases labour tends to be slow and pains poor. But often neither of these is present and the delay is then generally due to primary uterine inertia. If the membranes are unruptured the long first stage does not matter much, though it tries the patience of all concerned. If the membranes have ruptured prematurely, however, the life of the foetus is then endangered, because as liquor trickles away the uterus is allowed to retract excessively and progressively diminish the area of attachment of the placenta and the foetal and maternal circulation through it. If it proceeds far enough this will cause foetal asphyxia and intrauterine death. These cases can be very anxious ones, and little can be done except to give a sedative, keep the bowels and bladder empty (a warm soap-and-water enema often stimulates strong contractions), and avoid fussy and repeated vaginal examinations. The foetal heart sounds should, of course, be examined at intervals. Most of these labours end spontaneously or the patients are easily and safely delivered by low forceps; but in a few cases anxiety arises because the foetal heart sounds indicate foetal distress before the cervix is fully dilated. The best course to adopt then needs both experience and good judgment. It is often possible to complete the dilatation of the cervix manually and apply forceps, but this is always a hazardous procedure for the child, especially as the dilatation is apt to be imperfect. If the cervix is thinned out over the head it may be divided at 5 and 7 o'clock by scissors after applying the forceps and drawing the head gently downwards. But more and more of these cases are now being treated by lower-segment caesarean section, which is usually much safer for both child and mother.

Constriction Ring

This is one of the more infrequent causes of failure. It may be the only cause or it may complicate contracted pelvis or unrotated occipito-posterior cases. Usually the head descends fairly well when forceps traction is made, but it quickly recedes again when traction is stopped. This should make one suspicious, but the condition can only be diagnosed by removing the blades, inserting the finger, and feeling the ring as hard as iron around the neck. In such cases the forceps should be reapplied and one capsule (3 minims: 0.18 ml.) of amyl nitrite crushed on gauze under the patient's nose so that she may inhale it. Relaxation takes place immediately, but is temporary. The forceps should therefore be in place for delivery without unnecessary delay. An alternative method is to inject subcutaneously 7 minims (0.62 ml.) of 1 in 1,000 adrenaline.

Finally, a certain number of failed forceps cases arise through application of the instrument to a hydrocephalic head, an undiagnosed breech, a mento-posterior or brow presentation, or when there is obstruction by a pelvic tumour.

Should the Forceps be Applied or Not?

On each blade of the forceps there should be a large question mark. On taking up the first blade and seeing the question mark the operator should ask himself, "Are the conditions present that make the application of forceps safe—the membranes ruptured, a vertex presentation with the occiput anterior and the sagittal suture in the antero-posterior diameter, the bladder and bowels empty, the lower pole of the head at least at the level of the ischial spines, the os fully dilated?" On taking up the second blade he should ask himself, "What are the indications justifying the application of forceps in this case?" Apart from the

case when the head is on the perineum, when application of forceps (preferably preceded by episiotomy in a primipara) is always safe, there are only two real indications for forceps—maternal or foetal distress.

Finally, we should all of us keep in mind the words of our master, Smellie. "Students should never think themselves perfect; for after all the instruction that can possibly be conveyed there are many things in midwifery that can only be learned by practice and observation; and cases will sometimes arise which will puzzle and foil the best practitioners."

BIBLIOGRAPHY

- C'aye, A. M. (1931). *Practitioner*, 126, 611.
 Miller, D. (1928). *British Medical Journal*, 2, 183.
 Murray, E. Farquhar (1927). *Clinical J.*, 56, 488.
 Phillips, L. (1926). *Lancet*, 1, 113.
 Shaw, W. Fletcher (1928). *British Medical Journal*, 2, 188.
 Smellie, William (1774). *A Collection of Cases and Observations in Midwifery*, vol. 3, 6th ed. London.
 Stacey, J. C. (1931). *British Medical Journal*, 2, 1073.

Preparations and Appliances

HOSPITAL SYRINGE SERVICE

Mr. P. G. MERRETT, theatre technician, St. Thomas's Hospital, writes: In modern anaesthesia the use of 2-ml. syringes both for immediate and for intermittent use during operations seems to have increased with the introduction of curare and its antidotes. I thought, therefore, that a method of packing syringes ready for supply at a moment's notice would be acceptable, and have devised the following contrivance, which has met with the approval of most of the anaesthetists at this hospital. The type of sterilization is by dry heat.

As the accompanying diagram shows, the requirements are heat-resistant syringes, oven-glass tubes, "cellophane," and a small rubber band to seal and secure the end of the tube when ready for sterilizing. Care of the needle-point is important, and for this purpose use is made of a piece of the brass needle-holding tray supplied by most manufacturers with new needles. The unwanted clips are flattened, leaving one for each guard, and cut out as shown. This guard is easily clipped to the needle-butt and hand-shaped so as to be deflected from the point of the needle. The tapered end is curved inwards slightly to fit the bottom of the glass tube, and serves as a protection to the whole. This method has stood up to normal use. If required, an absorbent cotton roll, as used for dental purposes, can be inserted alongside the needle. This provides a swab for cleaning the area of injection, and can be used to protect the needle if the syringe is required for intermittent injection. Most anaesthetic trolleys could carry a small bottle of surgical spirit.

There is no doubt the method could be adapted for other branches of medical, surgical, and laboratory work, and it might be handy in a central supply depot.

A set of pictures for public display has been prepared on "Coughs and Sneezes" by the Central Office of Information. Local authorities may obtain these sets from the C.O.I. (Room 2301), Block 2, Montagu Mansions, Crawford Street, Baker Street, London, W.1.