It is reasonable to suggest, therefore, that abnormal sensitivity to streptomycin or a direct poisoning by the drug is the most probable cause of the symptoms.

For permission to publish the account of these cases I am indebted to the Director of Medical Services, Federation of Malaya.

REFERENCES

- Beham, H., and Perr, H. (1948). *J. Amer. med. Ass.*, **138**, 495.
 Pallister, R. A., and Hu, C. T. (1949). *Med. J. Malaya*, **3**, 204.
 Sumner, J. (1949). *Tubercle*, **30**, 62.
 Thevathasan, A. W. S. (1949). *British Medical Journal*, **1**, 665.

UNUSUAL MANIFESTATIONS OF MALARIA

BY

A. K. NANDI, B.Sc., M.B.Cal., M.R.C.P.Ed.

Additional Physician, Medical College Hospitals, Department of Medicine, Calcutta

Malaria is an arch-simulator and a great mimic of all diseases. The multiplicity and duplicity of the disease cannot be overemphasized. In spite of the classical division of pernicious malaria into types—cerebral, a'gid, gastro-intestinal, cardiac, renal, purpuric, etc.—a practitioner is sometimes faced with very peculiar symptom-complexes, and becomes bewildered regarding the probable diagnosis. During my practice in malarious districts for more than sixteen years I came across many such puzzling cases, some of which are worth recording.

Case Reports

Case 1.—In 1939, at Jalpaiguri, North Bengal, a healthy robust boy aged about 12 suddenly became unconscious and fell down while taking his meal before going to school. I saw him within an hour of the attack. He was then having epileptiform fits. His temperature and blood pressure were normal, the liver and spleen were not palpable, and no abnormality was detected in the heart and lungs. The pupils were moderately dilated and reacted sluggishly to light. The question was whether it was a case of epilepsy—although he had never had such an attack before—or of cerebral malaria, Jalpaiguri being a highly malarious district. Blood was immediately taken and a few M.T. rings were found in the blood slides. Quinine was given parenterally and the boy recovered after being unconscious for about 48 hours. The interesting feature in this case is that no rise of temperature occurred at the time of examination, nor was there a history of malaria within six months of the attack. The temperature rose, however, to 100° F. (37.8° C.) about four to five hours after the attack.

Case 2.—In 1941, also at Jalpaiguri, a postal clerk aged about 40 complained of a severe burning sensation and pain in both hands, including the fingers, for about one week. The pain was so excruciating that he had sleepless nights. On examination there was an erythematous flush of both palms and the fingers were slightly swollen, the tips being extremely tender. His temperature varied from 100 to 103° F. (37.8 to 39.4° C.). There was a history of recurrent attacks of malaria. He consulted local doctors, who prescribed sedatives, sulphonamides, and hot boric compresses, but without the slightest relief. When I saw him he appeared to have acute lymphangitis, and thinking it might be of filarial origin I sent the blood for examination for eosinophilia and microfilaria. I was extremely surprised to hear that the blood slides revealed M.T. rings. With hesitancy I gave quinine, and the result was spectacular. The temperature dropped to normal, and the redness and swelling of the palms and fingers, the pain, and the burning sensation all diminished considerably after one injection of 10 gr. (0.65 g.) of quinine bihydrochloride. The patient was fully cured after three injections.

I have never found any mention in the literature that malaria parasites can give rise to lymphangitis or lymphatic obstruction.

It may be that the syndrome was due to obstruction of the capillaries by the parasites.

Case 3.—In 1945 at Burdwan, West Bengal, a boy aged about 14 had an attack of fever on returning from school; this was accompanied by a chill but no definite rigor. The next morning he was seen by a local doctor, who gave him quinine. That afternoon, however, his temperature shot up to 104° F. (40° C.). He had great difficulty in swallowing food, his voice was nasal, and liquid food regurgitated through the nose. The doctor thought it was a case of diphtheria, and I was called in. On examination there was no patch in the throat; the boy was fully conscious, and had a high temperature, slight enlargement of the spleen, and paralysis of the soft palate; no other abnormality was detected. His blood was examined and M.T. rings were found. The boy made an uneventful recovery after one week's course of quinine.

Case 4.—In 1947, at Dacca, Eastern Pakistan, a man of about 30 returned home with severe headache after a morning walk. A little later he became unconscious and had severe convulsions—twitchings of face and arms were particularly severe. His pulse was very slow, varying from 30 to 35 a minute, and was irregular. The blood pressure was 120/78 mm. Hg. His pupils were slightly contracted and equal, and feebly reacted to light. Jerks were rather brisk, Babinski's reflex was flexor, and abdominal reflexes were sluggish. The axillary temperature was 98° F. (36.7° C.). The liver and spleen were not enlarged. The lungs were normal and there was partial heart-block. Adams-Stokes syndrome was diagnosed by a local doctor, who called me in for an opinion. On inquiry it was learnt that the patient had had two or three attacks of malaria about six months before. A specimen of blood was at once sent for examination, and the report came back that the slides contained many M.T. rings. Quinine was first administered intravenously with glucose, then intramuscularly, and the patient recovered after being unconscious for seven days. During this time he was fed by intragastric drip, and atropine, adrenaline, and nikethamide were given. The interesting feature of this case was heart-block giving rise to an unduly slow irregular pulse. There was a rather small rise of temperature throughout the whole course; on the first day it rose to 99° F. (37.2° C.) towards the evening, and on subsequent days it varied from 97 to 101° F. (36.1 to 38.3° C.).

Case 5.—A man aged about 24 was brought to the emergency room of the Medical College Hospitals, Calcutta, in January, 1949, with a history of severe pain in the abdomen and vomiting for about 48 hours. He was very dehydrated, the tongue was coated and dry, the pulse rapid and feeble, and the abdomen greatly distended, with marked tenderness and rigidity over the epigastrium. There was bilious vomiting and absolute constipation for two days. His temperature was only 99° F. (37.2° C.). The spleen and liver could not be palpated owing to pronounced distension of the abdomen. He was admitted to a surgical ward as a case of "acute abdomen" requiring immediate operation. While the house-surgeon was giving pre-operative treatment the patient had a severe rigor and the temperature shot up to 105° F. (40.6° C.). The blood was immediately examined and Q.T. parasites (band forms) were found. He was at once transferred to my ward and given 10 gr. (0.65 g.) of quinine bihydrochloride with normal saline and glucose intravenously. Quinine was given intravenously at intervals of six hours for two more doses and the patient made a dramatic recovery. The occurrence of a rigor immediately before the operation was providential; otherwise the result might have been disastrous. The presence of Q.T. parasites in such a case is extremely unusual.

Summary

Five cases presenting rare manifestations of malaria are recorded, each of which was confirmed by positive blood findings; in four cases the causative organism was *Plasmodium falciparum*, but in the fifth case it was *Plasmodium malariae*, which is extremely unusual. The peculiar symptom-complex in each case may be explained by the fact that sporulation of the malignant parasite generally takes place in the internal

organs and may affect any part of any organ of the body. Difficulty in diagnosis arises if the possibility of malaria is not thought of, and examination of the blood (if necessary, repeated examinations) is essential for a definite diagnosis.

Early detection and immediate specific treatment will save many lives: one cannot afford to wait long under such critical conditions.

Medical Memoranda

Vasa Praevia Simulating Ante-Partum Haemorrhage

The following case is recorded as a reminder of a rare cause of bleeding in the first stage of labour.

CASE REPORT

A primigravida aged 35 was admitted on April 24, 1949. Her last menstrual period started on July 25, 1948, and the pregnancy had been uneventful until two weeks before admission, when the blood pressure began rising and the urine contained a trace of albumin. On April 21 the membranes had ruptured spontaneously, and on the day of admission pains had started, followed two hours later by a sudden considerable loss of bright blood. From that time no foetal movements had been felt.

Photograph of the placenta, showing the torn umbilical vein. Black rods have been placed in the torn vessel.

On examination the patient's general condition was good. The pulse rate was 76, the blood pressure 130/85, and there was some oedema of the ankles. The uterus was at full term and the vertex presenting, deeply engaged; the foetal heart could not be heard. There was slight continuous vaginal bleeding. A diagnosis of toxic antepartum haemorrhage was made. Labour progressed rapidly, and five hours after admission the

os was fully dilated. After two hours in the second stage the infant was delivered by forceps—a male, 8½ lb. (3.85 kg.), still-born, fresh, and with marked pallor. The placenta and membranes were expelled intact within a few minutes, with very little blood clot. Examination of these showed that the insertion of the cord into the placenta was velamentous and bordered the hole in the membranes. At one point the membranes had torn into the umbilical vein as it lay separated from the arteries (see photograph). The placenta was ischaemic, but its maternal surface showed no sign of premature separation.

The sudden haemorrhage associated with cessation of foetal movements and absence of foetal heart sounds shortly after the onset of pains indicates that, as the cervix began to dilate, the opening in the membranes already present extended outwards and involved the more delicate vessel in the cord. Vasa praevia is an uncommon condition and provides an unexpected foetal risk.

My thanks are due to Mr. A. P. Bentall for permission to publish this case.

E. A. J. ALMENT, D.Obst.R.C.O.G.,
Late Obstetrical-Gynaecological House-Surgeon,
Norfolk and Norwich Hospital.

Experimental Ascorbic Acid Deficiency

The article recently published in the *Journal* on the Falkland Islands Dependencies Survey (Roberts, 1949) prompts me to report an experiment in ascorbic acid deprivation carried out on a similar expedition to Hope Bay, Graham Land, in January, 1945. It was desired to observe how long an interval was necessary for the clinical manifestations of ascorbic acid deficiency to develop in an individual living on the normal diet at the base, and on the unsupplemented sledge ration. As the observations were made on myself alone, they can be regarded as indicating only the approximate period required.

During the journey to the Falkland Islands I had been taking the normal diet, which included large quantities of fresh oranges, orange juice, and fresh vegetables. It is therefore reasonable to assume that on arrival at Hope Bay a condition of near-saturation had been achieved. Thereafter I ate the normal unsupplemented diet, which included fresh penguin meat and fresh seal meat, when available, but I omitted the 50 mg. of ascorbic acid which was taken daily by all the other personnel.

During the following two months no abnormal symptoms or signs were observed. At the beginning of the third month, however, an increasing degree of hyperkeratosis was evident. This particularly affected the extensor and flexor surfaces of the thighs and the extensor surfaces of the arms and forearms, but was noticeable on the abdomen, chest, and back also. Some desquamation was apparent, scales of about 1 mm. square being lost. No petechiae were seen.

In the beginning of the fourth month I experienced lassitude and increasing fatigue, so that travelling became an effort. On May 7 I had a short attack of giddiness: at the same time irritability to an extent unwarranted by the provocative incidents was noted. There was no abnormality of the alimentary or urinary systems. Two ulcers developed in the mouth, one on the gums and the other on the hard palate, but no bleeding occurred. On April 16 and May 11 the tourniquet test for capillary resistance was performed. No capillary haemorrhages were observed on either occasion.

After May 20, I took 400 mg. of ascorbic acid daily for one week, then 100 mg. daily for a further week, after which time I consumed the normal dose of 50 mg. daily. During the following month the hyperkeratosis, ulceration of the mouth, and other signs gradually disappeared, and after two months no trace of the condition remained. One interesting feature was the appearance on June 5 of oedema of both ankles, for which no other cause was apparent. By the following month the oedema had disappeared, and has not recurred since.