HOSPITAL OUTBREAK OF ENTERITIS DUE TO DUCK EGGS

BY

LAWRENCE P. GARROD, M.D., F.R.C.P.

*Bacteriologist and Sanitary Officer, St. Bartholomew's Hospital ;
Professor of Bacteriology, University of London*

AND

M. B. McILROY, M.B., M.R.C.P.

*Chief Assistant. Medical Professorial Unit, St. Bartholomew's
Hospital*

On the morning of July 7, 1949, about 40 nurses at St. Bartholomew's Hospital had reported sick with an acute febrile enteritis, and many patients throughout the wards of the hospital were similarly affected. Inquiry in the kitchen revealed that a lightly cooked pudding containing 200 duck eggs had been served to the nursing and other female staff and to patients on July 5. It was therefore no surprise when stool cultures revealed the presence of *Salmonella typhi-murium*. The outbreak was on an exceptional scale for one so produced, and had features which appear worthy of record.

Source of Infection

The incriminated dish was a "queen of puddings." In preparing this, yolks of eggs are beaten up with milk and added to bread crumbs in dishes in which the mixture is baked for 35 minutes, during which it becomes semi-solid. It is then smeared with hot jam and covered with a layer of beaten-up white of egg mixed with a boiling saturated solution of sugar. The pudding is then returned to the oven for five minutes to brown this top layer. The whole of a consignment of 200 duck eggs had been used to make this pudding: none of these were left, nor was any of the pudding itself, when its effects became manifest two days later. That it was the cause was proved conclusively by its effects on the family of a member of the kitchen staff who took some of the pudding home and gave it to them: these persons were admitted to another hospital, and *Salm. typhi-murium* was recovered from their stools.

It was found that the duck eggs had been collected by the wholesaler from 16 farms in Essex, at two of which conditions were reported by the local health authorities to be insanitary.

This pudding was served in the nurses' dining-room, where certain other female staff also take their midday meal, and to patients throughout all the wards. The amount prepared was said to suffice for 700 portions, but it is impossible to state how many people ate it, because an alternative sweet was also served.

Clinical Features

Varying numbers of patients in almost every ward of the hospital were affected, totalling about 60. A few, owing in part to their previous condition, were gravely ill, and one elderly woman died. That comparatively few patients were attacked is due to the preference of the average hospital patient for rice pudding, the alternative sweet served. It was impracticable to follow all these cases, and this account will deal in detail only with the disease as seen in the nursing and domestic staff, of whom one of us was temporarily in charge.

Attack Rate.—At the time of the outbreak the nursing and domestic staff numbered 351; of these, 197 ate the pudding and 136 (69%) were affected; 154 did not eat the

pudding and 11 (7%) were affected. Of those who ate the pudding 59 out of 136 were sufficiently ill to report sick.

Incubation Period.—The time of onset of the first symptom was determined to the nearest hour by direct inquiry or by a questionnaire distributed within a week of the start of the epidemic. Opinions differ on the limits of the incubation period of *Salm. typhi-murium* enteritis. The American Public Health Association (1945) quotes 6–48 hours, but Dack (1943) allows up to 72 hours. Savage and Bruce White (1925), in a report of 100 outbreaks of food-poisoning, give 48 hours as the longest incubation period of this infection. In the present epidemic there were a number of secondary cases, and it is difficult to define the upper limits of the incubation period, but of 136 persons who ate the pudding and were ill 120 (88%) noticed symptoms within 48 hours. The mean incubation period was 23.8 hours in these cases. Of the remaining 16, three had an incubation period of 95–96 hours: they are considered secondary infections. There is not enough evidence to be sure whether the remaining 13, with incubation periods of 48–76 hours, were primary or secondary infections (see Chart).

Chart showing time between eating of pudding and onset of symptoms.

Secondary Infections.—The 11 who developed symptoms without having eaten the pudding require further explanation. If the pudding was the source of infection, then either their evidence was incorrect or they were infected secondarily. All 11 had typical symptoms and four had positive stools. Nine had incubation periods of 42 hours or more and could be secondary cases. The remaining two, who were ill 23 and 27 hours after the pudding was served, ate lunch in the main dining-room on the day in question, but said they did not eat the pudding. One had a positive stool culture: the stools of the other were not examined until after recovery, when they were negative. It is just possible that these two were served the alternative sweet with a spoon used for the infected pudding. The total number of secondary infections among nurses and domestic staff is thus thought to be 12, including five with positive stools. There was ample opportunity for cross-infection, since many nurses were looking after affected patients in the wards. Secondary infections also occurred among the patients. Four patients in different wards developed *Salm. typhi-murium* enteritis 4, 6, 7, and 21 days after the start of the epidemic. All four had positive stools. The latest secondary case was that of a woman who was ill two months after the epidemic. She was thought to have contracted the infection from a baby born during the epidemic which was infected by its mother and continued to excrete bacilli for several weeks. Another secondary attack was produced in an unusual way: it occurred in a physicist who was performing estimations of the radio-iodine content of the excreta of a patient with thyrotoxicosis who unfortunately contracted the infection.

Symptoms.—The presenting symptoms in the 136 persons who ate the pudding and were ill were as follows :

Abdominal pain ..	37 patients	Backache ..	9 patients
Headache ..	27 "	Tiredness ..	3 "
Diarrhoea ..	22 "	Dizziness ..	2 "
Nausea and vomiting ..	12 "	Faintness ..	1 patient
Shivering ..	11 "	Stiff neck ..	1 "
Malaise ..	10 "	Fever ..	1 "

It will be seen that abdominal pain, headache, and diarrhoea were the commonest. The disease was more closely studied in the 59 who reported sick: in 48 of these stool cultures were positive. Fever occurred at some stage of the disease in 55 (93%), and 48 (81%) had a temperature of over 100° F. (37.8° C.). The average duration of fever was 3.1 days, and the longest 7 days. Diarrhoea with three or more stools a day was present in 45 (76%), and the largest number of stools per day was 26. The stools were watery, with mucus but no blood. No green stools were reported. Vomiting occurred at some time in 18 (31%), usually at the onset. Abdominal pain was almost invariably present, and was colicky. Three nurses developed transitory urticarial rashes while they were ill, and one ward maid had an epistaxis. The average duration of illness in the 59 cases reporting sick was 6.1 days; the longest was 12 days. In 10 cases (17%) there were sequelae—in three, mild diarrhoea persisted for two to three weeks; five had a further mild gastro-intestinal upset lasting two to three days in the next four weeks; and two of the more severely ill complained of weakness and lassitude during the next month. Stools examined during these possible relapses were all negative.

Treatment.—All cases reporting sick were given sulphaguanidine or sulphasuxidine, 3 g. at once and 1.5 g. six-hourly, without obvious effect. Three of the more severely ill nurses were given streptomycin by mouth in 0.5-g. doses four-hourly: this treatment was also given to several very ill patients, and was considered to have had a good effect. The organism was shown to be streptomycin-sensitive, being inhibited *in vitro* by a concentration of 1 µg. per ml.: in an infant similarly treated the organisms still present in the stools after recovery were found to be highly resistant. If streptomycin is used in this disease it should evidently be given by mouth. It is of interest in this connexion that two patients under treatment with intramuscular streptomycin for tuberculosis developed enteritis in spite of it, although in one of them the attack was mild and somewhat atypical.

Laboratory Investigations

Examination of Faeces.—Cultures of specimens from some of the original cases and certain others in the acute stage were made on desoxycholate-citrate medium and MacConkey's medium. No attempt was made, or could in the circumstances be made, to confirm the diagnosis in all the cases at the height of the epidemic, which accounts for the fact that positive stool cultures were not obtained in all evident cases discussed in the foregoing analysis. On the other hand, it was imperative to examine faeces from all affected nurses after recovery to be sure that none remained a carrier. Many had to return to duty before this had been done, and some had gone on more or less prolonged sick leave or annual leave; but arrangements were made to obtain a series of specimens at intervals of not less than four days from all affected nurses, until at least two negatives had been obtained. (Kitchen and other domestic staff and, so far as possible, affected patients were examined similarly. These examinations continued for several weeks and raised the total of specimens dealt with in connexion with the outbreak well into four figures.)

Such specimens from convalescents were cultivated solely by the heavy inoculation of a tube of tetrathionate medium, which was plated after overnight incubation.

These specimens were obtained at such irregular intervals that no adequate data exist for a detailed analysis of the rate of disappearance of the organism. The findings, such as they are, may be summarized by saying that almost all specimens examined during the first week were positive: the number of known positives fell by the end of the second week to 12 (8%) and after a third week to 4 (3%); thereafter only one positive culture was obtained (on the 22nd day). (These figures refer to nursing and domestic staff only: several patients, including two infants, carried the organism for longer periods.)

Agglutinin Formation.—The blood of 12 nurses was examined for agglutinin during convalescence, in most cases about two weeks after the onset. Agglutinin for *Salm. typhi-murium* was found in all but one, the titre varying from 1 in 25 to 1 in 500: there was a strong suggestion that a higher titre was reached after a more severe attack.

Cooking Experiment.—With the kind co-operation of the catering officer, a pudding identical to that responsible for the outbreak was inoculated with culture of *Salm. typhi-murium* and cooked in the same way, cultures being made from it afterwards. The object was to produce in both egg-containing parts of the pudding a bacterial content of about 100,000 organisms per ml., and viable counts before cooking showed that this had been achieved in the lower layer (yolk of egg and milk). It was not thought advisable to add culture to the white of eggs before beating and the addition of boiling sugar solution, owing to the danger of contaminating the power-driven machine used in the preparation of this material, and after its preparation it was difficult to achieve uniform admixture of a small amount of added liquid. Moreover, this is a material with which it is almost impossible to work quantitatively. This part of the experiment was therefore not wholly satisfactory.

The results can be expressed shortly by saying that whereas before cooking both constituents contained the organism, in the former case in very large and ascertained numbers, afterwards both were almost completely sterile, and from neither constituent in any of many cultures was *Salm. typhi-murium* recovered.

Discussion

It has been recognized for nearly twenty years that duck eggs, unlike hens' eggs, may actually contain pathogenic organisms of the *Salmonella* group, commonly *Salm. typhi-murium* and sometimes *Salm. enteritidis*, owing to the existence of this infection in the bird itself. Hens are susceptible to infection with certain other *Salmonellae* also pathogenic for man, but in their eggs contamination is confined to the outer surface of the shell, and has been an important cause of food-poisoning only as the result of the large-scale use of imported dried egg (Report of M.R.C., 1947). The risk of eating duck eggs unless thoroughly cooked is fairly well known in medical circles, but perhaps less so among caterers, and it seems that a useful purpose may be served by giving this episode wide publicity. It is believed to be unique in the scale on which the eggs were used and the number of persons affected: most cases of enteritis so produced are single, due to the consumption of one lightly boiled or poached egg, and it is likely that the origin of many remains unrecognized.

Experiment failed to confirm that the organisms can survive the cooking process employed. In view of their

low thermal death point, the survival of *Salmonellae* in even lightly cooked dishes is difficult to explain. Recent experiments (Report of M.R.C., 1947) with reconstituted dried egg known to be infected showed that the great majority of the organisms were killed by scrambling (producing a temperature of 83° C. for three-quarters of a minute) and in omelette-making (85° C. for one minute). Only when the reconstituted egg was first incubated overnight, so producing a bacterial content of between 10,000 and 1,000,000 per g., were positive cultures obtained after cooking, and then only from an inoculum of no less than 50 g. In view of these findings, and of failure to recover organisms from an experimental pudding of the kind concerned here, the survival of sufficient numbers of bacteria in the incriminated food to cause so high an attack rate in those consuming it remains a mystery.

This outbreak is also of interest on account of the hazardous measures without which the work of the hospital could not have been continued. The only persons forbidden to return to work until after repeated negative stool examinations were food-handlers, a kitchen employee, and two dining-room maids. It was necessary that nurses should return to duty as soon as they were fit to do so, and proof of their freedom from infection was not obtained until some time later: for a few days at least after their return to duty many were probably carriers, and some were shown to be. They were of course fully instructed in the precautions they must take, and evidently observed them faithfully. A greater problem at the height of the epidemic was the nurse with a mild or unrecognized attack of the disease who remained on duty. It is possible that some transmission to previously uninfected patients took place during this brief and strenuous period.

Summary

An account is given of an outbreak of *Salm. typhi-murium* enteritis involving nursing and other female staff and patients in a large general hospital. The vehicle of infection was a lightly cooked pudding containing 200 duck eggs. The clinical features of the disease are described, together with laboratory investigations and the steps taken to control the epidemic.

We are indebted to Dr. Joan Taylor, of the Salmonella Reference Laboratory, for kindly examining five of the strains isolated from patients and confirming their identity.

REFERENCES

- American Public Health Association (1945). *The Control of Communicable Diseases*. New York.
Dack, G. M. (1943). *Food Poisoning*. Chicago.
Report (1947). *Spec. Rep. Ser. med. Res. Coun., Lond.*, No. 260.
Savage, W. G., and White, P. Bruce (1925). *Spec. Rep. Ser. med. Res. Coun., Lond.*, No. 92.

A subcommittee of the Nurses and Midwives Whitley Council has made a report on working conditions in mental hospitals and recommended measures for the achievement of student nurse status. It points out that more qualified staff must be provided if the student is to do only the work that is necessary for gaining practical experience in training. The buildings should be reorganized to provide better facilities for the patients and less needless work for these students. Central teaching units should be established to which students could go for a period of special tuition each year. Training should be supervised by education officers (who should be qualified tutors) in co-operation with the matron and chief male nurse. The period at the preliminary training school should be extended as soon as possible to three months, and some of this time should be spent in wards. Lectures and demonstrations should be given in duty time, and no student should be on night duty for a month before the examination. Students should not be put in charge of wards and should not have to work more than 48 hours a week. Their uniform should be distinctive.

THE MARTIUS BULBO-CAVERNOSUS INTERPOSITION OPERATION

BY

WILFRED SHAW, M.A., M.D.Cantab., F.R.C.S.
F.R.C.O.G.

Surgeon in Charge, Gynaecological and Obstetrical Department, St. Bartholomew's Hospital; Gynaecologist, St. Andrew's Hospital, Dollis Hill

This operation has been devised and advocated by Professor H. Martius, of Göttingen, for the treatment of stress incontinence and of particular types of vesico-vaginal fistula. The principle of the operation is to interpose a graft of tissue obtained from the labium majus between the urethra and bladder above and the anterior vaginal wall below. A vertical incision is made into the labium majus, and the fatty tissues, together with the bulbo-cavernosus muscle, are mobilized to form a cylindrical graft. The upper end of the graft is then cut through in the vicinity of the mons veneris. The posterior attachment of the graft is left intact so that the bulk of the blood supply to its tissue is left undisturbed. A flap of anterior vaginal wall is mobilized laterally until the deep tissues of the labium are exposed. The free end of the graft is then drawn through so that the graft can be interposed and stitched into position. The operation is extremely simple to perform, and the only difficulty seems to be to obtain complete haemostasis.

For the treatment of stress incontinence the thick tissues of the graft act as a supporting shelf by pushing up the urethra and the neck of the bladder. In the treatment of vesico-vaginal fistula the bladder is repaired in the usual way and the graft is utilized partly to prevent leakage of urine and partly to separate the sutured bladder from the anterior vaginal wall. I have employed the operation in a small series of cases, the details of which are here given.

Case 1

A primipara aged 53 had suffered from stress incontinence for a year. She had been married for 20 years and her child was aged 19. The menopause had occurred four years previously. The pelvic organs were normal. There was a slight descent of the lower part of the anterior vaginal wall. The meatus was patulous and urine leaked away when the patient coughed.

A Martius operation was performed in July, 1948. A midline incision was made in the anterior vaginal wall, together with lateral incisions from the end of the vertical incision near the meatus. Lateral vaginal

FIG. 1.—Martius operation. An incision has been made in the right labium majus and a graft of tissue containing fat and the bulbo-cavernosus muscle has been mobilized.