X-ray examination following injection of "lipiodol" into the cisterna magna and into the lumbar region of the subarachnoid space with inversion of the patient revealed an extramedullary block extending from the body of the second thoracic vertebra to the lower border of the fourth. Laminectomy was contemplated, but by Nov. 10 slight objective sense perception was demonstrated and some control over micturition was evident. Steady improvement occurred, and normal function and normal signs in both lower limbs were established by May 15, 1933.

On July 12, 1933, an abscess made itself evident at the vertebral border of the right scapula. This was incised, and the subsequent sinus healed in six weeks. A pure culture of *Staph. albus* was grown from the abscess pus. There were no signs or symptoms referable to the back prior to the development of the abscess. Examination as an out-patient in May, 1934, showed the femoral lesions to be completely healed and epithelialized, and the spine and back seemed perfectly normal.

It remains a matter of speculation whether the above sequence of events was due to a metastatic abscess unrelated to a focus of bone infection, or whether the abscess developed secondarily to a "minimal" bone lesion too small to be detected by x-ray examination, as occurred in Case 1 of Mr. Martin's series (Nov. 9, 1946, p. 688).—I am, etc.,

Cleveland, Ohio, U.S.A.

T. FRANCIS JARMAN.

Air Crashes and Fractured Spines

SIR,—The development of civilian flying seems to be causing an increase in the number of fractured spines, and their treatment may now have to be undertaken by those inexperienced in the application of hyperextension plasters. Serious sores may be easily caused, and yet they can almost always be avoided if the two following precautions are taken:

(1) A strip of adhesive felt 6 in. (15 cm.) wide and about 36 in. (0.9 m.) long is fixed down the centre of the back over the spinous processes and smaller strips of the same material cover the iliac crests, symphysis, and manubrium. Stockinette or an old bathing costume may be put over this, but no cotton-wool should ever be used.

(2) The first plaster of Paris applied to the back of the hyper-extended patient must be a 6-in. longitudinal slab from the base of the neck to the sacrum. This is held in place by circular 6-in. plaster bandages, supplemented of course by additional slabs and in particular by the all-important anterior strut, which takes its bearing on the manubrium sterni and symphysis pubis. The common mistake is to start with circular bandages round the lumbar region (which is of course in marked lordosis), with the result that they tend to slip down into the lowest part of the hollow of the back and make a sharp transverse ridge which can and does cause disastrous plaster sores.

Every orthopaedic and fracture department has of course the necessary equipment, but it is my firm belief that even the smaller hospitals should always have a stock of 6-in. plaster bandages and adhesive felt.—I am, etc.,

London, W.1.

ERIC I. LLOYD.

Eye Surgery

SIR,—It is not my desire to counter any criticism in your review of my book *Eye Surgery* (Dec. 28, 1946, p. 989), but I feel unable to allow your reviewer's statement about Krönlein's operation to pass unchallenged, for if this is accepted by your readers it will give a wrong impression of a valuable operation. The comment in your review is thus: "Few surgeons who have tried it will agree that a Krönlein's operation is particularly valuable for the removal of retro-ocular new formations." All the cases which I have done of Krönlein's operation for an orbital neoplasm have recovered visual acuity, in some instances from perception of light to 6/6 (partly), and the visual field has improved very appreciably. In two of these cases orbital exenteration had been advised by more than one surgeon. In one case transfrontal craniotomy had been done at Oxford, where the report was that "the orbital neoplasm was not seen." Nine months later I removed it by Krönlein's operation; the patient was spared exenteration; the optic nerve which was stretched over the neoplasm recovered almost entirely, save for a small defect in the visual field.

Another point made in the review concerns the use of two needles for dividing thick capsule remnants. The reviewer writes: "Whether one needle bent at an angle should be used

in operations with two needles seems a moot point. Most surgeons prefer to avoid angled instruments, since direction is less certain than with straight instruments." He will have read earlier in the book my comments about straight instruments making precision work easier than is the case with angled instruments. The latter are, however, necessary in entering the eye from the nasal side, as in capsulotomy with two needles and in using a keratome on a deep set eye with prominent supra-orbital margin. It is quite impossible to use anything but an angled instrument under these circumstances, the nose and brow forming obstacles.

Another comment which I think is likely to mislead is: "The author likewise probably takes an extreme view in believing that malignant melanomata of the choroid are not so radiosensitive as glioma of the retina." Although I have had success in irradiating malignant melanomata of the choroid with radon seeds sutured to the sclera, it is a fact that these neoplasms are not so radiosensitive as gliomata retinae.

I express my grateful thanks and appreciation of your reviewer's generous remarks elsewhere in his review.—I am, etc.,

London, W.1.

H. B. STALLARD.

Defective Colour Vision in Industry

SIR,—In the opening sentence of the leading article (Dec. 21, 1946, p. 948) you make an excellent point when you say "Defective colour vision, more commonly and less accurately called colour-blindness . . ." That you should emphasize the frequency of terminological inexactitude in this connexion is of great importance at the present time when the role of colour vision in industry is receiving greater attention. So many of the so-called anomalous trichromats declare most emphatically that they are not colour-blind, and their contention is undoubtedly correct. The explanation of the defect as a faulty colour appreciation is well received, and now is the time to urge very strongly that the term colour-blind should be replaced by the term colour-defective.

The time is also opportune to suggest that the presence of defective colour vision is only in a limited number of cases a bar to the efficient performance of many of the tasks mentioned in your article. As is well known, the R.A.F. had many colour-defectives who, as members of aircrew, played an important role in all commands, and the salvage of manpower achieved by the use of the colour-defective safe category was considerable. This "safe" category has, however, been subjected to much criticism—perhaps the most important being that the personal equation of the examiner was too great. During the war much work was done in this country and in America to work out the permissible degree of colour-defect for certain tasks. Clearly there is a great field for the salvage of colour-defectives in industry just as in the Services, but it is now realized by all who have been asked to do colour vision testing on a large scale that the problem is exceedingly complex and admits of no easy solution. Difficult borderline cases will always appear.

Later in the article you are on less sure ground when you state that the mothers of male colour-defectives are not themselves affected. It is probably true to say that these relatives are not so grossly affected, but by more accurate testing methods—anomalouscope or colorimeter—quite a number of these women show a definite deviation from normal. Pickford² has investigated this problem, and he found among the female relatives of colour-defectives a considerably higher incidence of colour defects than the expected. I have seen one or two families where the sister of colour-defective brothers has had a marked degree of colour defect.—I am, etc.,

Dundee.

JOHN GRIEVE.

REFERENCES

- ¹ *Nature*, 1946, 157, 376.
- ² *Ibid.*, 1944, 153, 409.

Diet and Canine Hysteria

SIR,—As a member of the sister profession may I be allowed the courtesy of the use of your columns to comment upon the report (Dec. 14, 1946, p. 885) of Sir Edward Mellanby's feeding experiments? This piece of work has aroused lively interest in the veterinary world, and we would all most heartily welcome a final elucidation of the aetiology of canine hysteria.