WATER AND SALT DEPLETION*

BY

H. L. MARRIOTT, C.B.E., M.D., F.R.C.P.

Physician with Charge of Out-patients, Middlesex Hospital

Causation of Pure Salt Depletion

It has already been explained that the chief effect of salt depletion is a state of secondary or extracellular dehydration consequent upon loss of osmotic pressure in the extracellular fluid ; therefore to speak of "pure" salt depletion may seem paradoxical. The expression is meant in a causal sense—that is to say, the state which occurs in patients who become deficient in sodium and chlorine while being liberally supplied with water. This state is common in medical and surgical practice, but is often unrecognized for what it is. The two essentials for its causation are: (1) Abnormal losses of sodium and chlorine. Simple inadequacy of intake does not, as in the case of water, produce serious deficiency. However, low intake or lack of intake may play an important part in the rapid development of depletion when abnormal salt losses are occurring. (2) Adequate water intake. As mentioned earlier, the condition of adequate water intake is frequently fulfilled in these days of "dehydration consciousness," when so many patients are rightly put on fluid balance charts. Water is pressed upon them or isotonic glucose solution is parenterally administered. The intake and output figures may be gratifying till the late stages. Yet the patients become weaker and weaker and finally die. Abnormal losses of sodium and chlorine may occur in alimentary secretions, sweat, or urine (Addison's disease). Table IV gives approximate maximum daily losses

TABLE IV.—Possible Secretion Losses (24 hours)

Secretion	Circumstances of Abnormal Loss	Volume	NaCl Equivalence		
			NaCl	= Isotonic Saline	
				Litres	Pints
Sweat	Sweating	ml.	g.		
Saliva	Vomiting or gastric suction	14,000	35	4	7
Gastric juice ..		1,500	8	1	2
Bile		2,500	18	2	3½
Pancreatic juice ..		500	4	½	1
Intestinal juices ..	Diarrhoea, intestinal suction	700	4	½	1
		3,000	22	2½	4½
Mixed secretions:					
Vomit	Vomiting or suction	6,000	40	4½	8
Diarrhoea		7,000	50	6	10½

of sweat and alimentary secretions. It is obvious that smaller losses, prolonged over days or weeks, can also cause severe depletion, particularly if salt intake is lowered. In Great Britain vomiting and diarrhoea are the commonest causes of abnormal losses of sodium and chlorine. The large number of frequently occurring commonplace medical

and surgical conditions causing vomiting and diarrhoea need not here be enumerated. Acute poisoning may cause extremely violent vomiting and diarrhoea, with precipitous decline of extracellular electrolytes, and provides particularly striking examples of the development of extracellular dehydration within a space of minutes or an hour or two (Marriott, 1935). Reference to Fig. 8 shows that vomiting and diarrhoea respectively cause very different proportional losses of chlorine and sodium. Vomiting causes great relative depletion of chlorine ions and the well-known tendency to alkalosis. Diarrhoea, on the other hand, has the reverse effect, with disproportionate loss of sodium and tendency to acidosis. Therapeutic continuous aspiration of gastric or intestinal contents (Wangensteen, 1942) can cause great withdrawal of these ions. Lyall and Nicol (1939a and b) found that continuous gastric aspiration resulted in removal of 93 g. of chloride, expressed as NaCl, in six days. Such an amount represents more than half the body's total. Bartlett *et al.* (1938) produced severe salt depletion in normal subjects. Gastric or intestinal suction drainage is a valuable method of treatment, but if used without parallel administration of saline can be highly dangerous.

FIG. 8.—Showing the relative ratio of loss of sodium and chlorine in vomiting and diarrhoea.

The effects of pure salt depletion are perhaps best seen in severe sweating, accompanied by free drinking of unsalted liquid, because the relative proportions, though not the concentration, of sodium and chlorine in sweat do not deviate very much from the proportions in extracellular fluid and because there is little loss of other substances. Also, climatic and industrial conditions may permit mass observation of large numbers of normal individuals, suffering only from the effects of salt loss from sweating, without obscuring effects due to some underlying pathological condition. Such little understanding as I have of the effects of pure salt depletion comes mainly from observation, during four years in the Tropics, of men sweating severely while drinking water freely (Marriott, 1943, 1945a) supplemented by reference to the experience of others—notably Marsh (1937), whom I visited at Abadan, in the Persian Gulf, in 1942 ; Moss (1923), Dill (1938), Caplan (1942), Taylor *et al.* (1943), Ladell, Waterlow, and Hudson (1944) ; and last, but not least, McCance's classical experimental studies (1936). It was my duty in India to do special tours in the hottest weather (June) to observe heat effects in such particularly

* Part II of the Croonian Lectures delivered at the Royal College of Physicians, London, on Dec. 3 and 5, 1946. The first part appeared in the issue of Feb. 15 at page 245.

hot stations as Allahabad, Cawnpore, Lucknow, and Bareilly. The tour of 1942 was particularly instructive, because it happened to be an unusually hot season (maximum shade temperatures in the above stations were between 115° and 123° F. (46° and 50.6° C.) and sun temperatures very much higher) and because in that year—the first real war year for India—there was not adequate shade provision for men, nor was there general realization of the importance of extra salt intake. During this hot season there were 1,959 admissions to hospital for heat effects and 136 deaths. I personally saw some 400 cases. During my period in the Tropics I also saw salt depletion complicating the picture in many illnesses—for example, scrub typhus (Marriott, 1945b).

Much will be said below about the clinical and pathological features of pure salt depletion as seen, in uncomplicated form, in individuals whose salt loss arises from sweating due to heat. I wish to emphasize that the same clinical and pathological picture is frequently to be seen in the medical and surgical wards of our own hospitals in cases of vomiting, gastric or intestinal suction drainage, biliary or intestinal fistulae, diarrhoea (e.g., ulcerative colitis), and Addison's disease.

The main mechanisms by which the body may be most seriously injured by hot atmospheric conditions are: (1) loss of water and salt from sweating; (2) overheating from cessation of sweating. The first is called "heat exhaustion"—an unfortunate term, because it implies mere fatigue whereas it is a serious and may be a fatal state. In 1942 the incidence of this condition in India was 1,405 cases with 27 deaths in British troops and 255 cases with 13 deaths in Indian troops. There were many more Indian troops in the area at the time, and so the incidence in British troops was perhaps thirty to fifty times greater. The probable explanation would seem to be a lower concentration of NaCl in the sweat of Indians. Overheating from cessation of sweating is called "heat hyperpyrexia" or "heat-stroke," and its essential characteristic is a temperature of 107° F. (41.7° C.) or above. Its underlying cause seems most commonly to be some infection, often minor, which sets off the pyrexia reaction with its associated inhibition of heat-loss mechanism. In India in 1942 its incidence was 170 cases with 34 deaths in British troops and 129 cases with 62 deaths in Indian troops.

Heat hyperpyrexia is not relevant to the present discussion and will not be further considered. Heat exhaustion—heat exhaustion Type 1 in Ladell, Waterlow, and Hudson's classification (1944)—however, when occurring in men drinking water freely, presents a typical picture of pure salt depletion. Sweating without water drinking produces effects mainly of water loss, because sweat is much more dilute than extracellular fluid. Hence in men lost in deserts, and without water, the effects of water depletion overshadow the effects of salt depletion.

Effects of Pure Salt Depletion

Serious depletion of sodium and chlorine gives rise to the following conditions:

(A) An *invariable* loss of total osmotic pressure of the extracellular fluid. In consequence the kidneys excrete water and there is diminution of extracellular fluid volume—extracellular or secondary dehydration.

(B) A *possible* disturbance of acid-base equilibrium if loss of sodium and chlorine ions is not proportionate (see Fig. 8).

Both changes produce clinical effects, but of the two I would submit that the former is the more important. In severe vomiting and severe diarrhoea the main serious effects are the same in both conditions and are due to secondary dehydration. The manifestations due to alkalosis

and acidosis respectively are of secondary importance. It has been unfortunate that it has so happened that the historical progress of medical discovery caused attention first to become focused on acid-base disturbances. This accident of history has resulted in undue preoccupation with acid-base changes, in cases of the type described, to the relative exclusion of the more serious effects due to loss of extracellular osmotic pressure. In any event, if sodium chloride is administered, along with enough water, the kidneys will automatically restore the acid-base balance by appropriate differential excretion of sodium and chlorine ions. These remarks are not, of course, intended to apply to disturbances of acid-base equilibrium arising from causes other than asymmetrical loss of sodium and chlorine ions in alimentary secretions—that is, not to acid-base alterations from the excessive administration of acid or alkaline salts, overbreathing, or the accumulation of ketone bodies.

A. Effects due to Loss of Osmotic Pressure

Lassitude; Apathy; Stupor.—Anyone who has lived in the Tropics knows the feeling of extreme lassitude in the hot weather and that it is alleviated by ingestion of extra salt. Caplan (1942) found fatigue and asthenia in 243 of 244 cases of heat exhaustion. McCance (1936) states: "We have all suffered from excessive fatigue and a general sense of exhaustion—Whitteridge and Niven both got into an extraordinarily interesting state, in which they were content to sit and do nothing in a chair for hours on end." Lassitude and apathy, progressing to stupor, were marked features in the cases of heat exhaustion which I observed. In the hot seasons I was often struck by the apathy of men in hospital for sickness of various kinds, especially those eating poorly; examination of their urine for chloride generally demonstrated its absence. Black (1946) has recently published a paper showing how an increased sense of well-being and a return of vigour were induced in cases of chronic sprue in India by the administration of extra salt. Such men were not only losing salt by the skin but also in their copious evacuations. Lassitude is an earlier and more pronounced feature in pure salt depletion than in simple water deprivation.

Weakness.—Muscular feebleness, as distinct from the mere disinclination for effort arising from the lassitude, is definite. McCance (1936) describes how he found his arm getting tired when shaving and noticed his jaws tiring when eating. The myasthenia of Addison's disease is well known.

Headache.—This is often present. It has the character of a low-pressure headache; it is much less noticeable if the subject gets out of the vertical position and lies down.

Giddiness and Tendency to Orthostatic Fainting.—This is frequent. I saw it in many cases of heat exhaustion. Caplan (1942) noted it in 215 of 244 cases. Ladell *et al.* (1944) say: "The effect of standing was dramatic. The pulse became progressively more difficult to feel and more rapid. The blood pressure fell and often could not be taken because the sounds were inaudible. The patient became increasingly pale, began to sweat profusely, to yawn and to complain of dizziness. He was obviously on the verge of syncope."

Normal Urine Volume (till Late Stages).—If the subject is drinking freely, urine volume is normal, or may be increased until the late stages, when nausea and vomiting occur; there is, however, a very slow excretion of any particular quantity of water ingested. McCance (1936) states: "We have all noticed that our water metabolism was not normal. After drinking a large amount of water no diuresis would develop at the expected time, but then hours later—possibly during the night—the diuresis would

begin." In Addison's disease delayed diuresis has been noted (Rowntree and Snell, 1931). It is the basis of the water-elimination test described by Robinson *et al.* (1941). The delayed diuresis is usually ascribed to delayed excretion. I shall presently advance another hypothesis, which I have not seen elsewhere put forward—namely, that the delay may be chiefly or in part in *absorption* and that the water may remain overlong in the stomach and alimentary tract. Chloride is absent from the urine except in Addison's disease.

Absence of Thirst.—This is most important. It has been laid down as an essential condition in the causation of pure salt depletion that there must be liberal intake, oral or parenteral, of unsalted water. However, even if intake is relatively low but not low enough to make the case one of predominant water depletion, there is no thirst; indeed, water is often repellent. I have seen men suffering from severe secondary dehydration refuse water or spit it out.

Anorexia→Nausea→Vomiting.—These symptoms, in the progression indicated, are present in most moderate or severe cases. Associated with the anorexia is loss or perversion of taste. Caplan (1942) observed nausea in 152, and vomiting in 136, of 244 cases. He regarded 194 of these cases as mild and 50 as moderate or severe; vomiting was invariable in the latter group. Ladell *et al.* (1944) noted vomiting in 73% of 45 cases. My own experience embraces a large number of cases, but unfortunately is impressionistic and unsupported by factual data, as the conditions of my tours precluded spending time on recording and I did not then consider publishing my findings. Indeed, the significance of some of the observations made did not become apparent until there was subsequent time for reflection. Anorexia, nausea, and vomiting are particularly important,

because they inaugurate a vicious circle which causes rapid deterioration (Fig. 9). This train of events is well known in Addison's disease.

Weight Loss.—This is marked, especially in the early stages, in spite of free water intake. It is consequent upon the secondary dehydration. The weight tends to be of the approximate

FIG. 9.—Vicious circle of events in salt depletion.

order of 1 kg. per 3.4 g. of sodium deficit (McCance, 1936).

Changes in Appearance.—Sufferers show signs of subcutaneous desiccation. The eyes are sunken from loss of orbital tissue fluid. In the late stages the appearance resembles the Hippocratic facies. The skin loses its elasticity, so that pinched-up folds do not smooth out on release. The patient no longer fills his skin, and in extreme cases the skin of the fingers may be wrinkled ("washerwoman's fingers" of cholera).

Cramps.—Heat cramps have long been well known. Especially have stokers and firemen realized that they tend to be precipitated by water ingestion and prevented by drinking salted fluids. Caplan (1942) noted cramps in 142 of 244 cases; Ladell *et al.* (1944) in 70% of 45 cases.

Mental Confusion and Delusions.—I saw several men with marked mental changes during and shortly after heat exhaustion; some were just confused and disoriented—a similar picture to that seen in severe vomiting or diarrhoea. Others had delusions. One young soldier was possessed with the idea that he was for execution on the morrow; a nursing sister was stated to be mad. Saphir (1945) has commented on how salt depletion, of a sustained mild degree, may simulate psychoneurosis.

Blood Changes

Plasma Sodium and Chlorine.—When salt depletion is severe there is a fall in the plasma concentration of sodium and chlorine. Most published figures relate to plasma chloride, expressed as sodium chloride, which may fall in extreme cases from the normal levels of 560 to 630 mg. per 100 ml. to 300 mg. per 100 ml. or even slightly lower. As the sodium ion is the more important from the osmotic standpoint, data regarding falls in sodium levels would be more relevant.

Blood Urea.—The blood urea rises (McCance and Widdowson, 1938). Ladell *et al.* (1944) found it raised in all cases of salt-depletion heat exhaustion; 12 out of 26 cases had levels above 100 mg. In spite of high blood urea and low urine volumes in their cases, the urea concentration in urine was low. In 9 cases the ratio between concentrations of urea in urine and plasma was measured and found to average 40 : 1. In severe vomiting and diarrhoea figures of the order of 100 to 200 mg. per 100 ml. or higher are commonplace. In experimental adrenalectomy (Harrop *et al.*, 1933) or in Addison crises the blood urea may be similarly raised.

Plasma Volume.—There is shrinkage of the plasma volume leading to haemoconcentration. Leonard Rogers (1908, 1909a and b, 1911, 1915, 1916, 1921) demonstrated many years ago how in severe cholera cases as much as two-thirds of the plasma volume may be lost. A similar loss may occur as a result of vomiting. In salt depletion due to sweating McCance (1938) produced in volunteers extracellular-fluid reductions of 28 to 38%; Ladell *et al.* (1944) found plasma-volume reductions up to 50% in heat exhaustion. In extreme salt depletion from vomiting, gastric or intestinal suction, fistulae, diarrhoea, sweating, or Addison's disease, a loss of as much as one-half to two-thirds of the plasma volume is possible. It is probable that definite circulatory manifestations (see below) hardly become evident until the plasma volume has been reduced by 15%. A systolic blood pressure below 90 mm. Hg probably indicates at least a 25% reduction. The relative normal ratio of plasma volume to tissue-fluid volume is about 1 : 4, but when the extracellular fluid is reduced through salt depletion the reduction of the tissue-fluid portion is relatively greater than that of the plasma portion (see Fig. 4b) because of increasing colloid osmotic pressure as the plasma protein molecules become concentrated. A reasonable rough assumption is that loss of a litre of plasma volume (normal plasma volume being approximately 3 litres) means a total loss of extracellular fluid of the order of 6 to 7 litres.

Plasma volume may be measured directly, and Evans (1945), in a valuable paper discussing the significance of alterations in blood volume, has rightly stated that blood-volume determinations should be part of the laboratory routine procedure of all hospitals; or, in a non-anaemic subject, the plasma-volume reduction may be deduced from data relating to haemoconcentration. Leonard Rogers was a pioneer in this field. Relevant data are the haemoglobin percentage, the haematocrit reading, and the plasma and whole-blood specific gravities. The last-named, as Rogers realized, is perhaps the most convenient for quick practical use. It may be determined by the falling-drop technique in graduated mixtures of glycerin and water (Rogers) or in graduated copper sulphate dilutions as proposed by Phillips and van Slyke (1943). The correlation between blood gravity and haemoglobin and haematocrit percentages is shown with clarity in their line chart. If the haemoglobin percentage is determined directly, or indirectly, the approximate order of loss of plasma volume can be obtained by

adaptation of a form of calculation proposed by Black (1940b):

$$\text{Plasma loss (in litres)} = 5 - \frac{5 \times \text{Hb}_1}{\text{Hb}_2}$$

where $\text{Hb}_1 = 100\%$, $\text{Hb}_2 =$ value in patient, and 5 litres is the assumed normal blood volume. I would say that, as a very rough rule, in secondary dehydration the plasma loss may be multiplied by 6 to give the total extracellular fluid loss. In extreme salt depletion, with a loss of one-half to two-thirds of the plasma volume, the consequent haemoconcentration may produce such high blood values as the following: specific gravity, 1064 to 1070; haemoglobin, 20 to 23 g. per 100 ml.; haematocrit, 58 to 64%; red cell count, 7 to 8 millions per c.mm. Such findings indicate an extracellular fluid loss in the region of 7 to 10 litres for a 70 kg. man, or about 2 imperial gallons.

Blood Viscosity.—Haemoconcentration increases the viscosity of the blood (Whittaker and Winton, 1933; Holbrook and Watson, 1939; Evans, 1942). Kekwick and I (1940) studied the flow of blood through capillary tubes in devising a capillary regulator, instead of a screw-clip, for obtaining a more constant flow in drip blood transfusions. We then realized the tremendous part that viscosity must play in determining the flow through small blood vessels and capillaries. Poiseuille's equation is said to hold for such flow (Whittaker and Winton, 1933; Seligman *et al.*, 1946).

$$\text{Flow} = \frac{\text{Pressure}}{\text{Viscosity} \times \text{Resistance}} \times \text{Constant.}$$

Hence, if the viscosity should be doubled the flow will be halved. Wynne Willson and I (1940) did some work with various concentrations of red cells in citrated or heparinized plasma in an endeavour to ascertain the correlation between progressive haemoconcentration and increase of viscosity. Our experiments were of a rough-and-ready character, designed to produce a quick answer to urgent resuscitation problems, and were not brought to a stage fit for publication. However, they showed that when the red cell count is around $7\frac{1}{2}$ millions per c.mm. the blood viscosity is about double what it is when the count is 5 millions per c.mm.

The Circulation

If the decrease in plasma volume is at all severe the patient passes into a state of oligaemic circulatory failure or "shock." This is well understood in relation to severe vomiting and diarrhoea, and I have seen it often in heat exhaustion. Caplan (1942), reporting on 50 moderate and severe cases, states: "The patients exhibited all the signs of shock. The skin was cold and clammy—the blood pressure in the severe cases was indeterminate and in the moderate cases the systolic pressure was always below 95 mm. Hg—commonly 60 to 90—and the pulse pressure 2 to 10 mm. Hg." The peripheral failure of adrenalectomized animals or of patients with Addison's disease in crises is well known. The main factors in the production of circulatory failure are: (a) the reduction in blood volume, and (b) the increased blood viscosity. The effect of blood-volume reduction in producing shock (Robertson and Bock, 1919; Keith, 1919) has been much studied in the recent war (Kekwick, Marriott, Maycock, and Whitby, 1941; Dick, 1944; Evans, 1945). Less attention has been devoted to the effect of increased viscosity, which, of course, also occurs in plasma loss in burns and injuries due to crushing. I suggest that in the circulatory failure of severe salt depletion it plays an important part. Seligman, Frank, and Fine have recently published (1946) an experimental study on the haemodynamic effects of raised blood viscosity, and have shown its importance.

The clinical picture of the circulatory failure of salt depletion is the familiar one of increasing oligaemic

vasoconstrictive shock. The patient is pallid and often in a clammy sweat. The extremities are cold (if atmospheric conditions allow cooling). The blood pressure is at first maintained and the diastolic pressure may be raised (Ladell, Waterlow, and Hudson, 1944) but later falls, and the pulse pressure is diminished. The veins are empty. The pulse deteriorates, first in force and then in tension, and usually quickens. Cyanosis of lips and nails appears as the condition develops. When the systolic blood pressure drops to about 70–80 mm. Hg anuria occurs. The mental condition is not as alert and clear as in haemorrhagic shock. The patient tends to pass into a confused stuporous condition and finally into coma. An important point to note in patients with oligaemic failure is the danger of sudden death from changes of position (Evans *et al.*, 1944), such as sitting the patient up to examine the chest.

Water Absorption

I have mentioned above how the subjects of pure salt depletion are not thirsty, and how they develop anorexia, nausea, and vomiting. In patients with severe heat exhaustion to whom I endeavoured to give saline by mouth I noted that often the saline, if not vomited, would stay in the stomach, which could be seen, through the abdominal wall, to be greatly distended. There seemed to be definite pylorospasm coupled with atony of the stomach. Other observations led me to believe that a tendency to spasm of the sphincters and lack of tone of the gut prevailed throughout the alimentary canal. I consider that the delayed diuresis of salt depletion may be largely a question of delayed absorption. It is my impression that fluids remain much longer than usual in the stomach, and that even when the pylorus is passed absorption is slow. I have recently had under my care a case which was typical of many others I saw before the war without appreciating their significance.

A woman aged 58, with pyloric stenosis due to old duodenal ulceration, and with the usual history of vomiting large amounts, had been vomiting intermittently for 10 years and daily for about a month. On admission she was stuporous and appeared to be dehydrated, and there was loss of skin elasticity. Her blood pressure was 90/50 mm. Hg, the plasma chloride, as NaCl, was 286 mg., and the blood urea 190 mg. per 100 ml. During the first three days of treatment she received intravenously 23 pints of isotonic saline solution (= 13 litres containing 110 g. of NaCl) and 8 pints (4.5 litres) of isotonic glucose solution; she also was given 3 pints (1.7 litres) of water by mouth. Owing to incontinence, urine measurement was not possible during the first two days. Her output of urine on the third day was 33 oz. (937.6 ml.) containing 3 g. of NaCl. By the fourth day she appeared almost recovered and the plasma NaCl had risen to 524 mg. per 100 ml. Intravenous fluid administration was stopped, and on this and the next four days she received by mouth 19 pints (11 litres) of fluid plus some solid food. There was no abdominal distension; therefore this volume had successfully passed her previously impassable pylorus. Pyloric stenosis was subsequently confirmed by barium meal and at operation.

Sanchez-Vegas and Collins (1946) have reported upon 50 patients with pyloric obstruction due to duodenal ulcer. Gastric retention disappeared in all except 3 under treatment, averaging five days, in which the chief measure seems to have been administration of large amounts of sodium chloride in hypertonic solution. The main purpose of the communication of these authors was not to make the point, which I am tentatively

FIG. 10.—The vicious circle suggested in cases of salt depletion with pyloric obstruction.

suggesting, that pylorospasm may be a consequence of salt depletion, but to urge that urine chloride estimations are a better guide to progress than plasma estimations.

My observations on cases of heat exhaustion suggest that patients with pyloric obstruction may enter a vicious circle (Fig. 10). There is scope for radiological investigation of gastro-intestinal function, especially the emptying time of the stomach, in salt-depleted volunteers. It seems that salt depletion may also explain many cases of "acute dilatation of the stomach" and perhaps some of "paralytic ileus" in which there is no peritonitis. The effect of saline on peristalsis has been demonstrated by Eitel and Loeser (1934) and by Perazzo (1937).

B. Effects due to Disturbance of Acid-Base Equilibrium

When there is considerable asymmetrical loss of sodium or chlorine ions—such as the predominant loss of sodium in diarrhoea or the predominant loss of chlorine in vomiting (Fig. 8)—there is a disturbance of acid-base equilibrium, with a relative increase of bicarbonate with respect to chlorine in vomiting and a relative recession of bicarbonate in diarrhoea. The carbon-dioxide-combining power (alkali reserve: plasma bicarbonate) in ml. of CO_2 per 100 ml. of plasma may be over 100 after vomiting or under 20 after diarrhoea. As Gamble (1937) has said, the bicarbonate ion plays a "mendicant" part: it takes what base it can get.

Much attention has been directed in the past to these tendencies to alkalosis and acidosis, and many of the symptoms which have just been described, and attributed to loss of electrolyte osmotic pressure, were stated to be due to the distortion of acid-base balance. However, this is surely untrue, because all the manifestations, here called osmotic, may be seen in cases of severe sweating where there is no gross acid-base change; also all the manifestations are common to cases of either severe vomiting or severe diarrhoea. Indeed, apart from the different direction of ejection of alimentary secretions, there is little difference in the clinical picture presented by sufferers from emesis or catharsis. It is true that alkalosis or acidosis may have some intensifying effect on the anorexia, nausea, and vomiting which are certainly prominent enough in simple symmetrical sodium and chlorine depletion (sweating). Alkalosis may increase cramps through its tendency to cause tetany, but cramps are quite a marked feature of choleraic diarrhoea, in which there is acidosis, and are a characteristic feature of the salt depletion due to sweating. The only effect which seems mainly or wholly attributable to alkalosis or acidosis, when caused by vomiting or diarrhoea, is perhaps some difference, of little practical significance, in the character of the breathing.

The mechanism of death in pure salt depletion seems quite clearly to be circulatory failure due to a combination of reduction of blood volume and increase of blood viscosity.

Quantitative Correlation of Clinical and Pathological Manifestations with Amount of Salt Deficit

The ability to estimate the degree of deficit from the overall picture presented by the patient is a matter of great practical importance. It is so because, so far as I am aware, there is no other reliable index. The concentration of plasma chloride, expressed as NaCl, is often used as an index, but a little reflection will show, as Abbott (1946) has pointed out, that the concentration of chlorine or sodium cannot mirror the total loss of these ions because adjustment mechanisms, especially renal, work to preserve, so far as is possible, normality of concentration in the effort to sustain the prime essential—isotonicity. When plasma

sodium and chlorine tend to fall the kidneys excrete water and, as already explained, there is sacrifice of plasma volume in order to preserve electrolyte concentration. How, then, can any measure of this concentration determine total deficit? Significant falls in plasma chlorine or sodium values only occur when total deficits are very gross. This point is well brought out by Sanchez-Vegas and Collins (1946) in their study of 50 cases of pyloric obstruction. All these patients had either very little chloride in their urines or none at all, and all required large amounts of sodium chloride to restore renal excretion of chloride to normal. Yet only 10 out of the 50 had plasma chloride concentrations, expressed as NaCl, lower than 510 mg. per 100 ml., although 29 had blood urea concentrations exceeding normal limits.

From the basis of personal experience I would suggest that cases of salt depletion may be divided into three grades:

(1) *Slight to Moderate Depletion*.—Low concentration or absence of chloride in the urine (except in Addison's disease), with lassitude and maybe giddiness and orthostatic fainting, indicates a deficit of up to 0.5 g. per kilo of body weight; in a 70-kg. man this represents a deficit of up to 4 litres, or 7 pints, of isotonic saline.

(2) *Moderate to Severe Depletion*.—Absence of chloride in the urine (except in Addison's disease), with lassitude, giddiness, and fainting; with anorexia and maybe nausea and vomiting; and with a fall of blood pressure but with a systolic pressure above 90 mm. Hg. points to an approximate deficit of 0.5 to 0.75 g. per kilo, which in a 70-kg. man would represent a deficit of 4 to 6 litres (7 to 11 pints) of isotonic saline.

(3) *Severe to Very Severe Depletion*.—Absence of chloride in the urine (except in Addison's disease), with apathy, stupor, vomiting, and a systolic blood pressure of less than 90 mm. Hg. suggests an approximate deficit of 0.75 to 1.25 g. per kilo—that is, a deficit of 6 to 10 litres (11 to 18 pints) of isotonic saline in a 70-kg. man.

The above divisions represent a very rough classification and are put forward merely as a crude guide to appraisal of salt deficit. There will be individual cases forming exceptions, but, at the same time, it is suggested that even rough clinical criteria may be better than seemingly accurate and scientific measurements of plasma concentrations of sodium and chlorine—not, of course, that these should be omitted.

A Useful Test

It will be noticed that absence of chloride in the urine is accorded a place of first importance. For several years I have used the very simple quantitative test described by Fantus (1936). This can be done in a minute, and is simpler than testing for albumin or sugar as no heating is needed. The requirements for performing the test are: a small test-tube; a small pipette ("fountain-pen filler"); 20% potassium chromate solution; 2.9% silver nitrate solution; some distilled water for rinsing the pipette. It is essential that the same pipette shall be used throughout so that the size of the drops shall be the same.

Technique.—Ten drops of urine are measured into the test-tube. The pipette is rinsed and one drop of potassium chromate solution—the indicator—added. The pipette is rinsed again, and then the silver nitrate solution is added a drop at a time and the test-tube shaken after the addition of each drop. The end-point is a sharp colour change from yellow to brown. The number of drops needed to produce the end-point gives the concentration of chloride in the urine expressed as grammes of NaCl per litre—for example, 5 drops = 5 g. of NaCl per litre. It is important to make a preliminary test with distilled water to make sure that there is no contamination of the chromate with chloride.

This very simple test is most useful not only in diagnosis but also in the control of treatment. When used for

diagnosis, chloride should be regarded as "absent" if the end-point colour change occurs with the first drop. This may happen in normal persons if the urine is very dilute because a large amount of fluid has been taken, but if it occurs in the absence of diuresis it is indicative of salt depletion, though the depletion may not have reached a degree sufficient to cause symptoms. In concentrated urine (with a specific gravity of more than 1020) less than 3 g. per litre suggests depletion. If the urine contains 5 g. per litre or more the patient, except in Addison's disease or during the administration of intravenous saline, is unlikely to be suffering from salt depletion. The test was found useful in India and South-East Asia Commands during the war in the detection of salt-depletion heat exhaustion and of salt depletion complicating many medical and surgical illnesses. In M.I. room work it was valuable in the hot weather as a quick method of separating cases of salt-depletion heat exhaustion from those of neurotic collapse and ill-health from other causes. White (1943) so applied it at Air Headquarters, Delhi, during June and July, 1943, to a series of 140 patients, of whom 29 were proved to be suffering from heat exhaustion. Its potential scope is not as great in temperate as in hot climates but is still enormous, for it is applicable in the diagnostic assessment and therapeutic regulation of all cases of threatened or actual dehydration or of overt oedema. Whenever fluid tends to be lost from or retained in the body, the measurement of urinary excretion of sodium chloride is as important, in many instances more important, than the measurement of urine volume. It is my belief that the means for performing this simple test should be available in all ward test-rooms and in every doctor's bag. A similar simple quantitative test for sodium would be even more valuable if it could be evolved.

(To be concluded, with a full list of references, in our next issue.)

MEDICAL ASPECTS OF PENICILLIN TREATMENT

BY

B. A. YOUNG, M.D.

Medical Superintendent, St. Alfege's Hospital, Greenwich

This paper is based on a survey of the cases treated in the medical wards of St. Alfege's Hospital with systemic penicillin alone, or with systemic and local penicillin combined, from January, 1945, to September, 1946. In all, 129 patients were so treated. The cases are unselected, except that they were considered worthy of penicillin therapy on clinical or bacteriological grounds, and are consecutive cases. They were not the subject of special investigation as part of an experimental trial of penicillin treatment, but we have been continually alive to the fact that each case undergoing penicillin treatment is, even at the present time, an experiment in a new form of therapy, the application of which is by no means finally settled.

Our first regular supplies of penicillin became available in April, 1945, when we were allocated 5,000,000 units a month. This quota was gradually increased until at the time of writing our monthly consumption is between 150,000,000 and 200,000,000 units. The majority of cases were therefore treated in 1946. The 129 cases include the most serious and potentially fatal illnesses treated. An analysis according to the final diagnosis is shown in Table I.

The pneumonias form the largest group with 53 cases. Carbuncles, meningitis, and empyema are the next largest with 13, 11, and 10 cases respectively. The appearance of

carbuncles, cervical adenitis, and diabetic gangrene among a series of cases treated in medical wards alone is part of the revolution in medical practice which penicillin has

TABLE I.—Medical Cases Treated with Systemic and Local Penicillin

Disease	No. of Cases	Disease	No. of Cases
Respiratory diseases:		Streptococcal tonsillitis	4
Pneumonia	53	Diabetic gangrene	3
Acute bronchitis	3	Perinephric abscess	2
Empyema	10	Cellulitis	1
Lung abscess	2	<i>Staphylococcus aureus</i>	1
Carbuncles	13	Septicaemia	1
Meningitis	11	Vincent's angina	1
Secondarily infected skin lesions	8	Acute pyelonephritis	1
Acute nephritis	6	<i>E. coli</i> septicaemia	1
Subacute bacterial endocarditis	5	Lobar pneumonia	1
Cervical adenitis	5	Pemphigus, chronic nephritis	1

brought about. Particularly, and most dramatically, in the case of carbuncles is this change of practice welcome, most of all to the patient.

Below is given an analysis of the results of treatment of the more important groups of diseases, with a brief description of the methods used, and such points of importance bearing on treatment that have been brought to light.

Pneumonia

From January to December, 1945, 12 cases of pneumonia were treated, with 5 deaths, and from January to September, 1946, 41 cases, with 3 deaths. This represents an overall mortality of 15%. Of the 53 patients 26 were over 40 years of age and 10 were under 6 months; and the majority were cases of great severity. During the first period penicillin was scarce and dosages were in consequence low. The figures for the second period, giving a mortality of 7.3%, must be regarded as very gratifying. Four of the eight fatal cases were moribund on admission and death occurred within 24 hours; three had signs of congestive cardiac failure on admission; and one was a premature infant. Seven patients who recovered, two who improved, and five who died were also treated concurrently with sulphonamides. Fifteen cases received penicillin after the infection had failed to respond to sulphonamide therapy.

TABLE II.—Cases of Pneumonia

Variety	Recovery	Improvement	No Change	Death	Total
Lobar, unclassified	11	3	—	3	17
Bronchopneumonia, unclassified	11	—	2	5	18
Pneumococcal	4	—	—	—	4
Staphylococcal	4	1	—	—	5
Streptococcal	4	—	—	—	4
Mixed flora	4	—	—	—	4
Atypical	—	—	1	—	1
Total	38	4	3	8	53

The bacteriological classification of the cases is not complete, partly because many of the patients treated were too ill or too young to produce sputum, and partly because we were loath to ask too much of our ever-willing but over-worked laboratory staff. We now regard a bacteriological diagnosis, with a penicillin-sensitivity test of the organism, as an essential in the penicillin treatment of pneumonia. Such information offers valuable guidance to the physician, particularly on the necessity for increasing the dose or for prolonging the course, in a case not initially responding to penicillin. Without this information he works in the dark. It may be significant that of the cases bacteriologically classified none were fatal.

Indications for Penicillin Treatment in Pneumonia.—Administration of sulphonamides is still our method of choice. Our indications for penicillin are: (1) Cases with a bad prognosis owing to clinical severity with signs of extreme toxæmia, old age, and infancy, particularly prematurity. In these cases penicillin and the sulphonamides