marry, had delivered their children and their grandchildren, and had become "family" practitioners in the true sense of the word—voted 2,782 for and 5,249 against.

Unless the Act is amended what have our patients in store from the new Service under the present doctrine that the individual exists for the State? Willesden was the opening gambit. The recent overriding of a G.P. who was acting on the hospital consultant's advice on diet for a patient is quite logical—if one admits the State doctrine. The patient had a cancer and would have died anyway, so why feed a "useless mouth" when world rations are so short? The beloved physician does his traditional best for his patient. The State administrator to whom the patient is merely a "case" does his impersonal duty to the State. Granted the doctrine, what use has the State for incurable tuberculous patients? Why give them rations needed by the healthy? Indeed, why feed the useless mouths of those too old to work? This may become a major political problem when, owing to the birth rate having fallen during the last decades, there will not grow up a sufficient number of adults to support a rapidly increasing burden of old-age pensioners. There is nothing illogical in the train of thought. Euthanasia for the useless to prevent the diversion of rationed food from the healthy and productive may not be so impossible as may seem once the State doctrine has been forced on us.

Before 1939 many believed that another war was too ghastly to be possible. Such "realists," with their heads in the sands of their daily round and trivial task, were disillusioned. Many could not believe Belsen to be anything but propaganda until the irrefutable facts unwillingly convinced them. In 1945 few would have thought that the Willesden episode could happen here, but it has. It is our Munich, and we have to act now. What is the answer? Just that the individual does not exist entirely for the State, and the present political measure of potential tyranny disguised as a Health Act must be amended to give all a first-class Health Service—I am, etc.,

Derham, Norfolk.

E. PUDDY.

Medical Certification

SIR,—I recently issued a patient of mine with a certificate stating that he required a "thermos" flask for medical reasons. On presenting his certificate he received the following letter from the Board of Trade.

Dear Sir/Madam,

I have your application for a "thermos" flask permit, with which you enclosed a medical certificate. In view of extremely limited supplies of flasks, however, the exact nature of the illness must be advised; will you please therefore supply this information, stating in addition why a flask is essential. . . .

As I was asked to deal with the matter I replied to the Board of Trade as follows:

Dear Sir,

Reference application for "thermos" flask permit for Mr. X., of Eastbourne, this patient is suffering from myocardial degeneration with oedema, cerebral thrombosis, and bronchitis. He is a very ill man and his health is failing. He requires an odd hot drink in the night, and since his wife is worn out by attending to him during the day it seems to me that a "thermos" flask is a reasonable request and that a permit should be forthcoming immediately.

I wish to lodge a personal protest: (1) The nature of a patient's illness is a personal matter, and I am at a loss to see why such private details as these should be disclosed to the Board of Trade (quotation from your letter: "the exact nature of the illness must be advised"). (2) Doctors these days are overburdened with the signing of certificates, forms, etc., and an effort should be made to lessen this work—I did not sign the initial certificate either indiscriminately or for fun. (3) I cannot understand how an officer of the Board of Trade behind a desk in London can possibly assess a patient's condition in Eastbourne. I earnestly plead that steps be taken to eliminate this bureaucratic stupidity.

These details I send you not as a moan from a doctor but simply to bring them to your notice in case you are not already aware of the situation. My letter alone obviously will do no good, but backed by a protest from the B.M.A. it is just possible that the Board of Trade might remove their long noses from matters which do not concern them.—I am, etc.,

Eastbourne.

BASIL S. KENT.

To Negotiate or Not

SIR,—Contained in the annotation "Plebiscite Secret" (Nov. 23, 1946, p. 779) is the statement: "If the majority of the profession say 'No' on their plebiscite forms then the Minister will have no one else but himself to thank for such a refusal to operate the new Health Service Act, because this is what the answer 'No' will mean." Presumably this "refusal to operate" implied refusal to negotiate the Regulations of the Act as it stood and as it stands now. I do not know how closely this statement expressed the intentions of the Council, but subsequent events showed that the Council was in fact prepared to follow the wishes of the simple majority in the plebiscite.

The more recent decision to reopen negotiations, so lamely represented as the result of a more conciliatory attitude of the Minister, has restored in the minds of many practitioners the old mistrust of the Council's ability to represent them, and aroused bitter recollections of previous vacillations in policy. If in fact the Council's change of tune does bear relation to the expressions of personal opinion by the Presidents of the Royal Colleges it seems reasonable to point out that these were only another three votes to offset the larger majority of "Noes" expressed by doctors with closer associations and deeper knowledge of the public, whose best interests are at the heart of us all. So yet again:

"... the native hue of resolution

Is sicklied o'er with the pale cast of thought."

—I am, etc.,

London, E.C.1.

ANTHONY ALMENT.

Aspiration of Acute Pleural Effusions

SIR,—I would like to congratulate Dr. E. A. Wood on his letter (Jan. 25, p. 159) concerning the aspiration of pleural effusions. I wish that the indications and contraindications for removal were as clearly and correctly stated in all medical textbooks.

In the hospitals I visit I have often found that the M.O.s have only a vague idea of when and when not to aspirate. I do feel, however, that pleural effusion cases should always have a small specimen of fluid withdrawn for diagnostic purposes.—I am, etc.,

London, N.7.

J. WALLACE CRAIG.

Curare

SIR,—I hesitate to enter into argument with Dr. Massey Dawkins (Jan. 18, p. 171) and Dr. A. J. Gray (Feb. 1, p. 196) on the subject of curare, for their experience with this drug must surely be much wider than mine. At the same time I cannot but disagree with Dr. Massey Dawkins over the post-operative condition of patients who have been given curare. In my small series of cases I can only feel delighted with the results, and, on questioning, the nursing staff appear to be equally satisfied. Dr. Gray, in his excellent reply, deals with the points raised, but perhaps one might be allowed to elaborate further.

With each administration of an anaesthetic agent there are well-known difficulties and dangers to be borne in mind, avoided, and, less frequently, remedied. When dealing with curare the points to be considered are profound muscular relaxation with limited respiratory excursion, which together lead to some degree of anoxaemia, increased carbon dioxide tension, decreased venous return to the heart, and some fall in blood pressure. All these are dependent upon the dose of curare and the speed with which it is given. Little has been written about the rate of administration, although L. Watter (1944) reports that in one case a rapid injection of 2 ml. of "intocostin" near completion of an operation resulted in severe respiratory depression, and Lundy (1945) states that rapid injection can prove fatal. It would seem wise, then, always to inject the drug slowly. As regards size of dose there is a wide variation between 9 mg. and 30 mg., in my experience, required to produce upper abdominal relaxation in adults. The dose should not depend upon the patient's weight but rather upon the patient's needs.

With larger doses of curare there is considerable limitation of respiratory excursion, and it is not sufficient to stand some yards away from the patient and observe that he is breathing. There comes a time when this breathing is insufficient for the patient's metabolic needs, and so it is imperative, long before respiratory arrest is likely to set in, that the anaesthetist intermittently compresses the rebreathing bag, thereby favouring respiratory exchange and avoiding pulmonary stasis. A high concentration of oxygen in the inspired gases seems to be beneficial, and should there still be shallow breathing at the end of the operation one should not hesitate to give prostigmin. When breathing has returned to normal, hyperventilation with carbon dioxide and air will introduce the inert diluent nitrogen into the alveoli and thus help to guard against atelectasis. If these precautions are taken in every case then there should be no need to worry about respiratory depression during the immediate post-operative period and little fear of pulmonary complications at a later date.

Since curare is non-toxic, then in an anaesthetic sequence in which it is used one need only consider the slightly toxic effects of the small amount of anaesthetic agent required to keep the patient asleep. Dr. Gray quite rightly states that Crile's theory has been invalidated. R. T. Knight (1944) goes further by saying "... that deep general anaesthesia is more damaging and more shock-producing than even the trauma of surgery." Surely, then, the maintenance of light anaesthesia is a prophylactic against shock. Light anaesthesia is easily maintained with non-toxic nitrous oxide, but I do not think that this is entirely satisfactory with curare, since the shallow breathing associated with the latter tends to increase the anoxaemia usually present with the former.

The majority of the cases in which I have used curare have been upper abdominal operations and/or ill patients, and in these cases an intravenous drip is often required before, during, and after the operation. "Pentothal" is always used, the initial dose being supplemented when rise in pulse rate—a sign of lightening anaesthesia—occurs. Usually an endotracheal tube is passed under direct vision and pure oxygen is administered throughout the operation. Under these conditions perfectly satisfactory anaesthesia and convalescence are ensured.—I am, etc.,

Truro, Cornwall.

G. V. S. WRIGHT.

SIR,—Dr. Massey Dawkins (Jan. 18, p. 111) has drawn attention to three post-anaesthetic complications following the use of curare—viz., shock, respiratory depression, and paralytic ileus. While these conditions unfortunately do occur, may they not be attributed to an overdose of curare—that is, a quantity in excess of that necessary to produce the desired effect?

At présent there appears to be a tendency to use curare in dosage sufficient to produce immediate absolute relaxation, with little regard to individual susceptibility. The latter may be of more importance than the weight or physical fitness of the patient, and the adoption of a small initial "test dose" (5–10 mg. tubocurarine chloride), followed by minimal additions, may lessen these complications. I have observed marked hypotension following the administration of curare in light and deeper anaesthesia—even before the incision has been made. This is presumably due to the action on the autonomic nervous system in the curare-sensitive subject.—I am, etc.,

Llanelli, Carmarthenshire.

D. S. JONES.

Curare or Magnesium Sulphate?

SIR,—We have used magnesium sulphate as a relaxing agent in anaesthesia in a few cases with unconvincing results. One woman aged 32 was given "pentothal" 0.5 g. followed by 20 ml. of 25% magnesium sulphate solution intravenously and nitrous oxide and oxygen for appendicectomy. Relaxation appeared to be good, but it is difficult to say what part the magnesium sulphate might have played in this. A man aged 38 was given "pentothal" 0.5 g. followed by 25 ml. of 25% magnesium sulphate with cyclopropane and oxygen for repair of an incisional hernia of the upper abdomen. In spite of deep cyclopropane anaesthesia relaxation was not as good as with curare. In another case laryngoscopy and orotracheal intubation were performed without difficulty in a healthy woman

aged 24 after "pentothal" 0.5 g. and 20 ml. of 25% magnesium sulphate. Laryngeal irritability appeared to be less than after "pentothal" alone.

There were no adverse side-effects and respiration was not obviously affected in any of the cases. Magnesium sulphate has been used to mitigate convulsions in shock therapy,¹ and it is said to have bronchodilator and antihistamine effects.^{2,3,4} Anaesthesia has been produced with magnesium sulphate,⁵ but muscular relaxation appears first. It seems that if magnesium sulphate is to be used instead of curare much larger doses must be used than the 0.5–1 ml. of 25% solution suggested by Dr. W. F. Cooper (Feb. 1, p. 197).—We are, etc.,

GEOFFREY ORGANE.

IAN C. W. ENGLISH.

London, S.W.1.

REFERENCES

- ¹ Yaskin, H. Edward, *Arch. Neurol. Psychiat.*, 1941, **46**, 81.
- ² Haurey, V. G., *Pharm. Exper. Therap.*, 1938, **64**, 58.
- ³ Monchal, Lumière, *C. r. Soc. Biol.*, 1937, **124**, 178.
- ⁴ Haurey, J., *Lab. clin. Med.*, 1940, **26**, 340.
- ⁵ Meltzer and Auer, *J. Amer. med. Ass.*, 1916, **67**, 1131.

Prognosis of Hypertension

SIR,—Your contributor writes in the annotation on "Prognosis of Hypertension" (Jan. 25, p. 144), discussing Bechgaard's monograph: "By hypertension was meant a blood pressure of not less than 169/90; the average was 190/110 taken after 15 minutes' rest."

But although Bechgaard's material consists of patients who had a blood pressure of 160/100 or a systolic blood pressure of 180 mm. at the first examination, he defines normal blood pressure as being 140/90 mm. He regards people with a B.P. of more than 140/90 mm. as hypertonics, even if they be over 40 years of age. He admits that "with this limit a very large number of older people (in case of the oldest more than 50%) are regarded as hypertonics." This would hardly be accepted by English and American observers, who regard as raised blood pressure in people over 50 a reading of 170/105 taken under proper precautions.—I am, etc.,

London, S.W.3.

V. C. MEDVEI.

Gynaecomastia

SIR,—It may be of interest to your readers to know that this condition is relatively common among lepers, and cases may be seen in any large leper colony at almost any time. It has been described by Dr. James L. Maxwell in his textbook *Leprosy: A Practical Textbook for Use in China*. It is associated with leprosy leading to atrophy of the testis, followed by partial or complete sterility. It is commonly a bilateral condition and appears to be entirely painless.—I am, etc.,

London, S.W.16.

STEPHEN D. STURTON.

"Anaesthesia" or "Analgesia"?

SIR,—I am not concerned with the polemics of Dr. F. W. Roberts (Dec. 28, 1946, p. 1007) and Dr. J. N. Fell (Jan. 18, p. 115), but I must take the latter to task for being misleading, apparently because he thinks anaesthesia is more euphonious than analgesia and would make a more charming name for a girl than would the latter. The statements he makes might cause some members of the profession to think that those who chose these Greek words for descriptive purposes in their anglicized forms had been guilty of careless or bad selection, whereas this is not the case.

Dr. Fell states that there is no distinction between the words in Greek; that distinction does, however, exist in precisely the way which is intended in their anglicized forms. Anaesthesia (*ἀνασθησις*) is used by Plato as a noun to mean "want of perception," as meaning "stupor," and in the same work he uses it adjectivally to mean "not subject to the senses" or "insensible"; it is also used as an adverb in Isocrates in an exactly similar way. On the other hand analgesia (*ἀναλγησις*) is used to describe "want of feeling," and the adjective *ἀνάλγητος* to describe "without pain"; but let us discard the particle *ἀν* and look at the parent words. *αἰσθάνομαι* and *αἰσθησις* are used without exception to describe "perception" in one form or another, while *ἀλγέω*, *ἄλγησις*, and *ἄλγος* are always used to convey the "sense of pain."