Correspondence

Care of Children in Hospital

SIR,—I have read and re-read with interest and profit the lecture given by Prof. Spence on the care of children in hospital (Jan. 25, p. 125). I am sure it will be of great interest and a stimulus to all who are engaged in this important work. Prof. Spence has given us much food for thought—so much indeed that it would be impossible to comment fully upon it within the space of a letter. One hopes that it will be a basis of much serious discussion and, if necessary, controversy among people who are interested in children's hospitals. I should like, however, to comment on one feature, and that is the "atmosphere" of a children's ward, so ably described by Prof. Spence. This masterly description should cause serious reflection and perhaps give rise to a sense of guilt. The state of affairs described by Prof. Spence is unnecessary but often only too true, and I would suggest that we, as doctors, are not entirely without blame. Too often have we looked upon the sick child as a patient presenting interesting clinical features requiring investigation and treatment. Too often do we tend to forget that while the child is in hospital the ward which houses him is his temporary home. The child is entitled, while in hospital, to have that feeling of security and affection which every home should provide. I feel that we, as doctors, can set an example. If the doctor will try to make the children realize that he is their friend and interested in them as human beings as well as patients, then the ward round might cease to be a distraction and become an eagerly awaited event.

There are numerous other ways in which the medical staff can contribute to a "happy atmosphere," provided they have an intimate knowledge of the working of a children's ward. Would it be too much to suggest that the medical student might acquire some of this knowledge by working on a children's ward during his training in paediatrics—by working I mean carrying out nursing duties under the ward sister for several days. I venture to suggest that in this way the future paediatrician would learn some valuable lessons in practical paediatrics.

Prof. Spence has pointed out the demands made on a nurse's time in carrying out the nursing treatment of small children, and has stressed the importance of an adequate nursing staff. As he rightly says, we do not realize sufficiently the extent to which an inadequate nursing staff may adversely affect the welfare of sick children. How can we expect a nursing staff—whose numbers are insufficient for the job—to nurse the dangerously ill child, nurse and comfort the fretful not-so-ill child, and nurse, comfort, and entertain the child in the convalescent stage of its illness? In his lecture Prof. Spence has shown one way by which the burden may be lightened. This has one disadvantage in that it is not always reliable. Many mothers of sick infants have other family responsibilities—especially the care of the young children—which they cannot lay aside easily. Could his suggestion be carried a stage further (or should I say backward?) and the help of expectant mothers be enlisted? I have in mind the mother expecting her first baby. Her family responsibilities are light, and the experience she would gain would be useful in the months to come. It might even be the means of avoiding a future case of "mismanagement and feeding error."

Another method of easing the burden on the nurses' shoulders would be to cater for the convalescent child by providing "diversional occupational therapy." This once fashionable term has fallen into desuetude. Let us revive it in the children's hospitals, but with safeguards. I would suggest that every children's ward should have the services of a teacher specially trained in bedside education methods. Such a teacher must recognize that her duty, as part of the hospital team, is the special care of the child in the convalescent stage and to render that stage as interesting, pleasant, and as profitable as possible.

No doubt, Sir, many of your readers have other and better suggestions than mine to render more tolerable "the daily rhythm of anxiety, wonder, apprehension, and sleep." To my

mind it is most important that the train of thought started by Prof. Spence should not be allowed to lapse. It is not necessary to wait for the structural alterations which most of us would like to see carried out. Let us remember that "four walls do not a hospital make"; it is the character and capabilities of each member of the staff which alone can provide the right "atmosphere." Let us never be satisfied with any improvement we make but always keep in mind that "achievement is but a milestone in the march of progress, the end of the journey lies ever ahead."—I am, etc.,

W. E. CROSBIE,
Medical Superintendent.

SIR,—I have read with great interest Prof. J. C. Spence's article (Jan. 25, p. 125) on the care of children in hospital, especially the question of nursing, as I am myself faced with an incredible shortage of nurses in the babies' ward, where there is barely one nurse for every seven ill babies. I was very anxious to try his method of getting their mothers to look after them and have surveyed the situation with that view in mind. I found that of eight bottle-fed babies who are in the ward at present, the mothers of four have five other children to look after at home, the mothers of two are ill in other hospitals, another had a partial gastrectomy here the day before, and the last baby's mother is suffering from puerperal psychosis. I envy Prof. Spence in being able to treat babies whose mothers can spare all that time for them, but I doubt that it is a practicable proposition in most children's departments.—I am, etc.,

Dartford, Kent.

JOHN LORBER.

Chemotherapy of Virus Diseases

SIR,—May I say that I agree with Dr. W. N. Leak's letter (Jan. 25, p. 160) protesting against the statement in the leading article of Dec. 7, 1946, that there is scarcely a hint that viruses are susceptible to chemotherapeutic attack? In the case of both vaccinia and smallpox viruses there is definite evidence that they are extremely susceptible to the action of potassium permanganate, which inactivates them even when diluted to 1:100,000. This extreme susceptibility to permanganate was described and illustrated in the Medical Research Council's Special Report, Series No. 98, 1925, and greatly interested the late Sir Andrew Balfour for the following reason. During an outbreak of severe smallpox in the Sudan the cases had been treated with compresses soaked in permanganate and applied to the skin lesions with amazingly successful result. The object of the doctors was to counteract any secondary infection, but Balfour saw at the time that something more than this must be involved to account for the beneficial effect. When he read the report in question and realized the high degree of susceptibility of smallpox virus itself to permanganate, Balfour was more delighted than I ever remember to have seen him. The pronounced viricidal action of permanganate on vaccinia virus *in vitro* ceases in the presence of glycerin. From this it would seem likely that the well-established value of glycerin in conserving viruses is due to the protection it affords at their weakest point—hypersensitivity to oxidation. Bacteria are far more resistant to permanganate than viruses.—I am, etc.,

Molesey, Surrey.

M. H. GORDON.

A Review of the Dietetic Factors in Liver Disease

SIR,—While admiring the painstaking effort and industry of Prof. L. J. Witts in summarizing the existing knowledge on the above subject, I was a little surprised to read some of his categorical statements on the clinical features of acute necrosis of the liver (Jan. 11, p. 45). He states, "Jaundice is intense." Surely one of the most variable features of the condition is the depth of the jaundice. Sometimes it is slight in the acute stage, and in general experience it is seldom as deep as in cases with obstructive jaundice, such as is seen, for example, in carcinoma of the pancreas or stricture of the common bile duct. It was perhaps for that reason that the old term "icterus gravis" long ago fell into disuse. This variability in the depth of the jaundice in acute liver necrosis was stressed by my father in his Lumleian lectures on toxic jaundice in 1931 (Willcox, W., *Lancet*, 1931, 2, 57). He maintained that the grave signs were produced by the liver deficiency, which he ventured to call