and Young found the sodium and chlorine clearances in full-term babies aged 7-14 days to be about one-fifth of the value they would be likely to acquire in later life. Still lower clearances were found in premature babies (Young, Hallum, and McCance, 1941).

As is now well known, the cortex of the suprarenal glands plays an important part in regard to renal excretion of salt. In experimental adrenalectomy or in Addison's disease there is an excessive excretion of sodium chloride, and in consequence the plasma level falls. Excretion may continue in Addison's disease even when the plasma chloride, expressed as NaCl, is below 500 mg. per 100 ml. (Anderson and Lyall, 1937). The subject has been reviewed by Loeb (1941), whose own work in this field has been of such importance. Detection of the tendency to excessive salt excretion, under standard conditions of salt restriction, is the basis of the well-known test for Addison's disease which has been devised by Cutler *et al.* (1938a, b). There is similarity in failure of hormonal control causing excessive excretion respectively of water and salt in diabetes insipidus and in Addison's disease. However, in the former the body is saved from very serious consequences by the protective mechanism of thirst, while in the latter there is no equivalent salt-craving. In passing it should be observed that in diabetes insipidus there is no increase of salt excretion or fall in plasma chloride level; this was the finding of Blotner (1941) in 22 cases of that disease.

(Part II will be published in the next issue.)

THERAPEUTIC POSSIBILITIES OF RADIO-PHOSPHORUS

BY

J. S. MITCHELL, M.B., Ph.D.

Radio-phosphorus (P^{32}) has been used since 1936 in the treatment of patients with chronic myeloid and lymphatic leukaemia, polycythaemia vera, lymphosarcoma, and various related diseases. Most of this work has been done in the U.S.A. Important summaries of the American results have been given by Low-Beer, Lawrence, and Stone (1942); Kenney (1942); and Reinhard, Moore, Bierbaum, Moore, and Kamen (1946). Of great interest is a paper from Stockholm by Lindgren (1944).

Clinical trials of the therapeutic possibilities of radio-phosphorus have not yet been carried out in Great Britain, although tracer studies with P^{32} made by cyclotrons are in progress. However, it now seems reasonably certain that within the next year we shall be fortunate enough to obtain adequate and regular supplies of P^{32} made by means of a "pile." The isotope P^{32} is produced in the pile from ordinary phosphorus (P^{31}) by slow neutron capture and is used in the form of isotonic Na_2HPO_4 solution (15 mg. per ml.), with initial radioactivity of about 300 microcuries per ml. The pile can produce a specific gravity of one-third curie of P^{32} per g. of phosphorus. The P^{32} given therapeutically should be accompanied by the smallest possible amount of P^{31} .

Dosage

The radioactivity of the phosphate solutions used can be determined by means of a Lauritsen electroscope calibrated with a uranium standard, but for clinical investigations a Geiger counter is essential. It appears most satisfactory to give the radioactive phosphate solution intravenously, usually in unit doses of the order of 0.5-2 millicuries. However, oral administration is possible, and it is stated

that "it is the practice of most workers who give P^{32} orally to assume that 75% of any given dose is absorbed."

The physical properties of P^{32} are well known. This isotope has a half-life of 14.3 days and emits only beta particles. The maximum energy of the beta particles is 1.71 mev.; the mean energy of the beta particles is of great importance in clinical applications and has the approximate value of 0.70 mev.

Although as yet most of the clinical dosage of P^{32} has been necessarily empirical, it seems highly desirable to correlate the dose of radioactive isotopes with the roentgen unit. An important paper on this subject was published by Marinelli (1942). It can be shown that 1 microcurie of P^{32} per g. of tissue delivers 43 r. per day, 1 g. roentgen corresponds to 1.13 microcuries destroyed per kilogram of tissue, and if 1 millicurie is retained for 24 hours by a patient weighing 70 kg., approximately 0.6 roentgen equivalents of whole body radiation are delivered. In radioactive isotope therapy it is useful to measure the differential absorption ratio for the isotope, which may be defined as the ratio of the concentration of the radioactive material in a particular tissue to the mean concentration in the body as a whole. The possibility of delivering useful doses of radiation in any particular case may be determined by measuring the differential absorption ratio by means of tracer studies (Kenney, Marinelli, and Woodard, 1941). In this way it was found impossible to deliver therapeutically useful doses of beta radiation from P^{32} to the primary tumour or metastases in cases of carcinoma of the breast and of osteogenic sarcoma. The present tendency in all diseases except polycythaemia vera appears to be to follow the "fractional method" of dosage of P^{32} outlined by Low-Beer, Lawrence, and Stone (1942). It seems that one can treat a typical case of chronic myeloid leukaemia with an initial dose of perhaps 1 millicurie of P^{32} , followed by five doses of 0.5 millicurie at three- to four-day intervals, and then doses of 0.5 millicurie weekly until the total white cell count has fallen to about 30,000 per c.mm. Then one or two fractions may be given subsequently at longer intervals. The whole clinical and haematological picture must obviously be taken into account. It seems likely that the above unit doses can be safely doubled in some cases, with reduction of the over-all time of treatment. The average treatment for the 39 cases of chronic myeloid leukaemia reported by Reinhard *et al.* (1946) corresponds to 11.8 millicuries of P^{32} administered in 78 days. In the treatment of polycythaemia vera it is recommended that the initial dose is 3.5 to 4 millicuries of P^{32} . This may suffice, but in some cases it is necessary, after waiting three months, to give a second dose of 1 to 3 millicuries; ancillary venesection is often advisable.

Therapeutic Value

It is not yet possible to assess with confidence the therapeutic value of radio-phosphorus, but it is evident that the earlier optimism is not yet justified. The conclusions reached by Reinhard *et al.* (1946) may be summarized as follows:

1. Radio-phosphorus is probably the best therapeutic agent available at the present time for polycythaemia vera.
2. The results of treatment by radio-phosphorus of patients with chronic myeloid leukaemia are probably comparable with those of the usual x-ray therapy but are not substantially better. With P^{32} , "freedom from radiation sickness is a practical advantage which patients who have had previous x-ray therapy appreciate." Great care appears to be necessary in treatment with P^{32} to avoid serious damage to the bone marrow.
3. "In the treatment of chronic lymphatic leukaemia radio-phosphorus is probably as satisfactory as, but not better than, roentgen radiation."

4. Radio-phosphorus is of no value in the treatment of acute leukaemias or of monocytic leukaemia.

5. "Hodgkin's disease, lymphosarcoma, reticulum-cell sarcoma, and multiple myeloma do not respond as favourably to P³² as they do to x-radiation."

6. The limited evidence available suggests that radio-phosphorus is not likely to prove to be a satisfactory agent for the treatment of carcinoma of the breast, malignant melanoma, lympho-epithelioma, Ewing's tumour, or mycosis fungoides.

However, the work of Kenney (1942) suggests that further clinical trials of radio-phosphorus in lymphosarcoma are desirable.

It seems doubtful whether radio-phosphorus, in the form of inorganic phosphate, will stand the test of time even in the treatment of chronic myeloid leukaemia. In his report to the United Nations Atomic Energy Commission on Sept. 23, 1946, Dr. C. P. Rhoads, the Director of the Memorial Hospital, New York, discussed the treatment of chronic myeloid leukaemia and concluded that "a tendency in most institutions is to discard the use of radio-phosphorus and to stick to conventional x-ray as being somewhat safer and more predictable in its results and attended by less hazards to the patient, even though it does cause some illness in the course of treatment."

In conclusion, one can quote again from Dr. Rhoads's report: "To have expected inorganic elements to be picked up so selectively and concentrated so adequately by cancer tissue as compared to normal tissue was a very optimistic hope. I am not surprised that we are somewhat disappointed, but we have a vast field before us. We are beginning to study the selective pick-up by cancer of a vast variety of organic compounds which can be synthesized in the laboratories of organic chemistry to contain radioactive isotopes of a variety of elements. I am very hopeful."

A final aspect of the therapeutic application of radio-phosphorus is the possibility of development of beta-ray applicators using P³², perhaps incorporated in plastics. A report on human skin reactions produced by the external use of radio-phosphorus has been published recently by Low-Beer (1946). "The estimated number of roentgen equivalents dissipated in the first millimetre of tissue is 4.3 per hour per microcurie per square centimetre area." The maximum range of the beta-particles from P³² is approximately 8 mm. in water or unit-density tissue. It is stated that half-value layer measurements have shown that about 48% of the radiation is absorbed in the first millimetre of tissue. "The results of these studies suggest that the use of radioactive phosphorus in the local treatment of certain superficial skin diseases is reasonable."

REFERENCES

- Kenney, J. M. (1942). *Cancer Res.*, **2**, 130.
— Marinelli, L. D., and Woodard, H. Q. (1941). *Radiology*, **37**, 683.
Lindgren, E. (1944). *Acta Radiol.*, **25**, 614.
Low-Beer, B. V. A. (1946). *Radiology*, **47**, 213.
— Lawrence, J. H., and Stone, R. S. (1942). *Ibid.*, **39**, 573.
Marinelli, L. D. (1942). *Amer. J. Roentgen.*, **47**, 210.
Reinhard, E. H., Moore, C. V., Bierbaum, O. S., Moore, Sherwood, and Kamen, M. D. (1946). *J. lab. clin. Med.*, **31**, 107.
Rhoads, C. P. (1946). *Medical Uses of Atomic Energy*, vol. 5.

Sixteen years have been added to the average life span of Americans was one point made by Dr. James A. Crabtree, acting Surgeon-General of the U.S. Public Health Service, in a recent address. He said that in the problems of communicable disease control, emphasis was shifting from the diseases of early childhood to those of middle and old age. Americans were taking a great interest in public health figures as the result of Selective Service findings, which drew attention to various national health defects. The control of venereal disease was progressing, and the mortality rate had decreased from 15 deaths per 100,000 in 1939 to 10.7 in 1945.

PINK DISEASE: A REVIEW OF 65 CASES

BY

T. N. FISHER, M.B., M.R.C.P., D.P.H.

Physician, Royal Manchester Children's Hospital; Physician for Children, Manchester Northern Hospital; Consulting Physician for Children, Warrington Infirmary.

Pink disease sprang into prominence in medical literature in the early years of the century and seems to have established itself in this country as a not very common but, nevertheless, regularly recurring member of the group of diseases incident to infancy and early childhood. I have reviewed the case histories of patients who attended my out-patient clinic at the Royal Manchester Children's Hospital during the 14 years from 1932 to 1945 to see if any further light can be thrown on the nature of this still somewhat puzzling ailment. During this period 65 infants and young children—27 males and 38 females—suffering from the disease came under my observation. I followed the progress of 48 of these from the time of their first attendance to the end of their illness. The remaining 17 I saw on only one or two occasions, but I ascertained the subsequent progress of 14; the other three I was unable to trace. Of the 62 children whose subsequent progress is known, 53 made a complete recovery and 9 died.

Incidence of the Disease

Year:	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945
No. of cases	3	4	5	4	1	5	3	7	6	6	7	2	5	7

There was a fairly even distribution over the whole period. The drop in numbers in 1936 and 1943 can have no significance where such small numbers are concerned, unless other observers have noted a similar feature.

Month of Onset of the Disease

Month:	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
No. of cases	8	4	6	5	4	6	5	4	7	6	2	5

These figures embrace the whole period and show that there was no demonstrable variation in seasonal incidence. The impression that it is commoner during the winter months can, I think, be explained by the greater severity of the disease, due to intercurrent infection, during that period of the year.

Age of Onset

Age in Months:	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	25	38	50
No. of cases	1	4	3	6	6	9	2	8	7	0	4	2	1	1	2	2	1	1	3	1	1

The figures show a rapid increase from the age of 4 months (the earliest age at which the disease occurred) to the end of the first year, and thereafter a sharp decline. The two children who succumbed to the disease in their fourth and fifth years were quite outside the usual age period in this series. A calculation of the average age of onset in each year throughout the period shows no significant trend in the behaviour of the disease towards an earlier or later susceptible age of choice.

Average Duration for Each Year

Year:	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945
Average duration in months	4.5	3.3	3.7	3.8	—	4.1	3.0	3.7	4.7	3.0	4.6	—	5.3	3.9

In 1936 only one case was seen (duration 3.5 months); two cases were seen in 1943, but the duration of the disease could not be determined sufficiently accurately for inclusion in this table.