MARROW BIOPSY

PREPARATION AND USE OF PARAFFIN SECTIONS FROM STERNAL-PUNCTURE MATERIAL

BY

D. F. CAPPELL, M.D. H. E. HUTCHISON, M.B.

AND

G. HARVEY SMITH, M.B.

(From the Pathology Departments of the Western Infirmary, Glasgow, and Royal Infirmary, Dundee)

[WITH PHOTOGRAPHURE PLATE]

The value of sternal puncture as a diagnostic measure in disorders of the haemopoietic system is now widely accepted by haematologists, and the procedure is generally recognized as being safe and causing little or no discomfort to the patient. The usual method of dealing with the biopsy material consists in the microscopical examination of marrow smears stained by one of the Romanowsky stains, a differential count of the nucleated cells, and an estimation of the leucocytogenic-erythrogenic ratio. Other workers carry out total nucleated-cell counts, and supra-vital preparations may be made if desired. These procedures, while of great value cytologically, do not always give a true picture of the marrow as a whole and give no indication of its architectural arrangement. A simple method of preparing good histological sections from sternal-biopsy marrow would be of great service and the sections would yield valuable additional information about the architecture of the marrow, especially in cases of hypoplasia or aplasia, which may be difficult to assess from the smears alone. Many pathologists and clinical haematologists to whom we have shown our preparations have expressed surprise that such valuable information can be obtained so simply; since the technique for preparing satisfactory marrow sections from sternal-puncture material does not appear to be widely practised, this paper purposes to give a short account of the method adopted, after many trials, in our laboratories.

Method

After adequate local anaesthesia has been obtained, and with due aseptic precautions, marrow is aspirated from the sternum through a needle of the Salah type into a 10-ml. Record or all-glass syringe, which must have a tightly fitting piston. Strong suction should be applied when the marrow cavity is entered, in order to aspirate actual fragments of the tissue and to avoid undue admixture with blood, which tends to spoil the films; no more than 0.5-1.0 ml. need be aspirated. Two or three dozen or more thin films are made by spreading a minute drop of the marrow on microscope slides without attempting to select marrow fragments, and the remainder of the marrow is ejected into a 500-ml. Erlenmeyer flask containing the modified Zenker-formol which we have found to give the best results. This is prepared immediately before use according to the following formula:

Zenker's fluid (minus acetic acid)	45 ml.
40% formaldehyde solution	5 ml.
Formol saline 10%	50 ml.

Fixation is adequate in 20 to 30 minutes, and the marrow fragments must not be left for more than 30 minutes in the fixative, otherwise too much shrinkage will occur and subsequent staining will be poor. We have experimented with numerous fixatives and varied the length of fixation, but the best results were

obtained by the use of this half-strength Zenker-formol solution. When the marrow has been in the fixative for the above time 400 ml. of distilled water is added and thoroughly mixed; the tiny marrow fragments quickly settle to the bottom of the flask. The supernatant fluid is then siphoned off without loss of marrow fragments, using a Pasteur pipette with upturned point and a water pump. When the fixative is diluted the blood proteins precipitated remain as a flocculent suspension and are removed in the supernatant fluid. A further 100 ml. of distilled water is then added to the flask, which is now agitated gently a few times and allowed to stand until the marrow fragments have again settled to the bottom; the supernatant fluid is then siphoned off. A final washing in distilled water is carried out, and after standing for about 10 minutes the water is siphoned off and 100 ml. of 60% spirit is added.

The marrow should not be left for more than 15 minutes in 60% spirit, as maceration will occur; but this step is useful as it allows practically all of the protein precipitate to be discarded, and only marrow fragments should now remain. The next step is to pass the material through a series of graded alcohols in the same way—that is, siphon off the supernatant fluid and add 20 ml. of the appropriate strength, 70% spirit for 15 minutes, followed by 80% then 96% for 15 minutes each before final dehydration in absolute alcohol, two changes for 15 minutes each. The absolute alcohol is replaced by toluol and left for 10 to 20 minutes until cleared; the fragments are then transferred by a pipette to a small widely conical vessel with a flat bottom. The toluol is pipetted off, replaced by molten paraffin wax (melting-point 54° C.), and the vessel is then put in the paraffin oven for 2 hours, the molten paraffin wax being changed twice during this time. The marrow fragments, after the final change of paraffin, are allowed to settle by gravity and so gather into the small base; consequently if the wax is now hardened *in situ* the block for the microtome has the important material within a small compass. For occasional use a 2-inch (5-cm.) porcelain or silica crucible may suffice, but difficulty may be experienced in removing the paraffin block. Usually the block is removed from the vessel by running a hot knife round the edge and then rapping the crucible smartly on the bench. It must not be heated to melt the paraffin, as the marrow fragments lie directly on the flat base and are easily spoiled by overheating. For regular use we have found it convenient to employ a brass crucible with a detachable apex, so that by unscrewing the apex the block is exposed and may be easily removed. The marrow fragments lie close to the paraffin surface and are ready for immediate cutting; trimming of the block should be avoided in case the minute fragments are lost.

At the 70% spirit stage the specimen may be safely kept for a day or two, and, in fact, may be sent to a laboratory by post if the necessary apparatus for embedding and sectioning is not available. For economy the alcohols are used repeatedly by filtering back into stock bottles through fine filter-paper to remove any marrow fragments.

Sections are stained by the ordinary methods such as haematoxylin and eosin for rapid reports, but a more detailed

study of the marrow is made on sections stained by Maximow's eosin-azure method or by one of the Romanowsky methods. Special stains can, of course, be used; for example we have found Mallory's aniline-blue method of value in the recognition of Gaucher cells in sternal marrow. Microchemical tests for the presence of iron can be applied. Zenker-formol fixative does not prevent the application of the prussian-blue method if brief treatment with warm hydrochloric acid is used (Mallory).

Excellent histological preparations of marrow are obtained which show the general architecture, the proportion of fat and cellular tissue, and the relationship of leucopoietic and erythropoietic tissue to one another and to the supporting framework. The number and disposition of the megakaryocytes are readily assessed, and the presence of abnormal elements is more easily detected than in the smear preparations. We have found that unless large numbers of smears are examined a true impression of the marrow may not be obtained, especially in regard to the megakaryocytes, which are not usually evenly dispersed. Abnormal elements, such as myeloma cells, Gaucher cells, and possibly tumour cells, may be recognized in the sections, and confusion due to admixture of blood does not arise. In a case of aplastic anaemia the marrow sections remove all doubt as to whether the lack of haemopoietic elements in smears is due to failure to obtain a true sample of the marrow. These points are illustrated in the accompanying photomicrographs (see Plate).

Summary

For the interpretation of alterations in bone-marrow architecture the value of histological sections is emphasized as a supplement to the usual smear technique. A simple method is described for the preparation of paraffin sections from sternal-puncture biopsy performed with the ordinary type of needle. Representative photomicrographs illustrate the results obtained.

PAROTIDECTOMY: INDICATIONS AND RESULTS

BY

HAMILTON BAILEY, F.R.C.S.Eng., F.A.C.S., F.I.C.S.
F.R.S.E.

Surgeon, Royal Northern Hospital, London; General Surgeon,
Metropolitan Ear, Nose and Throat Hospital

[WITH PHOTOGRAVURE PLATE]

To most general practitioners and to many surgeons, the mere mention of complete parotidectomy (Fig. 1, Special Plate) conjures up an apparition with a hideous facial deformity. Superficial lobectomy (Fig. 2, Special Plate) is a far more desirable term than subtotal parotidectomy, but until it is generally conceded that the anatomical classics are inaccurate in their description of the facial nerve—that the facial nerve does not plunge into the main body of the gland and there divide within the parenchyma—the term lobectomy in connexion with the parotid gland is meaningless. To clear up misunderstandings concerning the relationship of the facial nerve, to beg contemporary anatomists to revise their teaching and alter their textbooks in this respect, is only one of the tasks that I have set myself. It is my ambition to dispel deep-rooted beliefs concerning parotid tumours, beliefs almost amounting to old wives' tales.

The Bogy of Facial Palsy

"Leave it alone! If you have it cut out you will probably get paralysis of the face." This was the advice given to the brother of one of my nursing sisters by a comparatively

young North of England practitioner; as a result, the patient adamantly refuses to submit to operation. Similar advice was given to another patient by a middle-aged doctor practising on the Welsh border. I mention these examples because most of my patients come from London or the Home Counties, where advice such as this seems to be the rule. Further inquiry convinces me that this attitude

FIG. A.—Ages at which mixed tumours of the parotid gland first appeared (J. McFarland's statistics).

towards operation is adopted by the majority of practitioners throughout the world. Amazingly enough, the profession does not realize that it is only a matter of time before the seventh nerve becomes implicated by the tumour, and that this is a far more frequent happening than facial palsy resulting from the surgeon's knife. Furthermore, when a neoplasm has advanced so far as to cause facial paralysis, it is unlikely that the tumour can be removed *in toto*.

Joseph McFarland,* who held the chair of pathology in Philadelphia, studied parotid tumours for thirty years. Unlike most pathologists, in addition to studying the macroscopical and microscopical characters of the specimens, he investigated the clinical histories of the patients. As a result he pronounced the recurrence rate to be much higher in the case of removal of small as opposed to larger tumours. Kennon, of Liverpool, substantiated this observation. Presumably as the outcome of these well-founded observations there started the pernicious practice of waiting until the tumour got larger. How large has never been stipulated; that has been left to the patient or his doctor to decide. In this connexion it is important to point out that clinically I myself have found it difficult to be sure that a given tumour is small. Some of these swellings feel definitely circumscribed and are movable on the deep structures, yet when they are displayed to the light of day they prove to be veritable icebergs—four-fifths being

*Joseph McFarland died in September, 1945. His daughter, Dr. Helen McFarland Woodbridge, is carrying on her father's collective study of parotid tumours at the School of Dentistry, University of Pennsylvania.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

FIG. 5.

FIG. 6.