In the *Practitioner*, 1946, 157, 391, Dr. D. M. Evans commented in a very valuable article on the benefits he has observed in skin and other diseases from the use of vitamin K analogue, and recently in "Any Questions" in the *Journal* (Jan. 11, p. 77) your expert stated that vitamin K seemed "to exercise a beneficial effect in some cases of perniosis—and this was borne out in the case I am recording where the patient said his cold feet were warmer after taking the drug. May I therefore suggest that vitamin K analogue should be given a trial in Behcet's syndrome and the result of such treatment recorded.—I am, etc.,

Paignton.

A. J. AMBROSE.

Vitamin H

SIR,—In a recent paper some preliminary observations were reported about the relationship of premature alopecia and biotin by Chavarria *et al.*¹ In Costa Rica a nutritional deficiency syndrome was studied by them during the last three years which appeared only in children and never in adults. Similar changes were described in Africa under the name of kwashiorkor² or infantile pellagra,³⁻⁸ but they were apparently considered manifestations of other deficiencies. The hair changes in Costa Rican children consisted of dryness, fragility, and loss of colour; the hair became loose and easily and painlessly pulled out, leading to calvities. Concurrently depigmentation of the hair varying from slight greying to intense whiteness occurred. The children often presented other deficiency symptoms such as: follicular keratosis, seborrhoeic-like dermatitis, occasionally diarrhoea, generalized oedema with normal content of blood proteins and hepatic cirrhosis, glossitis, and angular stomatitis. The diet of the affected children consisted chiefly of bananas and molasses.

Unless the patient died shortly after admission the ordinary hospital diet effected a cure. An addition of 0.25 mg. of biotin to the diet twice or three times a day appeared to accelerate the return of both pigment and growth of hair more quickly than the addition of thiamine, nicotinamide, and other vitamins.—I am, etc.,

London, W.1.

Z. A. LEITNER.

REFERENCES

- ¹ *J. Amer. med. Ass.*, 1946, 132, 570.
- ² *Arch. Dis. Childh.*, 1933, 8, 423.
- ³ *Ibid.*, 1937, 12, 193.
- ⁴ *Trans. roy. Soc. trop. Med. Hyg.*, 1940, 33, 389.
- ⁵ *S. Afr. J. med. Sci.*, 1943, 8, 106.
- ⁶ *Nature*, 1944, 154, 148.
- ⁷ *Arch. Intern. Med.*, 1945, 76, 63.
- ⁸ *J. Amer. med. Ass.*, 1945, 129, 12.

Pellagrous Encephalopathy

SIR,—I was interested in Dr. P. R. Graves's account (Feb. 15, p. 253) of pellagrous encephalopathy occurring among prisoners of war in Singapore. In our camp at Palembang, Sumatra, we had a similar syndrome, but had many more acute coma cases. These occurred mainly in the last three months of captivity, when the basic rice ration was 200 grammes per diem, with very little vegetable and practically no protein foodstuffs. Most of the acute cases had previously shown well-marked signs of pellagra and were all greatly emaciated. Some went from the ambulant stage into coma in the space of hours, and on occasion a man failed to report at the morning roll-call as he had in fact gone into a coma overnight in his billet. All showed quadriplegic rigidity, but I cannot remember any case which had convulsions. We were able to perform only one or two necropsies and these very much *sub rosa*. It was possible to examine the brain *in situ* only. The hemispheres showed generalized oedema, but apart from that little else was noted, and it was not practicable to open the abdomen.—I am, etc.,

Devonport.

J. P. CORCORAN.

Intravenous Procaine (Novocain)

SIR,—More is known about the effects of intravenous procaine (novocain) than is suggested in the reply to a question in the *Journal* of Feb. 15 (p. 282). Procaine has indeed quinidine-like actions on the heart which can prevent the development of auricular or ventricular fibrillation in experimental animals. Since, however, this action can only be produced on intravenous injection, and probably only by doses that cause central nervous side effects, it is not likely to be of much therapeutic value.

Procaine has also been given by intravenous infusion for the relief of pain due to burns (Gordon, R.A., *Canad. med. Ass. J.*, 1943, 49, 478), for the relief of the pain of labour (Allen, F. M., *Amer. J. Surg.*, 1945, 70, 283), and for sundry other types of pain and in the relief of pruritus (Lundy, J. S., *Clinical Anaesthesia*, Philadelphia, 1942). It has been suggested that this analgesic effect is due partly to a local action on a traumatized area, where increased capillary permeability would allow greater amounts than normal to escape from the blood, and partly to an action on the central nervous system. For this purpose procaine has been given intravenously in doses of up to 5 g. in 1½ hours.

It is important to realize that the toxicity of intravenous procaine is very closely related to the speed of injection, since the compound is rapidly detoxicated, partly by an enzyme in the blood (called "procaine esterase" by Kisch, Koster, and Strauss) and partly by a similar enzyme in the liver. The procaine is hydrolysed to para-aminobenzoic acid and diethyl-amino-ethanol, neither of which has the typical peripheral or central actions of procaine. If the rate of infusion exceeds the rate of breakdown the blood level of procaine will rise steadily, and toxic effects such as convulsions and respiratory failure will finally occur. Tachycardia usually appears when procaine is injected at rates exceeding 2 ml. per minute of a 1% solution (20 mg. per minute) and becomes marked when rates of over 5 ml. per minute (50 mg. per minute) are given. It is claimed that the convulsive action of procaine is prevented by pre-medication with barbiturates—e.g., pentobarbitone (1 gr. (65 mg.) by mouth).

The safety and value of intravenous procaine as an analgesic have not yet been adequately assessed; studies on this subject are in progress in my laboratory at the present time.—I am, etc.,

Department of Pharmacology,
Middlesex Hospital Medical School, London. W.1.

C. A. KEELE.

SIR,—With regard to the question (Feb. 15, p. 282) on intravenous novocain (procaine) dosage, it may be of interest to mention some unpublished facts on this subject. The writer has now administered procaine by vein in 48 cases, mainly for post-operative pain in place of morphine. The routine dosage has been 1 g. in 0.2% solution given over a period of 60–70 minutes. No deleterious effects have been noted and in general satisfactory analgesia has been obtained without the dangers of post-operative respiratory depression. This analgesia persists for as long as four days in some cases and its action is by no means clearly understood. Further investigations are at present being undertaken to determine effective blood levels consistent with safety and the mechanism of action of generalized procaine effects.

In addition to its analgesic effect intravenous procaine has a mild cerebral excitatory action which is beneficial in many post-operative cases. This effect may be very easily controlled by barbiturates, which have the further advantage of increasing the minimum lethal dose of procaine fourfold (Goodman and Gilman).—I am, etc.,

London, W.12.

E. J. DELORME.

Endocrine Receptors

SIR,—We have to be grateful to Dr. S. L. Simpson (Feb. 15, p. 270) for standing up in defence of endocrinologists who paid attention to the "endocrine receptors" discussed by Mr. Aleck Bourne (Jan. 18, p. 79) in view of your statement in the editorial review in the same number (p. 96).

Due tribute should be paid, however, on this occasion to Prof. Julius Bauer, of Los Angeles, California, formerly of the University of Vienna, who based his teaching of endocrinology on this assumption as long ago as 1917.¹ The piquancy of the situation is the fact that in the summary of his works, *Constitution and Disease* (Wm. Heinemann, London, 1942), he devoted a whole chapter to the discussion of this question (p. 53). The book was extensively reviewed in your leading article of April 24, 1943 (p. 510), from which I quote: "Like H. H. Dale, he [Bauer] emphasizes the importance of the receptivity of the organ to hormonal stimulus, which explains cases where the endocrine lack is expressed in only one organ." In a previous paper² Bauer explains the manifold activities of the pituitary hormones in this way.