On the other hand there are very serious reasons for disbelieving any possibility of these creatures having passed the intestine. The chemical objections have to some extent been considered by Dr. Sharpe, and he himself discounts the possibility of the soft-skinned larva resisting the protein-digesting secretions of the stomach. He therefore assumes that the eggs with their "resistant chitinous coating" can pass the stomach unharmed and that the larva can hatch from the egg in the more alkaline portion of the bowel. In actual fact the egg membrane of the *ptinids* is very fine and thin, and in the absence of experiments does not justify the assumption of its being much more impervious to pepsin and hydrochloric acid than the chitinous integument of the larva. In order to obtain conclusive evidence on this subject I have treated eggs, newly hatched larvae, and almost full-grown larvae of *Ptinus tectus*, which were kindly placed at my disposal by the Director and Mr. R. Howe of the Pest Infestation Laboratory, Department of Scientific and Industrial Research, with artificial gastric juice. The specimens were first admixed to chewed bread, and the experiment was kept at 36-38° C. for three hours. The material was then washed in distilled water and scrutinized under a dissecting microscope. Full-grown and newly hatched larvae were dead but not disintegrated, and larvae more or less ready for hatching had been killed inside the eggs. There is no evidence so far (after ten days) that any of the less developed eggs have hatched. The fact that young larvae have definitely been killed inside the intact egg-shell seems to prove at any rate that the egg-shell does not afford a considerable protection against the influences prevailing in the human intestine.

Dr. Sharpe has also stressed the mechanical difficulties for the larvae of remaining in the intestine for a period of a week or more in view of the action of the peristalsis. According to him the larva holds itself in position by its claws and jaws. In actual fact the rather thin claws would probably be of little avail, while the jaws would be wanted for the ingestion of the food, so that they could not possibly keep their grip. On the other hand this point is of minor consequence, as an insect larva of this type developing in the intestine would find a nook at the base of the villi or somewhere else where it would be relatively safe, and it would soon create a small excavation round its body in which it could rest securely.

Ptinus tectus lives in dried animal or vegetable matter, such as dried crayfish (e.g., of collections or of food for aquarium fish) and casein (in which latter it is a regular pest) on the one hand, and in rye, maize, cayenne pepper, cocoa, ship's biscuit on the other hand. It is therefore very well adapted to a life in a medium containing between 5 and 15% humidity. The larvae would, however, be quite unsuited to a habitat in a stiff semi-liquid material with a moisture content of more than 90%, as is found in the human intestine, and they would very quickly be clogged up and asphyxiated. While experimenting with artificial gastric juice I kept another batch of eggs and newly hatched and almost full-grown larvae in chewed bread in a moderate quantity of saliva without gastric juice for 31 hours at about 20° C. This treatment, too, killed the larvae, but it is too early yet to decide whether all the eggs have been killed in this experiment. Mr. Howe has informed me of some tests with much lower degrees of humidity in hygroscopic foods with an estimated water content of up to 40%. As only a small percentage survived in these materials it is obvious that a further increase of the water-content would soon prove lethal, as shown in my own experiment.

The gravest reason, however, for scepticism is afforded by the question of temperature. It is known from experiments with eggs of the flour-moth (*Ephestia kuehniella*, Zell.), an insect which lives under very similar ecological conditions, that they cannot hatch but will perish at a prolonged exposure to a temperature of 34.4° C. or more. Even in a moist cell it has not been possible to raise this temperature (Hase, A., 1927, *Arb. biol. Reichsanst. Land u. Forst-wirtsch.*, 15, 116). As to *Ptinus tectus* itself Mr. R. Howe informs me from his experience, gained at the Pest Infestation Laboratory, that only large larvae survive for long at 28° C. or above, and that only few first-stage larvae reach the second stage at all at as moderate a temperature as 29° C. The most conclusive point against Dr. Sharpe's view, however, is the fact that, according to Mr. Howe's experiments, eggs do not hatch above 28° C., so they would hardly remain alive for a day and certainly never hatch at the temperatures prevailing in the human intestine.

Lastly a point of terminology should not pass without comment. The term myiasis, derived from *μύια*, a fly, designates the "infestation of any part of the body of living animals by the larvae of diptera," or two-winged flies (Smart, J., 1943, *A Handbook for the Identification of Insects of Medical Importance*, p. 810, London: British Museum). A similar infestation by beetles or their larvae is called cantharidiasis. Entomologists have for some time been aware of the fact that many reports of myiasis and similar ailments collapse on closer examination, and modern textbooks like that of Smart duly stress the necessity

for a thoroughly critical examination of each reported case. My personal impression is that in temperate European countries the larva of the lesser house-fly is most likely to produce genuine cases of myiasis, and that certain tyroglyphid mites can probably produce similar and sometimes rather serious symptoms.

I would like to express my thanks to the Director and Mr. R. Howe, Pest Infestation Laboratory, Department of Scientific and Industrial Research, Slough, Bucks., for breeding material of *Ptinus tectus* and for the information cited above; to Mr. N. D. Riley, Keeper of the Department of Entomology, British Museum (Natural History), and Mr. F. Barnett of the same Department for their, unfortunately unsuccessful, efforts at obtaining a quantity of pepsin for me; and, lastly, to the John Innes Horticultural Institution for providing me with this substance.

—I am, etc.,

Imperial Institute of Entomology.

F. I. VAN EMDEN.

SIR,—I was most interested to read Dr. D. S. Sharpe's account of a case of intestinal myiasis caused by the larva of *Ptinus tectus* (Jan. 11, p. 54). I am, however, startled by his implication that the larva of *Ptinus tectus* is insectivorous, and I should be glad to know which insects normally form the food for this larva.

I feel also that Dr. Sharpe may be adding to the anxieties of parents by his lurid threat of intestinal myiasis to those children who eat fallen fruits. The only dipterous families which are likely to infest fruits attractive to even the most omnivorous (or insectivorous?) child are the Trypetidae or the Drosophilidae, and, so far as I am aware, no member of these families has so far been incriminated with causing myiasis. The adults of the Muscidae and the Calliphoridae—the myiasis-causing species *par excellence*—certainly feed on fermenting fruit, but I think they prefer a rather more decayed pabulum for breeding purposes.—I am, etc.,

Lagos, Nigeria,

P. J. L. ROCHE.

Control of Measles

SIR,—After a depressing week-end of Arctic weather, snow shifting, and fuel crises (national and domestic) it was a welcome relief to open the *British Medical Journal* and read the three leading articles (Feb. 8, p. 225). The first of these ably reviewed the present position with regard to the control of measles, a very topical subject; the second set out clearly the reasons why a severe outbreak of influenza is unlikely this winter; while the third had the significant title "The Minister Accepts."

I found myself in general agreement with the opinions expressed in the article on the control of measles, but would like to make one or two comments with regard to the availability and use of convalescent measles serum, normal adult serum, and gamma-globulin. The last is at present unobtainable, while convalescent measles serum is in my experience a very rare product indeed; but adult normal serum could and should be readily available for the use of any doctor requiring it.

From the beginning of this year Leeds doctors have been able to obtain adult measles serum through the Public Health Department. The supply is obtained by arrangement with the Regional Blood Transfusion Service, who kindly carry out the necessary bottling and undertake the essential tests. The few weeks that have elapsed since the inauguration of this scheme have already proved its value, and in addition to numerous inquiries from doctors within the city there have been several requests for serum from areas outside the city and two from as far afield as Newcastle, so that it would appear that as far as the north-east of England is concerned the facilities for obtaining this product either do not exist or are not sufficiently well known. In this latter respect the statement in the leading article that normal adult serum is available from most of the constituent laboratories of the Public Health Laboratory Service was news to me, and I am sure it would be helpful if further publicity could be given to this source of supply.

I have found that some confusion exists as to the use of adult serum for attenuation. In one or two instances the serum has been given to the case rather than to the contact. Claims have been made as to the value of convalescent and adult serum in the treatment of a case of measles, but these have not been substantiated. When using adult serum for prevention the usual dose recommended is 10 ml., and this should be given to the

contact not later than the day of the appearance of the rash in the infecting case. If used for attenuation the same dose of 10 ml. should be given to the contact on the 4th, 5th, or 6th day after the appearance of the rash in the infecting case.—I am, etc.,

Leeds.

J. F. WARIN.

National Health Service Act

SIR,—It is with diffidence that I enter the controversy on the National Health Service Act. It would appear that everything has already been said many times over on either side. However, it seems to me that the minority who voted "Yes"—a minority that was nearly 50%—should be more often represented in your columns. I particularly disagree with the claim made by so many of your contributors that opposition to the Act is not based on politics, or, at any rate, on party politics. Personally I am not a Socialist and have never voted for any party, and, considering the complete political unreality of so much of the argument, it is obvious that most medical men are not actively engaged in political life. Nevertheless, how many are there of those opposing the Act who do not cast their vote for a Conservative at an election? Professionally and socially we are predominantly Conservative and are naturally suspicious of a Labour Government and antagonistic towards its efforts. The people of this country, however, elected that Government with an overwhelming majority in Parliament. It is idle to pretend there was not a widespread determination to obtain social security, and votes were given to a party that could be trusted sincerely to support it. That was the working of democracy. Is it democratic for a section of the community to oppose the wishes of the majority as expressed in an Act of Parliament? Is it not dangerous to suggest that an organized professional body should try to make an Act unworkable in order to change it? Are we going to support miners or transport workers who might try to do likewise?

The Act sets out to effect a revolution—that no longer will poverty and lack of means be a bar to obtaining the best medical facilities and treatment available. Revolutions always disturb established customs. Such a great social advance cannot be made without some risks and disadvantages. Let us remember the advantages it will gain for our patients and potential patients, and be particularly careful that we are sincere when we claim to oppose the Act on behalf of our patients. When we complain because in time of scarcity extra amounts of rationed tit-bits are refused to a dying man who can afford them, it is well also to remember those who have been condemned to death or persistent ill-health through inability to afford necessary medical treatment.

It is for the individual doctor eventually to decide whether to enter the Service or not, but the recent plebiscite has finally destroyed the claim for unity within the profession. A large proportion of doctors, especially the younger ones, are willing to build up the best possible medical service on the present Act. Alterations in medical practice are part of wider social and economic changes hastened on by the war and are inevitable. The inherent danger of bureaucracy can be defeated by the efforts of those determined to make the Service successful. It is therefore to be hoped that in the coming negotiations the profession's representatives will display the utmost statesmanship and realism. Could the Minister be blamed for having failed to discuss previously the Act with a body so out of step with the times that it opposed a medical service covering 100% of the population? The Act is now on the Statute Book and we have our last chance to offer expert advice in clothing its bare bones.—I am, etc.,

Tunbridge Wells, Kent.

D. G. ff. EDWARD.

Freedom of Choice

SIR,—We are most of us concerned with the imminent prospect of State control and of course hold freedom of choice of doctor by patient to be essential. Just how much freedom of choice the patient will have may be surmised from the following recent example. Mr. A is a non-industrial Civil Servant. He is an insured person and is under treatment by me for severe thrombo-phlebitis of leg. A few days ago he received from his department—namely, the Assistance Board—a letter stating

that his case had been referred by them to the Treasury medical adviser—the object of this being to enable remedial action to be taken.

The Treasury medical adviser directs the Assistance Board to obtain an independent medical report from Dr. B, who is a general practitioner in the same area as myself. Since Mr. A did not chance to select Dr. B for his own doctor, however, he is treated with scant courtesy by Dr. B, who, when informed of the complaint, refuses to visit him and insists upon his turning out and walking to the surgery with a thrombosed leg on dangerously ice-bound and snow-covered roads. Mr. A is then informed that he will probably have to attend again for a further report. All this happens without any notification to me as Mr. A's doctor or any report on his condition being requested.

To comment on this it may be summarized by saying that Mr. A, while under treatment by a doctor he wishes to attend him, is suddenly directed by the State as his employer to visit Dr. B, whom he does not wish to attend, so that the Treasury medical adviser, who has never seen him, may decide what treatment should be given. All this takes place in a most underhand way completely behind the back of his own doctor, who is neither consulted nor informed of what is going on. Now Mr. A cannot be simultaneously treated by two persons, so pursuing this to its logical conclusion either the State removes him from his own doctor, or, if he elects to continue treatment from his own doctor, it may threaten him with withdrawal of his salary. The most disturbing feature is that this is apparently standard procedure, the letter to Mr. A being a cyclostyled copy bearing the form number G46/200/779.—I am, etc.,

Greenfield, Lancs.

R. S. HOLGATE.

Administrators of the Act

SIR,—There is one seldom mentioned side of the National Health Service which I hope our negotiators will not overlook: the method of appointing the whole-time medical administrators. Whatever the final form of the Service it is clear that considerable executive authority will be vested in the statutory councils, boards, and committees envisaged in the Act. They, however, will consist for the most part of part-time members, and therefore considerable discretion and authority must devolve on their senior permanent officials, who will include medical men. It is our concern to see that from the start the method of appointing these key officials avoids the worst features of bureaucracy.

In my view at least two conditions must be fulfilled: (1) All senior appointments must be advertised and so thrown open to the widest possible "field." (2) Selection to medical appointments must be made by bodies on which there are representative medical members. This could be done peripherally by the body for whom the candidate is going to work or centrally by the Central Health Services Council. Any system which makes these appointments plums with which to reward long and faithful service in the existing medical administrative departments will, I am certain, prejudice fatally the prospect of getting a smoothly running service. Moreover I fear that unless we take steps to ensure the contrary this is the method of appointment that is likely to be applied.—I am, etc.,

Plymouth.

T. A. A. HUNTER.

Remuneration under the Act

SIR,—Now that discussions have begun between the Negotiating Committee and the Minister may I draw attention to a fundamental point hitherto overlooked? It is the method of remuneration by capitation fee—surely one of the worst evils of the N.H.I. scheme and now, alas, threatening to be an even bigger evil in the future comprehensive Health Service. I think I am right in saying that the majority of doctors, like other members of the community, prefer to be remunerated in accordance with the amount of work they do. Conversely they do not wish to be paid for anything they have not done. Nor is this feeling confined to the doctors; it is shared by patients, many of whom feel, when they need attention, that they have to apologize to their panel doctor for bothering him. How different many patients would feel if they knew their doctors were being paid on a *pro rata* basis for their time and trouble. As an example, with the best intentions in the world, human