Correspondence

The Toxicity of Iron Compounds

SIR,—The paper by Dr. Gilbert Forbes (March 22, p. 367) has drawn attention to the toxic action resulting from accidental overdosage of ferrous sulphate taken by mouth. No mention is made of the name of the preparation, but we believe we are correct in saying that it was "fersolate" tablets, of our manufacture.

Experiments in progress in these laboratories indicate that for the more soluble iron compounds the toxicity is closely correlated with the iron content (calculated as ferrous iron) and strongly suggest that the action is due to the formation of ferric chloride in the stomach. The iron compounds were given, either in solution or suspension, to fasting rabbits by means of a stomach tube; the doses were adjusted to body weight. According to the dose, the animals either died in a few hours or recovered completely. The post-mortem findings for those that died were completely in agreement with those of Forbes, the cause of death being haemorrhagic gastritis. The animals that recovered were quite well on the following day; daily blood counts and urinary examination showed no abnormal findings. In some cases, but not all, the stomachs of these animals showed signs of irritation at the great curvature.

The results so far obtained indicate that the fatal dose of ferrous sulphate in rabbits is higher than those reported by Forbes for guinea-pigs. We find that "fersolate" tablets compare favourably for toxicity with other iron compounds so far tested, and the presence of small amounts of copper and manganese sulphates does not increase the toxicity, which is due to ferrous sulphate. The toxic effect of the iron in Bland's pills is less than half that of other ferrous salts; but relatively higher doses are required for haemopoietic action.

The lower toxicity of Bland's pills may be due to the sodium sulphate in this preparation having a cathartic action that hastens the excretion of the iron. Rabbits treated with high doses of Bland's pills had a profuse diarrhoea within half an hour of dosing. It would appear, therefore, that a gastric lavage with an alkaline solution would be of value after accidental overdosage with ferrous sulphate. This would neutralize the acidity of the stomach, cause precipitation of ferrous carbonate, and prevent the formation of ferric chloride and retard the absorption of iron.

The results of further experiments, now in progress, will, it is hoped, be published in detail later.

Ferrous sulphate is one of many medicinal substances having undesirable effects when taken in grossly excessive quantities. We consider that Dr. Forbes's paper serves to emphasize the need to keep such preparations out of reach of young children, and to guard against overdosage.—I am, etc.,

Glaxo Laboratories, Ltd.,
Greenford, Middlesex.

G. F. SOMERS.

SIR,—Dr. Gilbert Forbes (March 22, p. 367) has written a very important paper which has wide-reaching implications. Have we not been assuming too much that "harmless" drugs are innocuous even in large doses, and that the excess will be excreted without danger? The outstanding example is the vitamins which are found only in traces in natural products, and yet are prescribed medicinally in enormous quantities.

To return to the ferrous sulphate question, I remember some 12 years ago being stimulated by Dr. Helen Mackay's papers on anaemia in infants to order massive doses of ferrous sulphate as a routine procedure during the last months of pregnancy. After some months I abandoned this, as a number of women complained of headache, nausea, and malaise, and, if the iron were continued, developed a mild albuminuria. I was convinced that there was an association between these liver and kidney symptoms, resembling a pregnancy toxæmia, and the excessive iron intake, but I must acknowledge that I did not find anyone to agree with me. However, I have never given large quantities of iron in pregnancy since, and in the light of Dr. Forbes's cases

with the reported damage to liver and kidneys I venture to bring up this theory once more and to ask whether anyone else has had a similar experience?—I am, etc.,

London, W.8.

VIOLET RUSSELL.

Retrodisplacement and Fertility

SIR,—There is not the slightest doubt that retrodisplacement of the uterus is a cause of sterility. A woman with a backward displacement of the uterus *can* become pregnant, but if she is sterile and a backward displacement is present, the correction is followed by pregnancy in a very large number of cases. Needless to say, full investigation of both male and female is a necessity.—I am, etc.,

Dublin.

BETHEL SOLOMONS.

The "Costoclavicular Syndrome"

SIR,—Prof. E. D. Telford and Mr. S. Mottershead raise in their paper (March 15, p. 325) several interesting points and advance new concepts of great importance to a proper understanding of the various factors involved in cases of brachial neuralgia. Probably owing to the paper shortage clarity has been sacrificed to compression in certain places.

The authors describe the operative findings in fourteen cases associated with cervical rib and extensive arterial thrombosis; except in two cases the thrombosis extended no higher than the lower border of the pectoralis major. Are these findings meant to imply that the cervical rib played no part in the syndrome and that no case of arterial damage due to cervical rib was found in this extensive series? Is it believed that the arterial damage was due to compression between the heads of the median nerve and that this was due, either directly or indirectly, to the fact that the cervical rib altered the relationship between the plexus and artery?

The stated "textbook picture of neurovascular bundle compression" fails to convey what is found in practice. In our experience there is no such picture. These cases differ considerably in history, symptomatology, and signs, and this difference has valuable diagnostic implications. The authors are undoubtedly right in casting doubt as to whether their remaining 69 cases of cervical rib were in fact correctly described. Prof. J. R. Learmonth's recent paper in *Thorax* undoubtedly gives a balanced judgment of this problem.

We believe that the case in which the axillary artery was compressed by the two heads of the median nerve is the first to be described and as such is of considerable importance. It might be helpful to consider this concept from its obverse angle. Consideration of the respective resistances of artery and nerve would lead to the expectation that symptoms would arise from the nerve rather than the artery in such a case. From the rather meagre clinical data provided it appears likely that this in fact happened. It might be preferable to contend that the median nerve was compressed by the artery. In fact, quite possibly this man's symptoms might have been alleviated by resection of his first thoracic rib, as this would have led to alteration of the artery-plexus relationship.

The challenging statement that in postural manoeuvres of the shoulder resulting in obliteration of the radial pulse the axillary pulse can still be felt cannot be allowed to pass unquestioned. A small personal experience does not suggest its general validity and it is contrary to certain published evidence. It is not clear from the text or the table whether the authors in fact ascertained, in each case where the radial pulse was obliterated, that the axillary pulse was still present and undiminished. Further information on this point would be welcome and, in particular, if in fact all cases had not been examined, on which cases this statement was founded. Further, the statements in the summary, paragraphs 2 and 4, are contradictory; paragraph 4 is almost certainly a misprint, as the references quoted include several undoubted cases of costoclavicular compression proved at operation. The role of costoclavicular compression in causing a rise in venous pressure in the arm requires consideration (Sampson *et al.*, 1940). The evidence as a whole is suggestive that costoclavicular compression is common and cannot be disregarded. It is noteworthy that the rise of venous pressure on bracing back the shoulders cannot be attributed to median nerve compression. It is considerably lessened if the first rib is missing as in thoracoplasty. In these cases the average rise

in pressure was 2.3 cm. of water, while the rise in 26 normals was 6.7 cm. of water.

The section dealing with the anatomical dissections is interesting and informative but, in a way, unsatisfactory. A considerable amount of work has not been utilized to its full extent. Nerves are described as being "tightly stretched"—tension can be measured; "the clavicle moves from the rib"—distance and angles can be determined. Kelvin, throughout his life, stated a great though limited truth: "Science is measurement." There is no doubt that the rational assessment of anatomical problems is profoundly aided by the discerning analysis of even necessarily imperfect measurements. That this problem is not easy is appreciated, since we have been interested in its complexities and difficulties for some time. Telford and Mottershead have undoubtedly done a first-class piece of work, and this is properly acknowledged in your leading article, but their answers to some of the questions we pose would also be of great interest.—We are, etc.,

London, W.1.

ERNEST FLETCHER.
ANTHONY FLETCHER

REFERENCE

Sampson, J. J., Sanders, J. B., and Capp, C. S. (1940). *Amer. Heart J.*, **19**, 292.

Curare

SIR,—Dr. Massey Dawkins (Jan. 18, p. 111) has drawn attention to a condition of shock, as reported to him by nurses and ward sisters, among patients who have received curare, which, he says, is probably due to insufficient anaesthesia. The letters from Dr. A. J. Gray (Feb. 1, p. 196), Dr. H. G. Kahlenberg (Feb. 8, p. 234), and Dr. G. V. S. Wright (Feb. 15, p. 268) point to this condition of shock as being due to the profound muscular relaxation and anoxaemia from depression of respiration. Dr. D. S. Jones (Feb. 15, p. 269) suggests that post-anaesthetic complications may be due to an overdose of curare. From this one infers that the condition of shock results from, overdose of curare.

From these three theories surely far the most plausible is the second as proposed by Drs. Gray, Kahlenberg, and Wright. No patient should ever be allowed to return to the ward in such a condition, no matter what anaesthetic has been used. Precautions must be taken in the theatre during and at the close of the operation to counteract it, otherwise such excellent drugs as cyclopropane and curare become a veritable menace and fall into undeserved disrepute.

Certain points in the induction, maintenance, and recovery from anaesthesia where curare has been used seem pertinent. Where maintenance is to be by "pentothal" and endotracheal oxygen it seems important to administer *enough* curare to obviate laryngeal spasm and straining at the commencement of the operation; too little may lead to difficulties in intubation, with subsequent misuse of the drug's potent qualities and an upsetting of technique. Controlled respiration is then performed, and as soon as automatic respiration is resumed the partial rebreathing technique is resorted to, oxygen being the only gas inhaled. The soda-lime is then used in accordance with the desirability or not of quiet breathing. In this way the carbon dioxide in the circuit causes dissociation of oxygen from oxyhaemoglobin in the tissues. Tissue asphyxia and hypotonicity with diminished venous return are mainly due to depressed respiration leading to anoxaemia. This I submit is most often caused nowadays not from overdose of drugs but from the use of prolonged carbon dioxide absorption and endotracheal tubes.

Except in the case of certain operations, among which I would include those demanding the full Trendelenburg, the use of an endotracheal tube certainly predisposes to diminished respiratory excursion from the necessity of having to use more curare. It seems therefore wiser not to use an endotracheal tube as a general rule, thus cutting down the amount of curare required, using only sufficient to procure adequate muscular relaxation, and to use only minimal doses of "pentothal." An endotracheal tube is not normally necessary provided an adequate airway is ensured, as in any case curare counteracts reflex laryngeal spasm even though a muscle-relaxing dose might not be sufficient to allow direct laryngeal intubation.

During maintenance the soda-lime should be cut out of circuit and the tissues given the benefit of carbon dioxide either self-generated in a cyclopropane anaesthesia or from a cylinder with the expiratory valve open in a "pentothal"—oxygen administration. With cyclopropane maintenance only sufficient curare is used to procure relaxation, and the soda-lime is periodically cut out whenever the respiratory excursion can be deepened. Air is admitted to the circuit before the close of the operation and further amounts of carbon dioxide used to help eliminate the anaesthetic. If respiration is still depressed there should be no hesitation in giving physio-

stigmine and atropine intravenously before returning the patient to the ward.

Finally in cases where respiration is markedly depressed or where the blood pressure is not within 10 mm. Hg of the pre-operative level and where the pulse pressure is below 30, the usual clinical evidence of shock being present, intravenous plasma or adrenaline drip should be instituted without delay. Coramine and oxygen are prescribed and house-surgeons warned to withhold the use of morphine until the patient has fully recovered and it is certain that any restlessness is due to pain. For less serious degrees of shock, warmth and raising the foot of the bed and a rectal saline drip are measures to be instituted. As soon as possible the patient should be encouraged to sit up and moved from side to side and remedial chest exercises instituted, as pulmonary complications are prone to develop following upon such a condition of shock.

I have been interested to find that in one or two cases this condition of shock has been associated with a marked *diminution* in pulse rate to the region of 50 beats a minute. This is a common finding in cyclopropane overdose and may possibly be an indication of "pentothal" overdose as well, though I have not been conscious of administering an overdose in this respect. Again I can only surmise that the condition is due mainly to anoxaemia or oxygen lack in the tissues, occasioned by diminished respiratory exchange from too much carbon dioxide absorption, leading to a diminution in peripheral resistance. Coramine acts centrally so must be reinforced with an additional supply of oxygen. I would be interested to hear whether other anaesthetists have also experienced this state of affairs.

—I am, etc.,

Bristol.

W. M. MAIDLOW.

Clinical Freedom and the Board of Trade

SIR,—The Minister of Health recently declared that "it is a basic principle of the new Service that there should be no interference with the clinical freedom of any doctor—specialist or general practitioner." Why then has the Board of Trade this week issued an instruction as to the type of case for which a spinal support may be prescribed (*Supplement*, March 15, p. 33)? Thirty-nine diseases and deformities are listed, ranging from actinomycosis of the spine to ureterostomy. Apparently there will be no problem in securing a surgical appliance for spinal neurosis! Many of us may think that no treatment could be worse, but at least the Board of Trade has no objection. It is also gratifying to learn that a surgeon will have no difficulty when he makes a diagnosis of actinomycosis of the spine (of which I know of only six records in the literature of the world); a simple prescription will suffice. But for the scores of patients seen week by week and month by month with prolapse of an intervertebral disk, such a prescription will not be accepted. The surgeon must prepare "a fuller medical certificate," give "the precise condition," add "sufficient detail," send it to Horseferry Road, where the Board of Trade will examine it, and submit it to the medical advisers of the Ministry of Health, who in their wisdom will deliberate and in due course pass judgment, and then send it back to the Board of Trade for endorsement, who will return it to the surgeon and possibly grant permission for him to refer it to his instrument-maker. Does anyone believe that such a proposal is calculated to promote honesty in the minds of surgeons, or is it obvious that sooner or later the alternative of subterfuge will be adopted and, no matter what the true diagnosis may be, a suitable label will be found in one of the 39 listed conditions which is less of an untruth than others and the prescription made accordingly?

In short I do not complain that one of the most common of all spinal disorders has been left off the list; I complain that such a list has been made at all. I complain of legislation of such a type as to bring legislation into contempt. The object could have been achieved more successfully by reminding doctors that certain materials are still in short supply and that care is still needed in prescribing spinal supports and corsets. A simple letter of 20 words, conforming to the traditions of the profession, would have done much more than this blundering 500-word document with its central direction and civil service control.—I am, etc.,

London, W.1.

REGINALD WATSON-JONES.

Certificates for Corsets

SIR,—With laudable objective in mind brewers allocated a proportion of available supply of spirits for medicinal use, supplying and designating same as such and providing such in