and also in primary suppurative meningitis due to the pneumococcus and many due to *H. influenzae* (Pfeiffer) is not questioned, but in meningococcal cases its superiority to oral and systemic sulphonamides has not been established, and its use is attended with unjustifiable risks.—I am, etc.,

London, S.E.13.

L. J. M. LAURENT.

Epidemic Kerato-conjunctivitis

SIR,—The annotation on epidemic kerato-conjunctivitis (Jan. 25, p. 145) prompts me to report the common occurrence of K.P. in an outbreak among British personnel in Calcutta in the rainy season of 1945.

The usual history was of a sore eye 10 days after trauma occasioned by Rugby, soccer, or motor-cycling without goggles. Swelling of the upper lid and of the pre-auricular gland was present. Staining of the cornea with fluorescein developed a few days later, followed by K.P. in 20 out of 30 cases, sometimes with folds of Descemet's membrane and sometimes after the ciliary flush had subsided. The corneal picture bore a marked resemblance to the eye lesions of onchocerciasis, which, of course, was not involved. The K.P. was as often associated with the type of fine corneal lesion as with the disciform type, and was quite a feature in this epidemic.—I am, etc.,

Johannesburg.

J. GRAHAM SCOTT.

Endocrine Receptors in Relation to Cancer

SIR,—There can be little doubt that a living cell is ready and willing to receive an endocrine stimulus just as in plants and animals the female is ready and willing to receive the male element. But I fail to see why Mr. T. C. Clare (March 22, p. 390) labels a cancer cell as degenerate and therefore lacking in an endocrine receptor. A cancer cell is not degenerate by a very long way—it is, unfortunately, young, active, and vigorous.

For many years there has been an idea that cancer arises from some fault in the endocrines, the cell receptors, or both, but prolonged observation and clinical experiment have failed to substantiate this theory, promising though it appeared to be.

We know that cellular division is the result of nuclear activity—the greater the nuclear activity, the more rapid the cell division. And here we must hark back to the very beginnings of the organism. An organism was originally a small bit of our planet cast off when the latter began to cool. We are of the earth, earthy. In the course of long years of evolution it adapted itself to surrounding influences, and ultimately man appeared. But just as in our parent, the earth, there are some atoms, such as uranium, which are not too stable, so man may have some cells of a like character—liable to burst into exaggerated nuclear activity at slight provocation. Lucky are the people who have not inherited these unstable cells!

I submit, then, that cancer is brought about by irritation whipping up the electrical phenomena known as nuclear activity in cells which happen to be unstable. If this be so, what then? The Ministry of Agriculture has forbidden the propagation of plants which are liable to disease—for instance, some varieties of potato. Breeders of animals and birds choose healthy parents. As this naturally is impossible in the human race, the predisposition to cancer will continue to be handed down.—I am, etc.,

Wigan.

J. THOMSON SHIRLAW.

Amoebic Hepatitis

SIR,—During the last few months there have been admitted to my wards four cases of hitherto unsuspected amoebic hepatitis with abscess, two of them fatal. One gave a history of treatment with emetine in 1930 in India, and two others admitted to fleeting attacks of diarrhoea in the Far East, while the fourth assured me that he was the only soldier in the Middle East who had never had an attack of diarrhoea. A fifth case sent in for diagnosis for axillary glands was found to have a tender liver, and the sigmoidoscope revealed advanced amoebic ulceration of the bowel. This man had been in the Navy and had had attacks of diarrhoea on and off, but never enough to report sick.

As such a large part of the British Army served in "amoebic dysentery countries" in the late war, and as the original infection by the *Entamoeba histolytica* can be such a mild disease, it seems to me that such cases may become common during

the next few years, and practitioners should keep a careful watch for them. Admittedly the signs and symptoms of early amoebiasis of the liver are indefinite and are not confirmable by laboratory tests. In a doubtful case, however, the improvement with emetine is dramatic if one is on the right track. Few diseases are more treatable in their early stages, but if not diagnosed in time disaster may easily follow.—I am, etc.,

Bristol.

RICHARD C. CLARKE.

Arsenical Encephalopathy

SIR,—The case report by Drs. G. Hipps and R. Goldberg (March 8, p. 296) on a patient dying with arsenical encephalopathy calls for some comment. Their treatment was that advocated by Ransome *et al.* (1945)—i.e., the patient is sat up in a vertical position, receives calcium gluconate, calcium thiosulphate, vitamin C, intravenous glucose-saline, and is repeatedly lumbar-punctured. Stress is laid on the postural part of therapy in order to decompress the brain.

Whereas it is obviously difficult to be dogmatic in a condition which is rare and where spontaneous recoveries occur (mortality is estimated at 75%), the promise of British Anti-Lewisite (dubbed BAL by the Americans) appears to be considerable in reducing the mortality in arsenical encephalopathy. In a recent survey by the American Council on Pharmacy (1946) 44 out of 55 patients suffering from arsenical encephalopathy recovered after receiving BAL; of the 11 cases who died 5 were started on this treatment some 9–72 hours after coma had set in. Thus with early administration of BAL one may hope to reduce the mortality to less than 15%. The American Council recommends 3 mg./kg. four-hourly intramuscularly for two days, then four injections on the third day and two injections on the following days until the patient has completely recovered. They emphasize that BAL must be given early and in adequate concentration to be of use.

My own experience is confined to two patients, both of whom recovered with BAL as the principal therapeutic agent, although both were comatose when this treatment was begun. Incidentally the first patient was nursed flat on his back, the second sitting up. In view of the published results of the American Council it is considered that BAL should be the primary agent in treating patients with arsenical encephalopathy.—I am, etc.,

G. W. CSOKA.

REFERENCES

- Ransome, G. A., Paterson, J. C. S., and Gupta, L. M. (1945). *British Medical Journal*, 1, 659.
Report of the American Council on Pharmacy (1946). *J. Amer. med. Ass.*, 131, 824.

Electric Convulsion Therapy

SIR,—During the past nine months a new technique for the administration of oxygen and CO₂ throughout the convulsion has been used in the out-patient E.C.T. clinic at St. George's Hospital in an attempt to reduce the excessive cyanosis produced by the convulsion, with the resulting stress upon the cardiovascular system..

It has been our practice over many years to administer oxygen through a Hewett N₂O and O₂ mask at the conclusion of the convulsion, and in this way to cut short the period of cyanosis as much as possible. However, it was felt in the case of old patients (a number being over 70 years of age), and in plethoric and hyperpictic subjects and in those suffering a degree of cardiac insufficiency, that the less strain that could be placed upon the cardiovascular system the better. With this object in view, it was decided to administer oxygen throughout the convulsion. This at first sight may appear illogical in view of the supposed laryngeal spasm occurring at that time; in actual fact, however, this spasm seems to take on the characteristics of the general clonic contractions and in reality is only intermittent—this can be observed by watching the bag of the apparatus when oxygen is administered throughout the convulsion; it will be noticed that very shallow and rapid respirations are taking place in time with the convulsions.

That gaseous exchange is also taking place is demonstrated by the fact that at the end of the clonic phase, and before normal deep respirations are resumed, the patient will be seen to be a good colour with little or no evidence of cyanosis—this being a marked contrast to those cases which do not receive oxygen during the clonic stage.