The technique observed is as follows: The current is administered in the normal manner; after the original muscular contraction and cry it is rapidly verified that the mouth gag is in correct position, has not shifted, and is firmly gripped. A Hewett gas-and-oxygen mask is then immediately placed over the mouth and nose and kept firmly in position; the mask should be of sufficient size to cover the No. 9 size rubber gynaecological ring used as a mouth gag. The oxygen is turned on and the administration is continued throughout the clonic phase and until the patient has taken two or three normal respirations.

With this technique cyanosis is avoided, thereby reducing cardiovascular stress and allowing a wider margin of safety; at the same time the impression gained by this observer is that there is a reduction in distress experienced by the patients on recovery from the convulsion.

I should like to thank Sister B. N. Samuel for her efficient running of the E.C.T. clinic.—I am, etc.,

London, S.W.1.

S. M. WHITTERIDGE.

Control and Treatment of Diphtheria

SIR,—Dr. James Fanning's interesting account of an outbreak of diphtheria in a girls' school and your excellent leading article on the control and treatment of this infection (March 22, pp. 371 and 384) both draw attention to the necessity for boosting doses of prophylactic at regular intervals during childhood and the school years. Insufficient attention appears to have been paid to this latter aspect of diphtheria control by the propaganda departments of public health authorities, and when this weakness in a child's protective armour is pointed out one is frequently confronted with the parental argument that their 7-year-old is immune because he was injected at the age of 1 year.

There is little difficulty in obtaining permission to revaccinate the school child, but the public still requires considerable education before it will accept the idea of reimmunization to diphtheria, and it is hoped that your leader and Dr. Fanning's article will direct the efforts of public health authorities towards this end.—I am, etc.,

East Horsley, Surrey.

BASIL S. GRANT.

Treatment in Early Cases of Influenza and the Common Cold

SIR,—For the past three years I have been treating successfully cases of influenza and the common cold by intravenous injections containing quinine bihydrochloride. Cases of the common cold subsided within 12–48 hours after a single injection of 2 gr. (0.31 g.) quinine bihydrochloride dissolved in 2 ml. of normal saline, while cases of influenza subsided within 2–3 days with the same injection and repeated within 12 hours. A number of cases were specially recorded during 1946 and a large majority of them responded successfully. Of course I am referring to cases seen within the first 24 hours. The object of this report is to interest investigators of this universal malady.

Virchow (1927) thought there might be a small decrease or increase in gaseous metabolism. Hardikar (1925) also noted a change in metabolism. Brown (1912), Solis-Cohen, Kolmer, and Heist (1917) noted that quinine is quite bactericidal to pneumococci *in vitro*. Probably quinine acts by checking the abnormal metabolism which is going on during the infection, or it may be that it interferes with the action of the virus concerned.—I am, etc.,

Belfast.

E. O. BLAKE.

REFERENCES

- Brown, O. H. (1912). *Proc. Am. Soc. Biol. Chemists*, p. xxxv.
Hardikar, S. W. (1925). *J. Pharmacol.*, **25**, 175.
Solis-Cohen, S., Kolmer, J. A., Heist, G. D. (1917). *J. infect. Dis.*, **20**, 272.
Virchow, R. (1927). *Arch. exp. Path. Pharmacol.*, **127**, 28.

Technique of Venepuncture

SIR,—Further to Dr. J. A. M. Cameron's remarks on venepuncture (March 29, p. 425), I have found the essential points to be: (a) efficient local anaesthesia; (b) complete extension of the arm at the elbow and supination of the forearm; (c) congestion of the veins of the forearm by a constriction higher up the arm at about 70 mm. Hg pressure; (d) decision as to site of puncture to be made by palpation rather than inspection; (e) at the moment of insertion of the needle, traction to be made on the skin distal to the site of entry and at the same

time the patient or donor to be asked to clench the fist; and (f) attention to sharpness of the puncture needle.

I have found an oblique light shining on wet skin to be most helpful in demonstrating a slight convexity of the skin due to a palpable but not otherwise visible underlying vein, and whenever possible I try to avail myself of this aid, but do not regard it as essential. In many cases a suitable vein is plainly visible without this aid, and in most of the others the course of the vein will be identifiable by the palpating finger. In less than 0.5% of adults the only superficial veins are very small and close to the surface. No larger vein is palpable, and evidently most of the return circulation from the hand and forearm is by vessels deep to the bicipital fascia. This condition is not always bilateral, and both arms should be examined, but where it is, I do not regard these persons as suitable for venepuncture in the antecubital fossa. I agree with Dr. Cameron that attention to stance is of prime importance for successful results.—I am, etc.,

Cambridge.

C. B. V. WALKER.

Sphygmomanometer Armlet as Tourniquet

SIR,—The daily Press recently reported a case in which a tourniquet was not removed from a patient's leg following an operation, and in which the mishap had a fatal termination. I have for some time employed in orthopaedic operations requiring a tourniquet a simple apparatus connected with an ordinary anaesthetic machine, by which an inflatable sphygmomanometer armlet is connected by tubing to a reducing valve on the CO₂ cylinder. The latter is adjustable to give pressures between 150 and 250 mm. Hg, and, since the exact pressure maintained does not much affect the use of the gas for anaesthetic purposes, the supply is made by a branch tube both to the armlet and to the flowmeter of the machine.

The apparatus is indicated in the accompanying rough sketch, and has proved of real practical use. One becomes at once independent of the minute leaks which seem an inseparable feature of such pneumatic systems, and the surgeon is spared the irritation of the pressure running down in the middle of the operation. Above all, it is impossible for the patient by some mischance to be returned to the ward with the tourniquet still in action.

Some modification of the commonly used reducing valves and armlets is of course necessary for the use proposed, and unfortunately there does not appear to be any commercial source of such apparatus. I am, however, making inquiries with a view to their manufacture, and hope to write later when a more finished type of apparatus is available.—I am, etc.,

Wallasey, Cheshire.

R. L. WYNNE.

Control of Leprosy

SIR,—In the interesting leading article (Nov. 2, 1946, p. 655) on the control of leprosy occurs the statement: "It soon became evident that compulsory isolation had become worse than useless." At first sight this remark, from its context, might have been regarded as applied to India alone, but when coupled in the same article with a disparaging reference to "the Biblical and Middle Ages plan of compulsory segregation" and a later contrasting mention of "modern methods," it is evident that it was intended as an authoritative statement of the present position, and that by the up-to-date person the policy of compulsory segregation should be regarded as a futile anachronism.

Having been advocating and carrying out a policy of compulsory segregation for the control of leprosy in Fiji and the neighbouring islands of the Pacific since 1930, I am naturally interested in the