The erythrocyte sedimentation rate was 59 mm. in 1 hour (Westergren). The white cells numbered 12,700, with a normal differential count. The Wassermann and Kahn reactions were negative.

As these clinical investigations strengthened the diagnosis of a right suprarenal chromaffinoma, it was decided to perform an exploratory laparotomy. Before this could be done, however, the patient had another severe attack, which proved fatal.

Post-mortem examination.—This was carried out at Charing Cross Hospital on Dec. 5, 1944, and the report by Dr. Vines was as follows. "*Brain.* No abnormality was found. The pituitary appeared normal. *Thorax: Lungs:* Both lungs showed some emphysema and chronic bronchitis; they were closely bound to the thoracic wall by chronic fibrous adhesions. In both lungs there was extensive patchy fibrosis at the apices, but no active tuberculosis was seen. *Heart:* There was gross hypertrophy of the left ventricular wall. The tone of the muscle was good. The aorta was rather small and some early atheroma was present. *Abdomen:* The only pronounced changes were seen in the kidneys and the right adrenal. In the latter there was a spherical tumour about 1 in. (2.5 cm.) in diameter projecting forwards. It felt soft and almost cystic, and was not incised. It did not appear to infiltrate, and no metastatic deposits were found. In both kidneys the cortical band was reduced in width and the capsules were adherent, leaving a finely granular surface. In the lower two-thirds of the right kidney there were numerous areas of cortical necrosis suggestive of malignant hypertensive changes; the total changes in the kidneys suggested uraemia as the ultimate cause of death."

Histological examination of the adrenal tumour revealed a fairly typical medullary chromaffinoma. In accordance with the clinical evidence and the presence of an actively secreting tumour, the kidneys histologically showed well-marked hypertensive changes with arteriosclerosis and necrosis. Although the adrenal tumour should have been easily removable, in view of the extensive renal damage it is unlikely that the patient would have survived operation.

There are, of course, reasonable grounds for belief that the accessory chromaphil bodies also secrete adrenaline. It is not surprising, therefore, that similar tumour masses should be found associated with one or other of these bodies anywhere in the sympathetic autonomous nervous system, causing similar paroxysmal attacks of hypertension. For instance, cases of paroxysmal hypertension associated with tumours of the carotid sinus and carotid body and also of the aortic bodies have been reported. The structure and function of both are probably identical. They consist of masses of glandular-looking cells arranged in characteristic clumps resembling those of chromaphil cells. Such a tumour of the carotid sinus or carotid bodies has often been mistaken for an enlarged cervical gland.

In essential hypertension cerebral hypertensive crises may simulate paroxysmal hypertensive attacks such as I have described. It is important to remember these occurrences in order that errors of diagnosis, and thus of treatment, may be avoided.

Summary and Conclusions

A case of medullary chromaffinoma of the right adrenal medulla in a woman aged 38, diagnosed during life, is reported and the syndrome discussed. The patient, unfortunately, died from secondary malignant hypertension and uraemia before operation could be performed. The history of the case and the results of various examinations conducted during and after attacks are described.

The symptom-complex was caused by a tumour of the right adrenal medulla which consisted of adrenaline-secreting chromaffin cells discharging into the blood paroxysmally, though possibly also constantly, small quantities of adrenaline. This caused hypertension with periodic exacerbations to very high levels—sometimes to over 300 mm. Hg—glycosuria, and, when the kidneys became ischaemic, albuminuria and uraemia. The vasomotor crises were also due to a discharge of adrenaline into the blood stream.

The treatment is removal of the tumour, but this is likely to be successful only if it is recognized early and before permanent changes in the kidneys have occurred. The tumour, as in the present case, is seldom malignant and is usually unilateral. It can therefore be concluded that a certain proportion of all cases of paroxysmal hypertension are due to a hyperfunctioning medullary tumour. In this case there was evidently gross hyperfunction, but cases also occur in which the tumour is less active and the symptoms correspondingly less pronounced.

Within the past few years a number of tumours of chromaphil tissue have been reported, some authenticated at necropsy, others actually removed surgically with beneficial results.

I have to thank Dr. P. J. W. Mills, medical superintendent, Lister Emergency Hospital, Hitchin, for permission to publish this case; Dr. H. W. C. Vines for the post-mortem and histological findings; Dr. J. W. Linnell and Dr. A. A. Cunningham for their kind encouragement; and Dr. C. Hilliard and Dr. M. Brown and their technical staffs for the radiological and pathological investigations respectively.

BIBLIOGRAPHY

- Cameron, A. T. (1935). *Recent Advances in Endocrinology*, 2nd ed., p. 191, London.
 Goldzieher, M. A. (1929). *The Adrenals*; Macmillan, New York.
 — (1932). *Endocrinology*, 16, 20.
 Hick, F. K. (1933). *Arch. Path.*, 15, 665.
 Kalk, H. (1934). *Klin. Wschr.*, 13, 613.
 King, E. S. J. (1931). *Path. Bact.*, 34, 447.
 Labbé, M., et al. (1929). *Bull. Mém. Soc. méd. Hôp. Paris*, 45, 952.
 — (1934). *Ibid.*, 50, 337.
 Mayo, C. H. (1927). *Coll. Pap. Mayo Clin.*, 19, 732.
 Oppel, V. A. (1928). *Arch. klin. Chir.*, 149, 301.
 Porter, M. E., and Porter, M. F. (1930). *Surg. Gynec. Obstet.*, 50, 160.
 Rabin, C. B. (1929). *Arch. Path.*, 7, 228.
 Rogers, E. (1932). *Amer. Heart J.*, 8, 269.
 Shapiro, S. (1926). *Endocrinology*, 10, 413.
 Shipley, A. M. (1929). *Ann. Surg.*, 90, 742.
 Vaquez, H., Donzelot, E., and Géraudel, E. (1929). *Presse méd.*, 37, 169.
 Volhard, F. (1931). In Bergmann and Staehlin's *Handbuch der Inneren Medizin*. 2nd ed., vol. 2, Pt. I, p. 390; Pt. II, p. 1742, Springer, Berlin.
 Werner, A. A. (1942). *Endocrinology*, Philadelphia.

A CASE OF PHAECHROMOCYTOMA

BY

J. M. K. SPALDING, B.M., B.Ch.

The surgical removal of a chromaffin tumour or phaeochromocytoma of the adrenal medulla is rare enough to warrant the report of a recent case.

Case History

The patient, a 24-year-old secretary, was admitted on June 4, 1945, under the care of Prof. Ellis, complaining of blurring of vision and of attacks of upper abdominal pain and vomiting with palpitations and headache. There was nothing relevant in her past history. The attacks began two years before admission and at first recurred about once every four weeks. They became gradually more frequent, and on admission she was having one or more daily. Immediately before an attack she had a premonition that it was about to occur. She then felt a burning pain in the epigastrium, followed by severe headache, nausea, and often vomiting. She had severe palpitations, "as though her heart would burst through her ribs," her hands and face became pale, and she felt cramp-like pains in the limbs. The attacks occurred at any time of the day or night and usually lasted 10 to 20 minutes, but might last an hour. On three occasions she had lost consciousness. After an attack she felt weak, her arms and face were flushed, and she had a sensation of "pins and needles" in her arms. Five months before admission she was examined elsewhere and found to have a raised blood pressure. The fundi and urine were normal, but the cerebrospinal fluid contained 80 mg. of protein per 100 ml. No record of the pressure of the cerebrospinal fluid at this time was available. She was noted as unduly excitable and her condition was dismissed as functional, a

diagnosis for which there was ample foundation in her social history. Four weeks before admission she first noticed blurring of vision in the right eye immediately after an attack, and two weeks later the left eye was similarly affected.

On examination she was "nervous" and anxious to demonstrate that her disease was not functional. There were no abnormal findings apart from the blood pressure, fundi, and urine, and there was no demonstrable enlargement of the heart or peripheral arteries. Blood-pressure readings between attacks ranged from 120/80 to 250/170, and during attacks from 230/135 to 300+/205 mm. Hg. Prof. Mann reported on the fundi: "The ophthalmic condition resembled that seen in albuminuric retinitis in that there were numerous flame-shaped haemorrhages, fluffy patches of exudate, and a well-marked macula fan, with slight blurring of the disk but no true papilloedema. The condition was not absolutely typical in that the arteries, which were all reduced to threads, were narrower than is normal in albuminuria." The urine consistently contained a trace of protein, and its specific gravity varied from 1008 to 1024.

She had frequent attacks after admission, and a moderately severe one was followed in detail. It began while the patient was sitting quietly in a chair. When seen three minutes later she looked pale and her respirations were deep and rapid. The cardiac impulse was extremely forcible, but the radial pulse was barely perceptible. There was marked "goose-skin," and the pupils were widely dilated and did not react to light. The blood pressure was 300/200 mm. Hg, and the pulse rate 120. Ten minutes after the onset there was subjective improvement, after 15 minutes the systolic and diastolic pressure began to fall, after 30 minutes she became flushed, and after one hour the blood pressure reached a resting level. It was another hour before the pupils returned to normal.

Special Investigations.—The haemoglobin was 118%, the leucocyte count 7,000 per c.mm., and the Wassermann reaction negative. The blood sugar during an attack was 107 mg., during an interval 120 mg. per 100 ml. Ventriculography and examination of the cerebrospinal fluid revealed nothing abnormal. The patient was profoundly collapsed for 12 hours after ventriculography. An electrocardiograph showed left axis deviation and an inverted T₁ wave. Intravenous pyelography was normal; the right kidney was 4.5 cm. lower than the left. A barium meal revealed no abnormality. An unsuccessful attempt was made to demonstrate the presence of adrenaline in blood taken during an attack.

A provisional diagnosis of phaeochromocytoma was made, and it was decided to explore the suprarenals. In answer to direct questioning the patient admitted having slight pain in the right loin, and she believed that lying on the right side precipitated an attack; it therefore was decided to explore the right side first.

Operation.—This was performed by Mr. Corry on July 24 under ethyl chloride and ether anaesthesia. A tumour weighing 86 g. was removed from the suprarenal region through an incision in the right flank. Neither induction of anaesthesia nor handling the tumour altered the blood pressure, which remained about 180/150 mm. Hg throughout the operation. But half an hour after the removal of the tumour the blood pressure fell to 80/60 and the patient appeared profoundly shocked. During the next two hours she was given 20 ml. of "eschatin" 20 minims (1.2 ml.) of adrenaline, and 1 pint (568 ml.) of normal saline parenterally. At the end of this time the systolic blood pressure had risen to 100 mm. Hg and her condition began to improve. During the first 24 hours after operation she was given 120 ml. of eschatin. Convalescence was uneventful, and until she was discharged from hospital 18 days after operation her blood pressure maintained a steady level of 140/100. The lesions in her fundi were regressing when she left hospital and her eyesight was improving rapidly.

Fifteen months after operation she was doing a full-time clerical job, was having no symptoms, and was about to be married. Her blood pressure was 135/100 and the urine was clear. Her vision was 6/5 on the right and 6/9 on the left, and was not improved by correction. Prof. Mann reported: "The disks are normal and the veins and arteries are of normal calibre and have healthy walls. The whole fundus colour is slightly pale. The macula, however, shows gross abnormality in both eyes, being pigmented and covered with

small dots of exudate and minute atrophic patches all rather resembling senile macular degeneration. It is remarkable that with so definite an ophthalmoscopic change the visual acuity should be so good."

Pathology and Histology (Dr. Biggs).—The specimen consisted of an enlarged adrenal gland measuring 7.5 by 7.5 by 5 cm. and weighing 85 g. Externally there was a thin layer of cortical tissue with a ridge of cortex over the upper pole. On section the medullary tissue was replaced by a homogeneous grey-white tumour. Microscopically the tumour consisted of dense masses of darkly staining cells separated by sinusoidal blood channels. The tumour nuclei varied in size, but there were few nuclear divisions and no giant-cell forms. The cells stored neither fat nor glycogen, but chrome fixation showed that the cells had an affinity for chrome salts. The general structure and histochemical reactions are characteristic of a phaeochromocytoma of the adrenal medulla; the regular pattern, the infrequency of nuclear divisions, and the absence of giant cells suggest that the tumour is benign. Biological essay, kindly performed by Prof. Burn, showed that it contained 5 mg. of adrenaline per gramme of tumour (normal=0.4 mg./g.).

Comment

The characteristic syndrome associated with phaeochromocytoma is paroxysmal hypertension with palpitations, headache, nausea, and abdominal pain (Padis, 1945). Pallor and tingling of the extremities are also common, profuse sweating sometimes occurs (Biskind *et al.*, 1941; McCullagh and Engel, 1942; Evans and Stewart, 1942; Thorn *et al.*, 1944), and occasionally there is persistent hypertension (McCullagh and Engel, 1942, Thorn *et al.*, 1944). The adrenaline content of the tumour is variable, as much as 20 mg. per gramme having been reported (MacKeith, 1944). Raised blood-sugar levels and glycosuria have been found in several cases (Duncan *et al.*, 1944). Post-operative shock is greater than would be expected from the severity of the operation and responds to large doses of adrenaline (Biskind *et al.*, 1941; Duncan *et al.*, 1944), and parenteral fluid has also been recommended (McCullagh and Engel, 1942).

Biskind and his colleagues (1941) reviewed 29 reported cases treated by operation. Five patients died soon after operation, 22 remained well for a period of 7 months to 13 years, 1 had a recurrence after 6 years, and 1 died from metastases. MacKeith (1944) collected 165 case reports, mainly from necropsies. In 16 of these the tumour was bilateral and in 15 malignant.

I wish to thank Prof. Ellis and Dr. Alice Stewart for their help in preparing this report, Prof. Burn and Miss Bülbring for biological assay of the tumour, and Prof. Mann for reports on the eyes.

REFERENCES

- Biskind, G. R., Meyer, M. A., and Beadner, S. A. (1941). *J. clin. Endocrinol.*, **1**, 113.
Duncan, L. E., Semans, J. H., and Howard, J. E. (1944). *Ann. intern. Med.*, **20**, 815.
Evans, W. F., and Stewart, H. J. (1942). *Amer. Heart J.*, **24**, 835.
McCullagh, E. P., and Engel, W. J. (1942). *Ann Surg.*, **116**, 61.
MacKeith, R. (1944). *Brit. Heart J.*, **6**, 1.
Padis, N. (1945). *Clinics*, **4**, 87.
Thorn, G. W., Hindle, J. A., and Sandmeyer, J. A. (1944). *Ann. intern. Med.*, **21**, 122.

The L.C.C. has agreed, at the request of the Ministry of Health, to the establishment at Sutton Emergency Hospital of an industrial neurosis unit of 100 beds (50 for each sex). This will be used for the reception of persons apparently suffering from neurotic disorder and referred from employment exchanges. They will be given a trial, as in-patients, in industrial processes with a view to later treatment, training, or employment according to their ascertained need. The whole hospital (500 beds), which has hitherto accommodated a number of general cases, is to be reserved for patients suffering from psychoneurosis, and will continue to rank as an emergency medical service hospital. Two full-time psychologists are to be appointed, one for the industrial neurosis unit and the other for the remaining 400 patients.