THE USE, COST, AND PROVISION OF INFANT HEALTH HOSTELS

BY

J. TUDOR LEWIS, M.D., D.P.H.

Medical Officer of Health, County Borough of West Bromwich

It has previously been suggested that homes or hostels for mothers and their babies, which would bridge the gap between their discharge from maternity unit or hospital and their return to poor home conditions, should be a feature of the infant health services of the future (Lewis, 1946). Such hostels, however, might have a more important role in counteracting unsatisfactory social conditions than that of mere post-natal sanctuaries, and the subject clearly merits detailed thought. The local authority with which I was connected decided to set up a hostel, and the problem was examined from the local point of view. The results may be of general interest, because, apart from a review by the *Lancet* (1946) of convalescent-home facilities, this subject, especially as applied to the prevention of baby deaths and the promotion of infant health, does not appear to have received adequate objective treatment. For want of a better name, these hostels might be called infant health hostels, and their purpose would be to combat the unsatisfactory social and environmental conditions that only too often interfere with the rearing of healthy infants.

It would not be difficult to write at length on how they would do this. In brief, however, they would be used to conserve the health and well-being of the mother in the pre-natal period, and so prevent premature labour and reduce the incidence of prematurity; to act as the link between maternity units and hospitals and the home, and thus save the infant from being exposed to unsatisfactory home conditions until it could stand them—the baby would, in fact, be “hardened off” in a hostel; to assist certain mothers by providing means by which for a period they could give all their attention to the baby; to help selected unmarried mothers; and to give concentrated teaching in homecraft and baby-care, for most of the mothers admitted would need instruction in these important subjects.

To obtain the greatest benefit from an infant health hostel it would be necessary to ascertain the households where conditions were prejudicial to the healthy rearing of an infant, and to do this as early in the pregnancy as possible. We have called these households “observation households,” for they should be kept under careful observation throughout the pregnancy, with a view to ameliorating the adverse conditions.

Cost of Infant Health Hostels

General.—In setting up an infant health hostel there seem to be two principles to follow: (a) As the function of the hostel would be to give individual help to mothers beset with social difficulties and to impart concentrated instruction in homecraft and baby-care, the proportion of staff to patients should be relatively high. (b) Although the babies admitted would not be under 2 or 3 weeks old, there would still be a risk of the various forms of cross-infection to which babies in institutions are always liable. If a serious infection were to break out, the smaller the unit the fewer would be the babies exposed. Therefore small units are to be preferred to large ones. The cost of a hostel would depend on its size and the staff necessary.

Size of the Hostel.—What is the smallest unit that can be made an economical proposition? The nursing staff would have duties peculiar to this form of hostel, for there would be little nursing as such, and no need for night duty so long as some responsible person was available when necessary. In order to allow the nursing staff off-duty and holidays in accordance with the Rushcliffe recommendations a minimum of three would be needed. It is considered that a staff of three could run a hostel containing 10 to 12 mothers and babies, and this would accord with principles (a) and (b) mentioned above. Considering the fairly heavy overhead charges to which a smaller hostel would be subject, the availability of suitable premises, and the scale of provision of hostels, it is probable that a hostel of 10 to 12 beds and cots is the optimal size to aim at.

Staffing.—The matron, or sister-in-charge, would be the key to the success of the hostel and must be carefully selected. She

should be an S.R.N., possessing the S.C.M. and the Health Visitor's Certificate, and should have had practical experience of sociological and public health work. An active interest in and knowledge of baby-care and mothercraft should be an essential qualification. The other two nurses should have special knowledge and interest in young babies, should possess the S.R.N. with experience of infants, or should be State registered children's nurses. As in the case of the matron, an interest in the teaching of mothercraft would be essential. In view of the special qualifications required it might be possible to pay a small honorarium over and above the Rushcliffe scales in order to attract suitable people. It is anticipated that the mothers themselves would help in the domestic work of the lighter type, so there would be no need for a large domestic staff. One cook with two daily workers would be sufficient.

Capital Outlay.—When new maternity units or hospitals are being built it might be possible to incorporate an infant health hostel in the main building, thus reducing overhead charges, but for the present it would be necessary to adapt a large private house. Some guidance on what is required is given by Brodie (1946), who deals with homes for unmarried mothers; but, with minor modifications, her recommendations would apply to infant health hostels, as also would those put forward by the National Council for the Unmarried Mother and her Child (1946). The large old-fashioned type of house would generally have to be used, and it would need a good deal of alteration. A house of the size and type required could be purchased to-day for about £5,000; alterations might amount to £1,000 and equipment to £2,000—a capital outlay of about £8,000.

Yearly Expenditure.—The following estimate of yearly expenditure is partly based on the running costs of a small maternity home of 12 beds, allowance having been made for the differences in the types of institution.

Salaries of nursing staff	£770
" domestic staff	568
National insurance, rates, and insurance ..	100
Superannuation contribution	40
Printing and stationery	25
Cleaning materials, painting, repairs, renewals	100
Maintenance of patients and staff	400
Petty cash, sundries, telephone	30
Bedding, drapery, earthenware goods	60
Light and fuel	300
Furniture replacements	50
Drugs and medicine	30
Gardening	300
Total	£2,773

Assuming that the 10 beds and cots were occupied throughout the year, the approximate cost per mother and baby would be £5 a week.

Provision of Infant Health Hostels

The scale of provision of infant health hostels would vary according to the area. Where social conditions were very bad the need would be greatest, and this would usually be in industrial areas where housing conditions were bad and the social environment poor. Powers to set up infant health hostels are apparently given by Section 204 of the Public Health Act, 1936.

The following gives some rough idea of the extent of the need. In 1945, of the 77 infant deaths investigated by Lewis and Blackwood (1946) it was estimated that a hostel might have helped to save 19 babies; in 1946 in a similar survey the estimated number was 16. It would, however, be the greatest mistake to consider only mortality, for there would be many babies whose lives might not be in acute danger, but who would benefit in health and well-being from being admitted to a hostel. This problem is better considered in terms of infant health than of infant mortality. By so doing, the net would be widely cast, and in saving the lives of the few the many would benefit. In the ascertainment of observation households referred to above it was estimated that between July and December, 1946, 25 mothers or mothers and babies would have benefited by admission to a hostel; over the whole year this would mean that about 50 should have been admitted. During 1946 there were 61 infant deaths in the area, so that according to these figures provision might be made for approximately one bed for each infant death.

Assuming that each mother and baby remained in the hostel for one month, 10 beds would provide accommodation for 120 cases a year. It is likely, however, that many mothers would be unwilling or unable to stay that time; to offset this, some might stay longer and there might be an occasional long-period case. If this approximate standard were adopted it is clear that only those authorities with over 100 infant deaths a year would find these hostels an economic proposition, unless joint hostels were set up or arrangements made for the admission of mothers and babies from adjoining local authorities.

There is virtually no well-documented practical experience of the use of hostels for saving infant lives, for while many authorities have made arrangements for sending mothers and babies to convalescent homes, a definite attempt to combat the effects of adverse social conditions at all stages of foetal and infant life by the extensive use of a hostel has never, so far as can be ascertained, been made. It would, moreover, avail little to set up a hostel unless this were accompanied by the ascertainment of observation households as early in pregnancy as possible. Remembering the relatively high cost, both capital and revenue, of infant health hostels, and that they must at first be experimental, it is evident that no extensive provision on the scale of one bed per infant death could be considered, but as an experiment in social and preventive medicine their use in selected areas may rightly be advocated. They can do no harm; they are certain to do some good; they may prove to be a means of saving many infant lives. To achieve this any reasonable expenditure of public money would be justifiable.

Summary

The uses of infant health hostels as a means of combating adverse social conditions are set out, the functions of hostels are referred to, and the indications for admitting mothers and babies are considered.

The optimal size of a hostel is 10 to 12 beds and cots. Staffing is dealt with and an estimate of the capital and revenue expenditure made.

The need for hostels is assessed. It is suggested that provision might be made on the scale of approximately one bed and cot per infant death per year. Until, however, the uses and value of infant health hostels are more clearly defined, they should in the first instance be experimental, and be set up in those areas where the infant mortality rate is high, as an organized attempt to reduce it.

REFERENCES

- Brodie, G. I. (1946). *Mon. Bull. Min. Hlth. E.P.H. Lab. Serv.*, 5, 203. *Lancet*, 1946, 2, 568.
 Lewis, J. T. (1946). *British Medical Journal*, 2, 893.
 — and Blackwood, M. W. (1946). *Mon. Bull. Min. Hlth. E.P.H. Lab. Serv.*, 5, 190.
 National Council for the Unmarried Mother and her Child (1946). *Schemes for Homes and Hostels*, London.

SOCIETY OF BRITISH NEUROLOGICAL SURGEONS

The Society held its spring meeting in Lisbon on April 10 and 11 under the presidency of Sir Hugh Cairns. The meeting was arranged to pay tribute to Prof. Egas Moniz for his pioneer work. It was he who introduced arteriography (in 1927), which was used in cerebral work first. The leucotomy operation for certain mental disorders was a later conception and was first performed by Prof. Lima in 1935. The visitors, including members from several European countries, were welcomed by Prof. Flores, of the Lisbon Faculty of Medicine, Prof. Moniz's successor in the Chair of Neurology.

Prof. Moniz spoke on thrombosis of the internal carotid and its branches, and on the interpretation of cerebral angiograms, especially the striking difference in circulation rate through the internal and external carotid systems. Prof. Lima described the diagnosis of subdural haematomas by arteriography. Prof. Furtado and Dr. Marques spoke about the Arnold Chiari malformation, Prof. da Costa on the neuroblastomas, and Prof. Fernandes on leucotomy. The technique of carotid arteriography was demonstrated by Dr. Imaginário, and of aortic arteriography by Dr. dos Santos. An excellent exhibition of cerebral angiograms included the early experimental procedures and instruments used for them, now of interest historically. Several communications, mostly relating to Prof. Moniz's work, were given by the visitors, including Mr. Norman Dott's on intracranial aneurysms, and Prof. Geoffrey Jefferson's on frontal lobectomy.

Reports of Societies

COMMON COLD RESEARCH UNIT

INTERIM REPORT ON A TRANSMISSION EXPERIMENT

A meeting of the Epidemiology and State Medicine Section of the Royal Society of Medicine was held on May 2 at the Common Cold Research Unit, Harvard Hospital, Salisbury, where for about a year, under the auspices of the Medical Research Council and the Ministry of Health, an investigation into the aetiology of the common cold has been undertaken. A succession of groups of volunteers have accepted isolation for a fortnight at a time in order that they might be tested with the common cold virus. The conditions and the outcome of the research so far are described in this report, and are the subject of a leading article at p. 645.

Members of the Section and others, who attended in large numbers, were received by the staff of the unit. They visited the isolation quarters—flats—of the volunteers and the air hygiene laboratory, and then assembled in the lecture hall, where a film produced by the Central Office of Information was projected. Dr. H. J. PARRISH, who presided, said that the common cold was responsible for the loss of 40 million man-days a year; yet he believed this was the first time the Section had had a discussion on the subject.

History of Harvard Hospital

Dr. W. H. BRADLEY gave an account of the origin of the hospital and of the unit. At about the time of Dunkirk, Harvard University planned to send to this country a group of field and laboratory workers to assist in the control of epidemics which were expected to arise. The Harvard unit grew into a team of more than 100 doctors and others; 22 buildings, prefabricated in the United States, were transhipped to this country and erected on a site near Salisbury secured and prepared by the Ministry of Health, and the hospital was formally opened in September, 1941. In selecting the location regard was paid to the suitability of the place for epidemiological studies of the detailed and intimate type called "clinical epidemiology." This part of Wiltshire had for the epidemiologist much the same kind of advantages as the "Pickles" country of Wensleydale. The hospital was for a while perhaps overburdened with beds (125); it became an E.M.S. hospital for both civil and military cases of communicable diseases, and no group of patients suffering from such diseases ever before received so much investigation and therapy. The course of the war unfortunately prevented the full benefit of the studies from being reaped, but some of the work was published in the reports for the war years of the Chief Medical Officer of the Ministry. With America's entrance into the war the hospital passed into the hands of the U.S. Army, and became the central laboratory for the American Army Medical Corps in the European theatre. The laboratories were extended and adapted for special purposes, and research units were established—one, for example, for the study of rickettsial disease. The place rapidly acquired a reputation as one of the foremost laboratories devoted to preventive medicine. With the invasion of France in 1944 the central laboratory was transferred to the Pasteur Institute in Paris, and with the cessation of hostilities in May, 1945, the American military authorities finally withdrew, though the place remained an important link between the United States and Great Britain and between Harvard University and its Faculty of Medicine on the one hand and the Ministry of Health and the Medical Research Council on the other.

The hospital was built specifically as an institute for the study of communicable diseases, and it was decided to adapt it to the purposes of a long-term research into the aetiology and transmission of the common cold. The staff and personnel house was converted into flats for the use of volunteers, and the staff of the unit and of some of the laboratories was housed in what had previously been the wards. The work was started in May, 1946, on plans which were first discussed less than six months previously.

It was seldom clear how projects of this kind began, but his first knowledge of it was a memorandum by Dr. C. H. Andrewes