first by Lister, who describes mixing blood and carbolic acid into a paste and using it to fill wounds.

Results.—When the dressing was removed healing in all cases was either complete or sufficiently advanced to shield the deeper parts of the wound from infection. All wounds were dry in three weeks from operation, and have remained so since for periods varying from nine months to two and a half years. It may be noted that no particular benefit has been observed to follow the local or general use of penicillin in cases in which all the infected bony and soft tissue could not be removed. Such sites of infection as the upper end of the femur or the vertebrae still remain most intractable once a communication, temporary or permanent, has been formed with the outer surface of the body.

Summary and Conclusions

Three series of cases are reported: 31 of acute osteomyelitis; 4 of acute suppurative arthritis; 5 of chronic osteomyelitis.

Of the acute cases 50% were in infants under 1 year old.

Dosages of penicillin were relatively small—1,000 units per lb. (454 g.) body weight per twenty-four hours for three weeks.

In acute osteomyelitis and arthritis, surgery was confined to aspiration of pus. Immobilization was partial and for short periods.

Radiological changes observed were far less serious clinically than might have been expected from their marked character.

In suitable cases of chronic osteomyelitis excision of infected tissue and primary suture is advised.

Mortality was nil. Complete restoration of function occurred in all but one of the acute cases. There was apparent complete healing in all the chronic cases.

We wish to acknowledge the kindness of physicians and surgeons at the Hospital for Sick Children, Great Ormond Street, who have permitted us to include some of their cases in this paper. We are also much indebted to Miss Eileen Rawlings for her tireless clerical assistance.

BIBLIOGRAPHY

- Agerholm, M., and Trueta, J. (1946). *Lancet*, 1, 877.
Aird, I. (1945). *Proc. roy. Soc. Med.*, 38, 569.
— (1946). Fleming: *Penicillin, Its Practical Application*, p. 200, London.
Bodian, M. (1945). *Proc. roy. Soc. Med.*, 38, 572.
— (1946). *Arch. Dis. Childh.*, 21, 13.
McAdam, I. W. J. (1945). *Brit. J. Surg.*, 33, 167.
Paterson, D., and Bodian, M. (1946). Fleming: *Penicillin, Its Practical Application*, p. 252, London.
Royal Society of Medicine Discussion on Treatment of Acute Osteomyelitis with Penicillin, *Lancet*, 1946, 1, 236.

Extensions to Joyce Green Hospital (L.C.C.), Dartford, including new wards to accommodate 120 beds, a new main operating theatre, and a nurses' home, were completed in 1939, but the formal opening by Mr. A. Reginald Stamp, chairman of the Hospitals and Medical Services Committee of the London County Council, was delayed until April of this year. This was originally one of the three river hospitals under the control of the Metropolitan Asylums Board. The main hospital buildings were opened in 1903, following smallpox epidemics in the two previous years, to take the place of the Thames ships which were formerly employed as floating hospitals. Afterwards it was used for infectious diseases generally, and during the last eight years it has admitted general medical and surgical patients. As a war emergency hospital its previous accommodation of 986 beds was almost doubled. The wards newly constructed, however, were not brought into use during the war because the large amount of glass made them unsafe, and indeed the hospital suffered extensively and repeatedly from blast. The total cost of the new extensions is £134,000. One of the new wards houses neurosurgery and physiotherapy units and the other is for children. Ear, nose, and throat, ophthalmic, and dental clinics have been established in the new premises. At the opening ceremony Mr. Stamp said that the extensions were part of a great reconstruction programme for the Council's 76 hospitals, but he added that this was the last year that Joyce Green and all the other hospitals would be under L.C.C. control. At County Hall they were anxious that the hospitals should be transferred to the National Health Service as going concerns. Sir Allen Daley paid a tribute to the medical superintendent, Dr. Maurice Mitman, the matron, Miss Vivian, and the staff at Joyce Green for the first-class medical and surgical work which had been done at the hospital throughout the war notwithstanding recurring air-raid damage and continual danger.

REFRIGERATION ANAESTHESIA

BY

E. S. R. HUGHES, M.D., M.S., F.R.C.S.

Refrigeration anaesthesia for limb amputations was introduced into the Royal Melbourne Hospital towards the end of 1944. Since then 25 lower-limb amputations have been performed with the aid of this technique, and they are here reported. With one exception all the amputations have been through the lower third of the thigh. It is not possible to reduce results fairly to statistical form owing to the diverse nature of the patients' condition. In the table the predominant constitutional factor has been indicated as either "atherosclerosis" or "diabetes mellitus." Some of the former group have had accompanying cardiac failure, renal failure, or cerebral degeneration; some of the latter group have been admitted in a coma or semicoma, and all have had atherosclerosis to a varying degree.

The local condition that required amputation has varied; most frequently it was progressive gangrene of the foot or gangrene complicated by severe pain or infection. In addition there were two cases of popliteal thrombosis accompanied by severe pain; two cases of gross infection of ulcers of the leg causing deterioration of the general condition of the patient despite conservative measures; two cases with epithelioma of the leg, one of which was severely infected and had metastatic glands in the groin, and the other equally badly infected and complicated by a pathological compound fracture of the tibia and fibula; finally, there was one case of a mangled foot in a shocked patient aged 69 years.

Technique

The simplest technique has been used throughout this series; the more elaborate and costly apparatus employed by some clinics abroad has not been available.

The Ice.—The ice has been procured from a city firm in 50-lb. (22.68-kg.) bags. It is already crushed when delivered, though sometimes the pieces have to be broken further before use. When packing a limb it was found that ice obtained on the previous day and kept at room temperature overnight was easier to handle than that obtained on the same day, because the sharp edges had become rounded off.

The Apparatus.—Two methods have been used to retain the ice in position: a wooden box, 40 by 10 by 10 in. (100 by 25 by 25 cm.) lined with galvanized-iron sheeting and drained at one corner; or two mackintoshes held in position by long stout calico bandages or by a reversed single-limb cradle. Two bags of ice are required initially, and one or two daily for maintenance in cases of prolonged refrigeration.

The Procedure

There are three main sources of infection—the patient, the ice, and the surgeon and his staff. The ice cannot be sterilized, but by adopting the usual procedures with the patient and the operating team the danger of infection is lessened. The limb and pubic region are shaved and the patient is placed on a canvas sheet. The abdomen and affected leg are given the routine pre-operative preparation and covered with sterile towels; the limb is then ready for the application of the tourniquet and placing in the ice.

The tourniquet is not an essential requirement at any stage of the procedure and may be dispensed with altogether. A tourniquet—of soft rubber pressure-tubing—has been used here for reasons given below. The tourniquet is first sterilized either by autoclaving or by boiling. The former method is preferable as a firmer grip may be obtained on the dry tourniquet. After the operator has scrubbed and the limb has been elevated, the sterile tourniquet is applied as tightly as possible just above the site of proposed amputation. It may be held with a reef knot, with sterile Kocher's clamps, or with sponge forceps. Occasionally the tourniquet has snapped some time after the application; hence it is wise to inspect it regularly.

A bed of crushed ice about 4 in. (10 cm.) deep is made, extending from above the tourniquet site to beyond the foot. To ease the pain of the tourniquet application it is advisable to keep the leg encased in ice for a period of an hour beforehand; in such circumstances the leg is lifted clear of the ice, the tourniquet is applied, and the leg is lowered again and covered with ice. The head of the bed is raised on blocks to assist drainage of the ice. The patient is made comfortable and the rest of the body is kept warm. The period of time for which the limb is refrigerated depends on the relation of the local lesion to the general condition of the patient; if the former be the cause of depression of the latter, then by cooling the lesion to the approximate level of 5° C., at which level metabolism is minimal, it becomes physiologically disconnected

operating table with the ice still in position; the surgeon and his team have scrubbed and are ready. A large canvas bag accommodated on a wheeled stand is moved to the foot of the table. The leg is elevated above the ice and the latter is swept into the canvas bag. The towels around the leg are removed, the area is given a final preparation by the surgeon, and the operation proceeds as in cases with the usual forms of anaesthesia. To secure complete haemostasis the tourniquet is left in position until the surgeon orders its removal. There is no need to hurry through the operation, as good anaesthesia persists for an hour or more.

Certain special features of the post-operative period may be noted. The absence of any primary shock, despite the fact that the patient remains fully conscious of his surroundings, permits

Details of 25 Limb Amputations in which Refrigeration Anaesthesia was used

Case No.	Age	Sex	General Condition	Local Condition	Remarks	Duration of Refrigeration	Result
1	60	M.	Atherosclerosis	Gangrenous foot; continued spread	Auricular fibrillation	3 hours	Discharged 4 months after operation. Gas gangrene second post-operative day
2	83	M.	"	Gangrenous foot; severe infection	—	6 days	Discharged 5 weeks after operation. Convalescence uneventful
3	78	M.	"	Huge infected ulcer of leg	Amputation of other leg 6 years previously	4 "	Death 12 days after operation. Bronchopneumonia
4	70	M.	"	Gangrenous foot; intractable pain	Cerebral degeneration	3 "	Discharged 8 weeks after operation. Mild secondary shock and wound infection
5	85	M.	"	Gangrenous foot; intractable pain	—	3 hours	Death 5 days after operation. Severe secondary shock followed by bronchopneumonia
6	67	F.	"	Grossly infected ulcers of leg	Nephrectomy 10 years before. Poor renal function	10 days	Death 8 days after operation. Renal failure
7	65	M.	"	Gangrenous foot; severe infection	—	7 "	Death 17 days after operation. Severe septic stump
8	88	M.	"	Gangrenous foot; continued spread	Cardiac failure	7 "	Death 8 days after operation. Renal failure
9	78	M.	"	Gangrenous foot; intractable pain	Paralysis agitans	6 "	Death 24 hours after operation. Cause undetermined
10	73	M.	"	Popliteal thrombosis	Advanced cardiac failure	7 "	Death 14 days after operation. Cardiac failure; bronchopneumonia
11	78	F.	Diabetes mellitus	Gangrenous foot; severe infection and pain	—	8 "	Death 2 days after operation. Bronchopneumonia
12	74	F.	" "	Gangrenous foot; continued spread	—	4 "	Discharged 4 weeks after operation. Convalescence uneventful
13	73	F.	Atherosclerosis	Epithelioma of leg; severe infection	—	3 hours	Discharged 4 weeks after operation. Convalescence uneventful
14	63	F.	"	Gangrenous foot; severe infection	Diabetes mellitus suspected	3 "	Death 9 weeks after operation. Cardiac failure; stump healed
15	80	F.	Diabetes mellitus	Gangrenous foot; severe infection	—	3 days	Discharged 8 weeks after operation. Mild secondary shock and wound infection
16	78	F.	" "	Gangrenous foot; continued spread	Carcinoma of common bile duct and metastasis in liver; jaundice	3 hours	Death 4 weeks after operation. Stump healed
17	56	F.	" "	Gangrenous foot; severe infection	Coma	3 "	Death 6 days after operation. Remained moribund; bronchopneumonia
18	68	F.	"	Popliteal thrombosis	—	3 "	Discharged 5 weeks after operation. Convalescence uneventful
19	73	F.	Atherosclerosis	Gangrenous foot; intractable pain	Cerebral degeneration	3 "	Death 3 weeks after operation. Cerebral thrombosis; stump healed
20	69	F.	Diabetes mellitus	Gangrenous foot; severe infection	Semicoma	9 days	Discharged 6 weeks after operation. Convalescence uneventful
21	74	F.	Atherosclerosis	Epithelioma of leg; severe infection	Marked debility	3 hours	Discharged 8 weeks after operation. Convalescence uneventful
22	76	F.	"	Gangrenous foot; severe infection and pain	—	2 days	Discharged 5 weeks after operation. Convalescence uneventful
23	59	F.	Diabetes mellitus	Gangrenous foot; severe infection	Semicoma	3 "	To be discharged. Convalescence uneventful
24	75	F.	" "	Gangrenous foot; severe infection	—	24 hours	To be discharged. Convalescence uneventful
25	69	M.	Atherosclerosis	Mangled foot	Shock	3 "	Discharged 10 weeks after operation. Wound infection

from the body. Therefore if the patient is suffering from severe pain or from toxæmia the affected area of the limb is refrigerated for several days without a tourniquet; several hours before the operation, as in all other cases, the refrigerated area is extended by increasing the amount of ice to just above the tourniquet level. The minimum time for which a limb may be safely refrigerated before operation varies from two to three hours.

It is not essential to give any pre-operative medication, but those who are apprehensive may have injections of morphine —1/8–1/4 gr. (8–16 mg.). In those cases of this series requiring such sedation the drug was given before the application of the tourniquet. The complete relief from a severe and persistent pain in the foot has been a most noticeable feature of refrigeration.

Transport to the theatre with the ice in position has not been difficult, as the patient and the ice-container have previously been placed above canvas. The patient is lifted on to the

him to continue his meals without interruption. Systemic sulphonamide therapy has been used in all cases post-operatively and has been a useful measure in avoiding or lessening local and pulmonary infections. A careful watch is maintained on the chest; a cold wet bed, so often seen in the pre-operative period, predisposes to infection. After amputation the stump may be kept cool with ice bags. This has not been done in the present series for two reasons: first, the results have seemed satisfactory without such cooling; secondly, a dry bed after operation is essential. A thick layer of cotton-wool acts as a moderately satisfactory insulating dressing. Stitches have been removed from the wounds at the same time as after operation under the usual anaesthetics.

Complications

Shock.—Primary shock did not occur, but secondary shock was not absolutely prevented, though it was minimized. It would probably have been still further decreased

if the post-operative cooling had been satisfactory in all cases. But it was found that ice without waterproof bags was difficult to handle even by the experienced individual; post-operative cooling so often meant an icy-cold wet bed, with all its attendant disadvantages, that this step was omitted.

Bronchopneumonia.—This was the cause of death of five patients. Wet sheets and blankets, caused by leakage from the ice despite elaborate precautions, were an important contributory factor. The most certain method of prevention is the use of ice-bags or electrically controlled apparatus in place of the bare ice. The ability of the patient to move about and sit upright in bed from the outset and to continue with nourishment has partially neutralized the effects of this unfortunate complicating wet bed.

Local Complications.—Surrounding the limbs with ice has not caused pain. The tourniquet occasioned temporary discomfort in some who had not been previously cooled at the site of application. In all cases the operation has been performed painlessly, and post-operative pain in the stump has not been a feature. No delay was observed in the healing of the stump except when infection supervened. There was no evidence of any sloughing of the flaps despite amputation below a tourniquet in every case (Perlow, 1944). The incidence of infection in this series actually appeared to be lessened, so conforming with published results. In the first case gas-gangrene developed; but at the same time there was a case of similar infection in hospital which was also a sequel of an amputation, but not a refrigeration amputation. Refrigeration has not increased this form of infection despite fears to the contrary (Kirz, 1944). Cultures were made from ten consecutive specimens of ice, but in no case were pathogenic bacilli isolated aerobically or anaerobically.

Miscellaneous Complications.—Two cases are noted in the Table as resulting in death from renal failure. At necropsy both cases showed advanced renal disease of long duration. No ill effects were observed from the use of the tourniquet or cold temperatures. There were no cases of thrombosis or embolism, nor were there any cold sensitive stumps as might perhaps have been anticipated. The skin of the amputated leg sometimes showed slight scaling when refrigeration had been prolonged for some days, but only the most superficial of the epidermal layers were involved.

Discussion

Since the first large-scale trial of refrigeration anaesthesia in America in 1941 (Crossman *et al.*, 1942) many independent clinical reports have been published abroad with a uniform expression of satisfaction at the reduction of mortality in poor-risk cases (Batalha, 1946; Cayford and Pretty, 1945; Hinchey, 1944; Lobachev, 1946; McElvenny, 1942; Massie, 1944; Mock and Mock, 1943; O'Neil, 1944; Richards, 1944; Yudin, 1945a, 1945b).

The very foundation of the technique is the application of cold temperatures with or without a tourniquet, and since these have been regarded with suspicion and disapproval a brief discussion is necessary.

The temperatures used in refrigeration anaesthesia are not freezing temperatures. They are never less than 0° C. The skin, by virtue of its capacity for supercooling, does not freeze unless the temperature is below -2.2° C. (Lewis and Love, 1926; Lewis, 1941). Admittedly moisture on the skin reduces the degree of supercooling, but freezing never occurs at 0° C. Cooling the skin to 0° C., or even supercooling to a considerable extent, very rarely causes injury (Lewis and Love, 1926). Temperatures resulting in frost-bite are much lower than those used in refrigeration

anaesthesia (Lewis, 1941; Greene, 1941). Immersion-foot lesions have been reported in survivors adrift for several days in warm sea-water (White and Scoville, 1945). In these circumstances, however, the circulation and general resistance are depressed by starvation, exposure, and possibly other injuries. Cramped or dependent positions, tight clothing, or other local factors embarrass the circulation still further. Finally, the deleterious effects of moisture in the causation of thermal damage are implied in the term "immersion." In refrigeration anaesthesia the general condition is maintained throughout so far as is practicable; the circulation within the limb is either completely stopped or is maintained at maximum efficiency; the unfavourable effects of moisture have not been observed with cracked ice draining freely.

The tourniquet is not essential for adequate anaesthesia (Cayford and Pretty, 1945), but its employment is advised for the following reasons: (a) When the inflow of warm blood is stopped the tissues can be chilled thoroughly and rapidly. Radical lowering of the temperature in a limb is practically only obtainable with a tourniquet, unless there is gross pathological obstruction of the arteries. With the thorough reduction of the temperature that is possible with a tourniquet all protoplasmic activity is abolished; primary shock is non-existent in amputations on such limbs. (b) When the return flow of cold blood is stopped reflex vasoconstriction is prevented in various parts of the body. This might be harmful in cases of Buerger's disease and other conditions resulting in ischaemia to a part (Neller and Schmidt, 1944).

But the tourniquet is regarded with misgivings and its prolonged application with concern. The correctly applied tourniquet, however, is not as dangerous as believed (Allen, 1945c). The tourniquet that serves best is good soft rubber tubing (Allen, 1944); hard inelastic materials create trouble either through looseness or through crushing. In the leg, unlike the thin upper limb, very strong tension seems to cause no harm. The time factor has been greatly misjudged by surgeons in the past. Hinman (1945) has reviewed the factors that limit the duration of application. The most important of these are anoxia in the tissues distal to the tourniquet and shock on its removal. But the cold temperatures reduce to a minimum the oxygen requirements, and amputation of the distal tissues largely avoids the danger of shock (Allen, 1938, 1939, 1944, 1945a, 1945b; Allen *et al.*, 1942a, 1942b, 1943; Brooks and Duncan, 1940; Wilson and Roome, 1936).

The most convincing evidence, both experimental and practical, supports refrigeration as a valuable form of anaesthesia and as a sure means of excluding a constitutional depressant (Mock, 1946). However, I do not agree with those who claim that all patients to be subjected to amputation should be so anaesthetized (Crossman and Allen, 1946). Anaesthetists in England have attained a remarkably high standard, and few patients suffer from anaesthetics skilfully administered. Nevertheless it is submitted that there are certain bad-risk cases in which life can be saved by resort to this technique.

Summary

Twenty-five consecutive cases of amputation under refrigeration anaesthesia are reported. The details of individual cases are provided in the Table.

The method used in this series is described. The simplest technique has been used.

Bronchopneumonia was the most frequent complication.

Recent opinions concerning the effects of cold temperatures and tourniquet application are discussed.

It is submitted that this method be reserved for certain bad-risk cases.

I wish to express my gratitude to the Royal Melbourne Hospital Committee for permission to publish this series and to Mr. J. B. Turner and Mr. Paul Jones for allowing me to study their cases.

REFERENCES

- Allen, F. M. (1938). *Amer. J. Surg.*, **41**, 192.
 — (1939). *Ibid.*, **45**, 459.
 — (1944). *Canad. med. Ass. J.*, **51**, 220.
 — (1945a). *Amer. J. Surg.*, **68**, 170.
 — (1945b). *Curr. Res. Anesth. Analges.*, **24**, 51.
 — (1945c). *J. Amer. med. Ass.*, **129**, 980.
 — et al. (1942a). *Curr. Res. Anesth. Analges.*, **21**, 241.
 — (1942b). *Arch. Physical Ther.*, **23**, 711.
 — (1943). *Curr. Res. Anesth. Analges.*, **22**, 264.
 Batalha, E. (1946). *Int. Abstr. Surg.*, **83**, 499.
 Brooks, B., and Duncan, G. W. (1940). *Ann. Surg.*, **112**, 130.
 Cayford, E. H., and Pretty, H. G. (1945). *Ibid.*, **121**, 157.
 Crossman, L. W., Ruggiero, W. F., Hurley, V., and Allen, F. M. (1942). *Arch. Surg.*, **44**, 139.
 — and Allen, F. M. (1946). *J. Amer. med. Ass.*, **130**, 185.
 Greene, R. (1941). *Lancet*, **2**, 689.
 Hinchey, P. R. (1944). *New Engl. J. Med.*, **230**, 63.
 Hinman, F. (1945). *Int. Abstr. Surg.*, **81**, 357.
 Kirz, E. (1944). *British Medical Journal*, **2**, 662.
 Lewis, Sir Thomas (1941). *British Medical Journal*, **2**, 795.
 Lewis, T., and Love, W. S. (1926). *Heart*, **13**, 27.
 Lobachev, S. V. (1946). *Curr. Res. Anesth. Analges.*, **25**, 22.
 McElvenny, R. T. (1942). *Amer. J. Surg.*, **58**, 100.
 Massie, F. M. (1944). *Sth. Med. J.*, **37**, 1.
 Mock, H. E. (1946). *J. Amer. med. Ass.*, **130**, 187.
 — and Mock, H. E. (1943). *Ibid.*, **123**, 13.
 Neller, J. L., and Schmidt, E. R. (1944). *Wisconsin med. J.*, **43**, 936.
 O'Neil, E. E. (1944). *New Engl. J. Med.*, **230**, 209.
 Perlow, S. (1944). *Nav. Med. Bull.*, Wash., **42**, 433.
 Richards, V. (1944). *Ann. Surg.*, **119**, 178.
 White, J. C., and Scoville, W. B. (1945). *New Engl. J. Med.*, **232**, 415.
 Wilson, H., and Roome, N. W. (1936). *Arch. Surg.*, **32**, 334.
 Yudin, S. S. (1945a). *Curr. Res. Anesth. Analges.*, **24**, 216.
 — (1945b). *Ibid.*, **24**, 252.

PRIMARY MALARIA IN LONDON CHILDREN

REPORT OF TWO CASES

BY

C. BLAXLAND LEVICK, M.B., F.R.C.P.

Physician, St. George's Hospital and Victoria Hospital for Children

AND

M. E. MacGREGOR, M.D., M.R.C.P.

Paediatric Chief Assistant, St. George's Hospital and Victoria Hospital for Children

Two cases of primary malaria in London children have been discovered in one out-patient clinic within the space of a few months. They are reported now as a reminder of the necessity for keeping this diagnosis in mind when confronted with cases of unexplained fever in children who have never been out of this country.

Case 1

A girl aged 6 attended the Victoria Hospital for Children as an out-patient on May 9, 1946, with a history of loss of appetite and a heavy head cold for one week, nocturnal delirium for 48 hours, and vomiting, fever, and headache for 24 hours. At the time of examination a severe rigor was taking place; the oral temperature was 102° F. (38.9° C.), but there were no other abnormal physical signs. She was admitted to hospital immediately. Her temperature remained about 100° F. (37.8° C.) for the first 48 hours after admission, then fell to normal for 12 hours, and thereafter showed spikes of 100°–104° F. (37.8°–40° C.) on alternate days; each rise occurred between 1 and 8 p.m. and was preceded by a rigor lasting about half an hour. She complained of frontal headache and became confused at the height of each paroxysm. Between attacks her temperature was normal and she was alert and well. At first all investigations, including three thin blood films, proved negative. The spleen tip became palpable on May 24. Quinine sul-

phate, 15 gr. (1 g.) daily, was started empirically the same day and the pyrexia ceased at once. Quinine was discontinued after 10 days, but on June 6, coincident with another rigor and a pyrexia of 104° F., many benign tertian ring forms were demonstrated for the first time in a thin blood film. Quinine bihydrochloride, 12 gr. (0.8 g.) daily, with pamaquin, 15 mg. daily, was given for another ten days. Progress was uneventful apart from one sharp rise of temperature on June 12. A blood film on the 14th showed many red cells containing Schüffner's dots, but no parasites were seen then or thereafter. The spleen could not be felt after June 10. After two more courses of quinine, the patient returned home on Aug. 15 and remained symptom-free until Dec. 10, when her original symptoms were repeated. On Dec. 17 she was readmitted to hospital. A course of quinine and pamaquin was administered, and she has since continued free from symptoms.

Her past medical history revealed nothing of importance. The father, a Cypriot, came to England from Cyprus 15 years ago, and is now a waiter in London. The mother has never been out of England. Both parents and two siblings are well; no history of malaria was ascertained. There has been no contact with any relative from overseas. The patient remained in Worcestershire from infancy until September, 1945 (during the war there was a large American military camp near her dwelling-place), went to Welling, Kent, for six months, and then moved to Pimlico, where the family now live.

The medical officer of health for the City of Westminster states that in the past year, apart from relapsed cases in military personnel treated in hospital, no other case of malaria has been reported to him.

Case 2

A boy aged 2 was seen as an out-patient at the Victoria Hospital for Children on Sept. 26, 1946, with three weeks' history of a daily shivering attack between 1 and 5 p.m., not missing a single day. During these he was stated to be at first cold, blue, and shivering; then burning hot; then sweating; and, finally, half an hour later, quite well again, the whole attack lasting about three hours. There was no vomiting.

He was admitted to hospital immediately, and was afebrile. He was a well-nourished child of 28 lb. (12.7 kg.), with shadowing under the eyes and a yellowish pallor of the skin. The spleen was palpable two fingerbreadths below the costal margin. There were no other abnormal physical signs. At 3 p.m. on the day of admission and on the day following he had a rigor, the temperature rising to 103° F. (39.4° C.). A large number of benign tertian malarial ring forms were seen in a thin blood film on Sept. 27. A blood count showed: red cells, 3,000,000 per c.mm.; haemoglobin, 42%; white cells, 8,400 (polymorphs, 50%; lymphocytes, 45%; monocytes, 3%). He responded well to oral quinine and has had no recurrence since being discharged home on Oct. 23.

The patient is an only child. The parents are separated; neither has ever been abroad. The father was in the R.A.F., and was under investigation in 1941 for unexplained fever; he still has occasional fever, but has refused to allow examination of a blood film. Father and child were in close contact during the first six months of 1946. Apart from a grandfather, who had malaria in 1918 without subsequent relapse, there is no known contact with a malarial case. His only previous illness was bronchitis. He was born in Scarborough and was taken to London when 2 weeks old, to Bristol when he was 6 weeks, and back to London in October, 1945. Since then he has lived in Chelsea, apart from a visit in August, 1946, to the Isle of Sheppey, where the whole family were severely bitten by mosquitoes.

The medical officer of health for Chelsea states that no other case of primary malaria has been reported in his area during 1946, and the medical officer of health for the Isle of Sheppey gives the same report. Anopheline mosquitoes, however, are known to be present in the Isle of Sheppey.

Discussion

Unfortunately the source of infection is not absolutely clear in either case. It seems unlikely that primary malaria would remain symptomless for more than a week or two in a child in England. Therefore infection must have occurred