should be most grateful if they would send me full particulars of the cases and ask the patient to communicate with: The Blood Transfusion Officer, University College Hospital, Gower Street, London, W.C.1 (Euston 5050).—I am, etc.,

London, W.1.

W. N. GOLDSMITH.

Water and Salt Depletion

SIR,—There are some reactions affecting the distribution of salt which seem to need further study. After the masterly presentation of the role of salt in the body by Dr. H. L. Marriott (Feb. 15, March 8 and 15) one hesitates to intervene and can only plead interest as an apology. The reactions in question involve the little-explored field of the absorption thresholds of colloids for different concentrations of electrolytes. To illustrate: Gelatin placed in cold water swells. If salt be added gradually the gelatin continues to swell until a critical point is reached; then if more salt be added it will abstract water from the gelatin. Finally the salters' process is reached, used for centuries to cure skins. The significant points are that for each concentration of the electrolyte there is a different degree of absorption, and when this is active the force developed against resistance is very great. This pressure is masked in the body by elasticity in the tissues. In an organ with a capsule, like the kidney, it is greater; the rock is split. Evidently the force of colloidal absorption cannot be ignored.

Its action in the body is shown when oxyhaemoglobin is converted into the carboxy form. This causes the venous blood to absorb from 5 to 10% of its bulk of water, aiding the venous circulation and that of the tissue fluid. The water is released in the lung and excreted mainly by the kidney. That the absorptive capacity of the colloid is not exhausted is shown by the fact that in cases of suffocation the bulk of the venous blood is increased by 30%.

The typical cell is stated to retain potassium and phosphorus, but contains little salt although bathed constantly by a solution of it. The differences in the other types of cell must be due to some extent to the colloids they contain. The admission cells of the kidney pass most electrolytes in solution but refuse feebly ionizable proteins and neutral glucose. The conservation cells of the organ absorb water and salt, which they return to the blood freely, but refuse all other constituents of primary urine. Why does the colloid of one cell combine with potassium while another refuses? Why does the earth retain potassium while salt is washed into the sea? Apparent suspension of the laws of osmosis is evident in different reactions. During life the cell excretes actively but when dead ceases to do so but absorbs avidly. Why is osmosis active in dead cells but not in the living? Why does the cancer cell absorb so rapidly as to cause pressure which favours cell division?

If a red blood cell be filled with molecules the extent of its internal surface can be expressed in square metres. As electronic change is interatomic rather than between molecules, the area at which surface interchange, the base of chemical action, can take place is very great. Considering the diversity of action in the cell, which is so small that we know but little about it, it must be the seat of intense activity, much of which must be maintained by colloidal surface. The little-explored field of colloid absorption does seem to need investigation.—I am, etc.,

Montreal.

RICHARD KERRY.

Prognosis in Acute Nephritis

SIR,—Several statements are made in the interesting survey (April 5, p. 457) of Rudebeck's important paper on the prognosis in acute nephritis that seem to me to require qualification and upon which I would venture to comment.

After remarking that the prognosis must be guarded if residual symptoms are present on discharge from hospital (with which I agree), it is stated that if they (i.e., residual symptoms) persist for more than one year, or are severe, chronic nephritis may confidently be awaited. With this I cannot agree. In the paper by me cited in the annotation and by Rudebeck I referred to 586 patients with war nephritis in whom residual symptoms persisted for from one to ten years after the onset, all of whom eventually recovered completely. While the majority cleared up spontaneously in the earlier part of this

period, in 14 patients residual signs finally disappeared only in the ninth and tenth years after the onset.

It is, I believe, generally agreed that the prognosis in war nephritis differed little if at all from that in glomerulo-nephritis in civilians of comparable ages. In the past twenty-five years I have certainly seen quite a number of civilian patients of all ages in whom residual signs persisted for years who eventually recovered completely as judged by standards not less rigorous than those used by Rudebeck. On the other hand the proportion of patients suffering from latent or, as I prefer to call it, "unresolved" nephritis that ultimately recover is undoubtedly small. Those who do recover are fortunate exceptions. Nor in my experience is it correct to assert or to imply that chronic nephritis inevitably ensues when residual symptoms are severe, for I have often seen patients recover completely with appropriate treatment who were suffering from massive albuminuria with anasarca and ascites that had persisted for more than a year after the onset of the disease and before treatment was instituted.

It is not necessarily, in my view, the severity of the residual symptoms that determines the issue, but their nature. When they include hypertension or diminished renal function, however slight, and these persist for a year, and often for less than a year, it is a very different story. It is doubtful if complete recovery ever then occurs. I believe that hope of eventual complete recovery should never be abandoned however long residual signs persist, provided these do not include hypertension or diminished renal function; especially the former. The continuance for more than three or four months after the onset of considerable and undiminishing numbers of red blood cells in the urine is also in my experience a sign of grave import.

The second point to which I would refer concerns the effects of prolonged bed rest on the ultimate prognosis in this disease. In referring to Rudebeck's conclusions it is stated in the annotation that "it could not be proved that the prognosis was any better if the patient was kept at rest (i.e., in bed) for long periods." On consulting the original paper, however, I find that only 50 of the patients in Rudebeck's series were confined to bed for more than ninety days and only 7 for between six and nine months. Seven patients in the series were regarded as cured on discharge (not those who were confined to bed for the longest periods), and 43 were classified as not completely recovered. Concerning the latter Rudebeck remarks: "... among the non-recovered cases we also meet with many in which bed rest may be designated as insufficient." I entirely agree.

Observations extending over many years, but particularly during the past five years in the Renal Unit at Pembury, where it is possible to keep patients in bed as long as is considered necessary, indicate that complete recovery from unresolved nephritis results more often than is commonly supposed from treatment in which prolonged confinement to bed is probably the most important factor. In many patients six to nine months' carefully regulated bed rest is not the maximum but the minimum period required to achieve success. So far as I am aware the potentialities of prolonged and accurately controlled bed rest in renal diseases have not previously been systematically exploited on any adequate scale. The effects are, of course, quite different from those seen in patients who languish in bed in their own homes or in hospitals where bed rest is prescribed more or less as a matter of routine. Our experience at Pembury, where the objective evaluation of the effects of this measure has been a main line of research for some years, indicates that it may well prove to be as important in the treatment of unresolved nephritis and some other forms of renal disease as it is in pulmonary tuberculosis, for example. That prolonged and properly controlled bed rest for from six to fifteen or twenty months is sometimes the only means of procuring complete and lasting recovery in unresolved nephritis I have no doubt, though there is little to suggest that this measure will prove successful in more than a small proportion of these cases. In the absence of any speedier or better method of treatment it seems right to urge that patients suffering from unresolved nephritis should be given the chance of permanent recovery, however slender, that adequate bed rest occasionally offers.—I am, etc.,

Pembury, Kent.

A. A. OSMAN.