quently observed after cortical mastoidectomy performed by the post-aural route has not been found following this endaural technique. The lesion in the cases in question was acute or subacute mastoiditis; the ear drums and wounds were healed within a week of operation.

Convalescence appeared to be aided by pre- and post-operative systemic penicillin treatment.—I am, etc.,

Warwick.

J. A. HARPMAN.

Treatment of Acute Mastitis

SIR,—From the beginning of August until the end of October I encountered twelve cases of acute mastitis in thirty-two consecutive midwifery cases of which half were confined at the local maternity hospital. Cases showed evidence of acute inflammation of a breast from the 7th to the 21st day after confinement, local inflammation, high fever, and rigors. All cases except two resolved without surgical interference with penicillin *cum* stilboestrol therapy: 100,000 units twice daily and stilboestrol 1 mg. orally t.i.d. 600,000 units was administered in nine cases, 1,000,000 in one case. In the two remaining cases evidence of abscess was evident on the 3rd day of treatment. Both were dealt with surgically, and penicillin was continued until temperature and local condition indicated that inflammatory mischief was well controlled. These two received 1,000,000 units.

Occurring so late the infection was deemed to be not of haematogenous origin, while on the other hand careful investigation failed to reveal evidence of case-to-case infection. Moreover, inquiry in domiciliary practice showed that cases were common in the district in the practices of different midwives. The contingency of the epidemic is therefore difficult to explain.

The following preventive treatment appears to have been successful, as no case has occurred in my practice since it has been carried out as routine. The nipple is cleansed with warm normal saline and dried with sterile cotton-wool. It is then smeared thinly over with fresh penicillin cream. I have great hopes that many cases of mastitis can successfully resume breast feeding after treatment has been successfully completed, but so far I have failed to persuade any mother to make the attempt.—I am, etc.,

Fraserburgh, Aberdeenshire.

J. MACLEOD.

Poisoning by "Lethane" Insecticide

SIR,—In the article by Dr. C. V. Harrison (May 24, p. 722) the composition of lethane 384 special is given as: "12.5% of lethane 384 (N-butyl carbital thiocyanate), 37.5% of lauryl thiocyanate, [etc.]." So far as my information goes this is incorrect. I believe the composition to be (as stated in the paper by Prof. Buxton and myself in the *Journal* of April 11, 1942, p. 464) as follows: 12.5% N-butyl carbital thiocyanate (or, alternatively, *beta*-butoxy *beta*'-thiocyano diethyl ether), 37.5% *beta*-thiocyanoethyl laurate, 50% refined paraffin.

The main constituent (*beta*-thiocyanoethyl laurate) was not investigated by Cameron, G. R., Doniger, C. R., and Hughes, A. W. M. (1939), *J. Path. Bact.*, 49, 363, but some data regarding its toxicity to mammals are given by Main, R. J., and Haag, H. B. (1942), *Industr. Med.*, 11, 531. The M.L. doses found for intraperitoneal and subcutaneous injections to guinea-pigs and rats were 1.48 and 4.3 ml./kg. These figures fall into the range given for lauryl thiocyanate, and the error (if it be such) does not necessarily invalidate Dr. Harrison's conclusions. Nevertheless it may be well to publish this correction unless he has information that the composition of lethane special has been changed.—I am, etc.,

London, W.C.1.

J. R. BUSVINE.

"Tuberculous"

SIR,—I should like to draw attention to the all too frequent and improper use of the word "tubercular" when "tuberculous" is what is really meant. "Tubercular" is a general term used to describe any condition characterized by rounded nodules—for instance, a new growth—and exostoses might be described as tubercular in appearance when they are nodular. "Tuberculous" has a limited and special application; it is the adjective which should be used instead of "tubercular" to describe a condition due to tuberculosis. One should say a tuberculous (not a tubercular) lesion, a tuberculous (not a

tubercular) abscess, when it is intended to convey that these conditions are due to tuberculosis. The distinction may appear to be meticulous, but precision in language makes for clarity of thought and exposition, and the reader of an article is entitled to expect this from the writer.—I am, etc.,

London, W.14.

J. M. HAMILL.

Sir Joseph Barcroft and Drugs in Asphyxia

SIR,—I am sorry that my paper on the use of drugs in asphyxia appeared by some means as a letter in the *Journal* (June 7, p. 825). It was originally prepared for the use of a medical committee in I.C.I., and it was my intention, before it appeared in print, to preface it with a paragraph explaining the part that the late Sir Joseph Barcroft had in its composition. As it stands it might well be the theorizing of a medical scribbler. In fact it was the culmination of a correspondence over many years, and it incorporated several alterations made by Sir Joseph himself. As the matter is one of great practical as well as theoretical importance I think it is only right that it should be known that my "letter" had this authoritative backing. Sir Joseph's letter returning it (with the caveat that he would not personally like to be responsible for giving strychnine to infants) was one of the last he wrote, and I should like to take this opportunity of recording that his interest and help in the work of a general practitioner who did not happen to have been one of his pupils were as keen and as generous as in that of the many research workers who did have that privilege.—I am, etc.,

Winsford, Cheshire.

W. N. LEAK.

State Medical Service in New Zealand

SIR,—In response to Sir William Fletcher Shaw's letter (May 31, p. 782) I would point out that my quotation from Mr. Porritt's article in *St. Mary's Hospital Gazette* contained the whole of his comments on medical service. I must dissent from Sir William's description of *St. Mary's Hospital Gazette* as "almost a private publication"; Sir William will find a copy, I believe, in every medical library. That Mr. Porritt's article was not intended to be a private communication is perhaps shown by the introduction to the paragraph on medical services, which runs as follows:

"This leads me to a few words on things medical in New Zealand. For ten years the country has had a partial State medical service and a dispassionate appraisal of this is perhaps not without interest at the present juncture in this country."

I would also point out that Mr. Porritt, as a native of New Zealand, would perhaps be in a better position than Sir William to appraise the present position in New Zealand as compared with that position when he left the country 23 years ago. But it so happens that Mr. Porritt's conclusions, to which Sir William takes exception, are strongly fortified by a much fuller exposition furnished by Mr. Douglas Robb in a recent little book, *Health Reform in New Zealand*, sent to me for review. Mr. Robb is a distinguished consulting surgeon, a member of the Council of Auckland University College and of the New Zealand Medical Council. He criticizes the absence of arrangements for postgraduate teaching and research. He remarks that the "impetus for introduction of the State service was a political one and the conception of the details of the service was almost 100% political—few medical men appear to have been consulted as such." While recognizing the increased financial prosperity of general practitioners, he states his belief that "the service has been accompanied by a steady demoralization of the profession." His sense of general dissatisfaction with the present medical position is summed up in his request, to urge which is the real purpose of his book, for "the appointment of a Health Service Commission to inquire into all the aspects of the subject, with a view to plans for the future." This suggestion is of particular interest to me as I begged the Prime Minister in a Parliamentary Question (*Hansard*, Jan. 21, 1941) to appoint a Royal Commission which would take evidence from all sections of the community concerned with the Health Service before any definite steps were taken which would revolutionize existing arrangements. This request was refused, but was repeated later by the Negotiating Committee of the medical profession, and again refused.—I am, etc.,

House of Commons.

E. GRAHAM-LITTLE.