of Health Education in the headquarters office, to a Bureau of Legal Medicine and Medical Legislation, to the publication of *Hygeia*, the health magazine of the American Medical Association which now has a circulation in excess of 200,000 subscribers, and to other agencies. *Hygeia*, the health magazine, is now published as a part of the editorial functions of the Association.

As the interest grew in the field of industrial medicine, a Council on Industrial Health was established.

The most recent of the Councils is the Council on Medical Service, which is concerned with the development of voluntary insurance against the costs of hospitalization and of sickness. In association with this council there is a Bureau of Medical Economic Research and the information bureau maintained in Washington, D.C., the national capital.

Another bureau of the Association is the Bureau of Exhibits, which arranges not only for the scientific exhibit at the annual sessions of the American Medical Association but for similar exhibits at the meetings of the State Medical Societies and at meetings of voluntary philanthropic organizations in the field of health and other exhibits of medical interest. It has taken part in the encouragement of the establishment of health museums, of which there are now about six in the United States.

Quackery

Quackery is perennial. The charlatans have preyed on the public in the field of disease and health since the beginning of time. As the *Journal* began to gain prestige it undertook to expose from time to time various charlatans who had achieved great followings in the United States. As a result of these exposés the *Journal* and its editor have been sued for libel—the total number of such suits approaching 40 and the total amount of damages sought being more than \$30,000,000. The *Journal* has gone into court at least six times, since it has never compromised or settled a libel suit. In only one instance did it ever have an unsatisfactory verdict, and in that instance the award to the plaintiff, who was a manufacturer of an alcoholic nostrum for women, was one cent in damages.

Research

Early in its career the American Medical Association offered prizes for essays and other literary contributions to the advancement of medical knowledge. As its funds began to accumulate, the number and the value of these prizes increased. As each of the councils was established, funds were made available for encouraging research, and the Association established also a Committee on Scientific Research, which makes grants to medical investigators. The Association has spent more than \$1,500,000 for such research during the last forty years and has set aside \$2,000,000, the income of which is regularly used for research.

The single room with four employees which represented the headquarters of the American Medical Association in 1899 has grown to a building of half a block in width and almost a block in length, in which more than 700 employees carry on their activities. For these employees the Association has established hospitalization insurance and insurance against the costs of surgery or catastrophic illness.

Early in its career the American Medical Association recognized the importance of the State in the development of preventive medicine and public health. More than any other organization it has the credit for the development of the State Health Departments and of the United States Public Health Service. It has co-operated with every voluntary health agency in assuring adequate return for

the funds contributed by the people in promoting research and in education both of the public and of the medical profession regarding tuberculosis, cancer, blindness, deafness, paralysis, heart disease, and the work of the Red Cross in disaster and catastrophe. Some of its special campaigns have led to the elimination of the hazards of fireworks and gunpowder in celebration of the Fourth of July. It has led in the campaign against motor accidents and the provision of first aid when such accidents occur. It has carefully recorded the progress of the campaigns against death from typhoid and diphtheria. It has stood as the bulwark of medical science against the vast army of the ignorant, the stupid, and the fanatic, the antivivisectionists, the antivaccinationists, the antimicrobial science groups. All of these functions it has carried on with never an appeal for funds from any agency outside the medical profession itself, never an appeal to any other agency than the medical profession itself to eliminate the evils and misrepresentations. Always it has been the pride of Medicine that it has cleaned its own house; that it has recognized its obligations; that it has met every crisis in its career with its eyes set on greater goals for the good of mankind than have been its objectives in the past.

THE SULPHONE TREATMENT OF LEPROSY

BY

ERNEST MUIR, M.D., F.R.C.S.Ed.

Medical Secretary, The British Empire Leprosy Relief Association

The drugs commonly known as sulphones are derivatives of diaminodiphenyl sulphone.

It was their antibiotic effect in controlling the growth of *M. tuberculosis* in experimental animals and *in vitro* (Feldman *et al.*, 1942; Petter and Prenzlau, 1944) that first suggested trial in leprosy. Strangely, they have so far shown much more evidence of usefulness in leprosy than in tuberculosis.

The sulphone derivatives which have so far been tried in leprosy are "promin" (U.S.A., "promin"; in England, "promanide") (diaminodiphenyl sulphone-*nn*-didextrose sulphonate), "diasone" (disodiumformaldehyde sulphonylate of diaminodiphenyl sulphone), "promizol" (2,4'-diamino-5-thiasolylphenyl sulphone), and "sulphetrone" (tetrasodium phenylpropylamino-diphenyl-sulphone tetrasulphonate). Promin was first used in the National Leprosarium, Carville, U.S.A. (Faget *et al.*, 1943). It was found to be excessively toxic by oral administration but to be tolerated intravenously in daily doses up to 5 g. Following on the first published results on promin, Muir (1944) began a trial of diasone in Trinidad. Fearing that diasone might be toxic by mouth like promin, he first gave it intravenously, but later found that it was well tolerated orally in daily doses up to 2 g. A more recent sulphone derivative, promizole, is considered by the Carville workers (Faget, Pogge, and Johansen, 1946) to give possibly even better results than the first two preparations, though it has not yet been tried out sufficiently. This can be given orally in daily doses up to 6 g. Lastly, sulphetrone (B.W. & Co.) is under trial, and the first reports indicate promising results.

The results obtained with these sulphones appear to be similar, though they may vary in the degree of toxicity, the amount of the effective dose, and the speed of results.

Mode of Administration

Suitable Type of Case.—It is the lepromatous or severe type of leprosy to which sulphone treatment has been applied—that is to say, the type which so far has been least amenable to treatment by chaulmoogra and other drugs. I have had no experience with sulphones in tuberculous and uncharacteristic cases, nor have I seen any published results in these types.

Toxic Effects.—Faget *et al.* (1943) mention the following toxic signs as occurring: decrease in blood cells, leucopenia, allergic dermatitis, allergic rhinitis, mild and ephemeral headaches and nausea, and lepra reaction including iridocyclitis. Further observation has, however, shown that with suitable dosage most of these can be avoided. In my experience anaemia (chiefly due to destruction of red cells) and increased reactionary exacerbations are the two toxic indications to be guarded against, and these are apt to occur principally in patients in poor general health and/or in an advanced stage of the lepromatous type, and in the initial stages of treatment. Before beginning treatment the patient should be examined for anaemia, and when this is present, as it often is in severe lepromatous cases, a preliminary course of full doses of ferrous sulphate or carbonate should be given.

Dosage.—My own experience has been chiefly with diasone (Muir, 1944, 1946), and the dosage here described is what is recommended in the use of this drug. Considerably higher doses have been used with promin and promizole (Faget *et al.*, 1943, 1946). It is advisable to begin with small doses in all cases, and gradually raise the amount according to the tolerance of the patient. In doing this the important indications are: the stage of the disease; the general condition of the patient, especially as regards febrile and focal reactionary signs; the condition of the blood; and the presence of anaemia. If the haemoglobin percentage is below 70 a preliminary course of iron should be given, and if this is not effective injections of liver extract should be added.

Diasone is generally made up in 5-gr. (0.32 g.) capsules or tablets. When the haemoglobin is at or over 70%, begin with 1, 2, or 3 tablets according to the general condition of the patient. This should be taken in one undivided dose, preferably an hour after food, so as to promote quick absorption and the highest blood concentration. Reactionary signs are not a contraindication to beginning the treatment, but are a warning not to raise the dose too rapidly. Whatever the initial dose, repeat it every second day for the first week unless they are signs of exacerbation. In strong early cases without septic or other complications one tablet may be added after the first week on the intermediate days and increased gradually till, after three or four weeks, the patient is taking 3 tablets daily for six days a week. It may, however, be a considerable time before a weaker or more advanced case reaches this dosage. The haemoglobin should be tested every week to begin with and iron (and if necessary liver extract) continued in all cases with a percentage below 80 or 90. A fall below 70% or the intercurrent of increased reaction calls for temporary suspension of diasone or diminution of the dose. When a dose of 3 tablets a day for six days a week has been reached it is well to suspend treatment for one week every month. Apparently when full dosage is suddenly resumed after this temporary stoppage the blood concentration rises to a higher level for a time. In patients who have improved

and are free from signs of anaemia and reaction the dose may be gradually raised to 6 tablets daily six days a week for three weeks a month—this being regarded as the maximum average dose, though it may be increased or diminished according to the body weight of the patient.

Effects of Treatment

These vary according to the stage of the disease. In a severe advanced lepromatous case with ulcerating leproma of the limbs, ulceration of the nasal and other mucous membranes with obstruction of breathing, and with severely affected eyes going on to blindness, the first effects, which often take place within a few weeks, are the healing of ulcers, clearing of the nose, and arrest of the eye condition with improved vision.

The next effect is seen in patients with chronic or sub-acute allergic reactions indicated by slight rise of temperature and/or the frequent appearance and subsidence of nodules and other inflammatory skin lesions; these symptoms gradually subside. At the same time permanent nodules and other raised skin elevations become slowly absorbed and flattened out, leaving marked wrinkling of the integument. This improvement generally requires a few months.

The third stage in recovery, and the only one in early or less advanced cases, is the gradual lessening of the number of bacilli found in sections and smears from the skin and in smears from the nose. Thus three-plus cases become two-plus and then one-plus, and one-plus cases become negative. The time required for this appears to vary with the advancement of the disease: some early cases have become negative in four to six months, but more advanced cases may require four, five, or even more years. One of the most striking signs of improvement is that seen in the affected eye and especially in the cornea, where, because of its transparency, the arrest and a certain amount of recession of the lepromatous infiltration can be observed with a lens or corneal microscope.

There is one important point on which evidence is not yet available—that is, whether or not a point is reached at which the type changes, the lepromatous case being transformed into a tuberculous one with a negative "lepromin" test becoming positive. Theoretically the lepromin test is negative in the lepromatous case because the antigen (i.e., lepra bacilli) are in such great excess of antibodies formed that an effective allergic reaction, which would destroy the bacilli, cannot take place. As the bacillary antigen becomes less and less, is a point reached at which an effective reaction can take place? If so, it would be indicated by a positive lepromin reaction, and we should have an additional factor helping to clear up the residual bacilli. I have found indications in one or two cases that this may occur to some extent, but further evidence is needed.

Length of Treatment

In any case, treatment should be continued at least until repeated bacterioscopic examinations have given negative results over a period of six months to two years, varying directly with the advance of the disease at the beginning of treatment, and with the length of treatment required to produce the first negative bacterioscopic results. Faget (1947) reports that "19 promin-treated patients have been discharged as disease-arrested following twelve consecutive months of negative bacterioscopy. Of this number three were under treatment for 1½ to 2 years, three from 2 to 3 years, six from 3 to 4 years, and seven from 4 to 5 years. There have been no relapses. . . . The period of

observation following arrest of the disease has varied from a few months to 2½ years."

Iodides have a specific effect in showing up concealed lepromatous foci. It is possible that their careful administration in cases that have become bacterioscopically negative may be of use in determining the length of further treatment required, and perhaps in speeding up the elimination of residual foci; but this matter still calls for very careful investigation.

Regulation of Established Treatment.—Particulars have been given of the dangers attending the initial treatment with diasone. But once the patient has taken his maximum dose for a few months without diminution of haemoglobin content or reactionary symptoms, and has made definite improvement as regards both leprosy and in his general health, the same precautionary methods cease to be necessary and treatment may be continued by an intelligent and reliable patient with a minimum of supervision. A visit by the doctor once a month is often all that is needed. This fact is particularly helpful to patients under domiciliary treatment.

Mode of Action

It is still uncertain how the sulphones act. Fite and Gemar (1946), after examination of biopsy sections from 32 patients under promin, concluded that tissue changes are "atrophic in character with extremely slow and gradual lessening of numbers of organisms in the lesions to the point of final disappearance in 10 of 32 cases examined. . . . The important finding is that promin appears to eliminate bacillary infection of the blood vessels and blood stream, thereby preventing the formation of new lesions. The atrophy of focal lesions is also more apparent in areas with a more generous blood supply. The results indicate strongly that the best results may be expected in those cases in which treatment is begun in a comparatively early stage of the disease."

A satisfactory hypothesis is as follows. Ordinarily in lepromatous cases the cellular ingestion of bacilli and their multiplication in the cytoplasm result in the formation of "globi," the rounded masses of bacilli found in all typical lepromatous cases. The global bacilli gradually die and are transformed into a non-acid-fast lipid matrix which imbeds the still acid-fast bacilli. Thus in untreated cases the living bacillary element in a globus gradually becomes extinct, and its place is normally taken by fresh globi. It may be supposed that the sulphones act not by destroying bacilli but by preventing the multiplication of bacilli and the formation of fresh globi.

If this hypothesis is correct it would account for: (a) The finding of globi in smears from the nose and skin of treated patients, in which there are no acid-fast bacilli or in which the bacilli are thin and ghost-like (b) The long time required in the clearing up of bacilli in fairly advanced cases. This will be made clear when it is mentioned that human leprosy bacilli, killed by boiling and injected into rats, can still be found retaining their acid-fastness after 18 months.

The Future of Sulphone Treatment

The history of leprosy in the last 30 or 40 years is strewn with the wrecks of so-called "cures." Knowing this, one is hesitant to put forward claims for the sulphones which might fail to be confirmed. It will take years before we can say with any confidence what the final results are, whether a complete and lasting cure is possible, what is the relapse rate, and whether relapsed cases will yield to further courses of treatment.

Confidence is given, however, by the fact that experienced workers all over the world who have carried out the initial trials have without exception obtained favourable results such as they had never before found with other remedies. Many have reported definite improvement in practically all cases that have been under treatment for at least six months.

The history of a rising scale of effectiveness in other drugs, such as the sulphonamides, gives reason for hope that sulphone derivatives may be produced which will give even more rapid and effective results.

Sulphones and the Anti-leprosy Campaign

The Supply of the Drugs.—Few of the countries where leprosy is rife are wealthy, and the patients who require treatment most are as a rule the poorest of the poor. In a campaign against leprosy there is, therefore, need for a drug which is not only effective but which can be made available to the poorest patient in countries like Africa, India, and China.

Fortunately the raw material of the sulphones is not expensive, and although the outlay on manufacture must at first be considerable the cost should rapidly diminish with mass production. It is calculated that 1,000 5-gr. (0.32-g.) tablets of diasone should, on an average, suffice for one patient for one year, and the present cost of this is in the region of £6. If treatment is to be extended to a gradually increasing proportion of the one or two million lepromatous cases calculated to exist in the world, there will be a considerable call on the budgets of governments, and on the generosity of philanthropic bodies.

Effect on Control.—The two chief difficulties in the control of leprosy have been: (a) open infectious cases, through ignorance, fear, and shame, or want of opportunity, are not isolated until they have had a chance, often lasting several years, of spreading the disease to their relations and other contacts; (b) that the treatment available has been painful and hard to persist with over a long period and, even when persisted with, has in the majority of lepromatous cases been unable to arrest the disease.

It may be expected that once the effects of sulphones become widely known they will attract early cases which would otherwise have concealed the disease as long as possible. Thus to a gradually increasing extent the danger of infection should be cut off at its source.

Need of Personnel.—One of the chief difficulties in anti-leprosy work has always been the lack of doctors. In the medical profession there has been a prejudice against leprosy, either on account of fear or because it was considered that so little could be done for it. In Brazil the Government offers an extra allowance beyond the ordinary service rates. But the best work has always been done by those who have felt they have a vocation, and many such doctors and other workers, once they have overcome the prejudices which surround the subject of leprosy, have become enthralled by the interest of the work. Now that there is a clearer prospect of effective treatment and control, it is to be hoped that more doctors with a pioneering and venturesome (not to say philanthropic) spirit will offer to do this work.

Summary

The sulphones have been found of definite benefit in the severe or lepromatous type of leprosy, clearing up complications, causing a slow but steady diminution of the bacilli, and in some cases bringing about arrest of the disease.

The method of administration of diasone, its possible mode of action, and length of treatment are given in detail.

The effects of sulphone treatment on the campaign against leprosy, the supply of sulphones, and the need of suitable personnel are discussed.

REFERENCES

- Faget, G. H. (1947). *Int. J. Lep.*, **15**, 7.
 — Pogge, R. C., Johansen, F. A. (1946). *Pub. Hlth. Rep.*, Wash., **61**, 957.
 — — — Dinan, J. F., Prejean, B. M., and Eccles, G. C. (1943). *Ibid.*, **58**, 1729.
 Feldman, W. H., Hinshaw, H. C., and Moses, H. E. (1942). *Amer. Rev. Tuberc.*, **45**, 303.
 Fite, G. L., and Gemar, F. (1946). *Sth. med. J.*, Nashville, **39**, 277.
 Muir, E. (1944). *Int. J. Lep.*, **12**, 1.
 — (1946). *Leprosy Rev.*, **17**, 87.
 Petter, C. K., and Prenzlau, W. S. (1944). *Amer. Rev. Tuberc.*, **49**, 308.

TROPICAL EOSINOPHILIA IN EAST AFRICA

BY

H. T. H. WILSON, M.B., B.Chir., M.R.C.P.
(Late M.O., Colonial Medical Service)

The syndrome now generally known as tropical eosinophilia or eosinophilic bronchitis was first described by Frimodt-Møller and Barton (1940) in Southern India. Under the title "a pseudotuberculosis associated with eosinophilia" they published an account of a syndrome characterized by cough, fever, and loss of weight which ran a prolonged but comparatively benign course. There was a fairly constant similarity in the radiographs of these patients, the characteristic picture showing extensive mottling of both lung fields, resembling somewhat miliary tuberculosis or silicosis. Blood counts revealed a high degree of eosinophilia, and in over half the patients examined 5,000 or more eosinophils per c.mm. were found.

Weingarten (1943) recorded 81 cases of tropical eosinophilia. He found that expiratory dyspnoea was a prominent symptom and that the spleen was often enlarged. After trying various remedies without success he discovered that dramatic improvement resulted from a short course of neoarsphenamine. Further cases were described in India by Simeons (1943), Treu (1943), Menon (1945), and others, while a similar condition was reported from Ceylon by Carter, Wedd, and d'Abrera (1944), Carter and d'Abrera (1946b), and Soysa and Jayawardena (1945). The Ceylon workers discovered mites of various genera in the sputum of more than half their cases, and considered that tropical eosinophilia might "in part at least be explained on the basis of a mite infestation of the respiratory system." Treatment by inorganic arsenicals was found to be effective with neoarsphenamine and "acetylarsan" by injection, and acetarsol or carbarsone by mouth.

While tropical eosinophilia can no longer be regarded as a rare disease in India and Ceylon, its existence in Africa has not yet been generally recognized. Isolated cases have, however, been recorded by Parsons-Smith (1944) and Stephan (1946) in Cairo, while Ritchie (1944) described the condition in a Tanganyika native serving in the Middle East Forces and Hunter (1946) in a European recently returned from West Africa. The present communication describes an investigation carried out in Dar-es-Salaam, Tanganyika. Differential white counts were performed on a consecutive series of 34 patients attending the native civil hospital with symptoms of recurrent bronchitis or asthma. The patients were all males, and consisted of 29 Indians, four Africans, and one Arab. In 28 cases an eosinophilia of less than 20% was found, while in six cases the eosinophilia ranged from 40 to 78%. The latter group, and a seventh case which was referred on account of an unexplained eosinophilia, were submitted to detailed investigation. All were considered to be suffering from tropical eosinophilia;

six were treated successfully with arsenic, while the condition regressed spontaneously in the seventh case.

Clinical Findings

The characteristic picture appeared to be recurrent attacks of cough, wheezing, and low fever separated by intervals in which the only symptom was a spasm of unproductive coughing in the early hours of the morning. In four cases retrosternal pain was present, and three complained of acute irritation of the larynx and upper respiratory tract. After a variable number of attacks the intervals of freedom became shorter, and cough and dyspnoea caused increasing distress. Early cases showed no constitutional symptoms, but those who had been ill for several months complained of lassitude, anorexia, and loss of weight. Case VII conformed to the clinical type described by Frimodt-Møller and Barton as "pseudotuberculosis." That patient complained of chronic cough, pain in the chest, and occasional small haemoptyses without asthmatic attacks.

X-ray Findings.—In three cases no abnormality was seen. In Cases VI and VII there was diffuse mottling of both lung fields, characteristic of the "eosinophil lung pattern," and a suggestion of mottling was seen in the radiographs of Case IV. These cases gave a history of one year's, two years', and ten months' duration respectively.

Pathological Findings.—(a) *Haematological.*—An initial blood count was done on all cases by Dr. Vivarelli. The red cells varied from 4,650,000 to 5,150,000 per c.mm. and the haemoglobin from 94% to 110% (Lovibond)—comparatively high figures in a country where anaemia is prevalent. Leucocytosis was a constant finding, counts ranging from 11,250 to 34,000 cells per c.mm., with an eosinophilia of 52–78%. Except for an occasional eosinophil metamyelocyte no immature cells were seen; the granules varied in size and tended to be larger than normal; they frequently did not fill the cell, and left spaces like vacuoles in films stained by the Leishman method. Whitby and Britton (1942) noted a similar appearance of the cells in eosinophilic leukaemia. Differential counts were repeated two or three times weekly on 200–400 cells, and the percentage of vacuolated cells was seen to diminish in the course of treatment. The blood sedimentation rate was estimated by Westergren's method in three cases; in one case it was raised to 47 mm. after 1 hour, but in the other two cases it was within normal limits. Films from all patients were examined for filariae, but none were found.

(b) *Urine and Stool Examinations.*—No abnormality was found on naked-eye or microscopical examination of the urine, and specimens of stools were negative for parasites, except in Cases V and VI, in which a few ankylostome ova were discovered by concentration methods. In both cases a rise in the total eosinophils was found when the blood was re-examined after deworming.

(c) *Sputum.*—Specimens were repeatedly examined for tubercle bacilli, but none were found. Sputum from Cases II–VI was examined by Dr. Vivarelli for mites by the method described by Soysa and Jayawardena. Twenty-four-hour specimens were collected into sterile stoppered bottles, and the mucopurulent material was disintegrated with 10% potassium hydroxide. After it had cleared, enough formalin to give a 10% concentration was added and the liquid left for 24 hours. It was then centrifuged and the deposit systematically examined under the microscope. Four mites of *Tyroglyphus* and a few ova were found in specimens from Case II, one mite of *Tyroglyphus* in Case VI, and an unidentified mite in sputum from Case IV. No mites were found in sputum from Cases III and V.