setting up a local focus where staphylococci from the infected tooth socket lodged.

Case 5.—A naval rating aged 25 was admitted to hospital because of a sinus of the posterior chest wall following the incision of an abscess at this site eight months previously. A skiagram showed an increased density of the left side of the bodies of the third and fourth thoracic vertebrae, and the intervertebral space between these was narrowed. An oblique view showed beak formation of the

FIG. 4.—Lateral view in Case 3 seven months after the onset of backache, showing massive bony bridging.

lower border of the third dorsal vertebra. The adjacent heads and necks of the third and fourth ribs showed widening and irregular thickening, and a similar appearance was evident in the corresponding transverse processes. Examination of the pus from the sinus indicated a mixed staphylococcal and streptococcal infection, and biopsy of the sinus wall showed no evidence of tuberculous granulation tissue.

The records of this case are incomplete, but according to the patient he was feverish at the time of the dorsal backache. The x-ray appearances point to a pyogenic infection, although an original tuberculous infection with a secondary pyogenic one cannot be ruled out. It is in such a case that inoculation of a guinea-pig with sinus scrapings would be valuable, but this was not done owing to the patient's repatriation to Canada.

Summary

The aetiology, pathology, clinical signs and symptoms, and treatment of pyogenic osteomyelitis of the spine are described.

It is suggested that the disease is more common than has been thought.

It seems that many cases of osteomyelitis of the spine, labelled tuberculous, are in reality pyogenic in origin.

At least in adults the disease is less fatal than has been considered.

The notes of five cases are presented.

REFERENCES

- Browder, J., and Meyers, R. (1941). *Surgery*, 10, 296.
Butler, E. C. B., et al. (1941). *Lancet*, 1, 480.
Kulowski, J. (1936). *J. Bone Jt. Surg.*, 18, 343.
Turner, P. (1938). *Brit. J. Surg.*, 26, 71.
Wilensky, A. O. (1934). *Osteomyelitis*, Macmillan Company, New York.

TREATMENT OF OSTEOARTHRITIS BY LACTIC ACID INJECTION

A SERIES OF 26 CONSECUTIVE CASES IN GENERAL PRACTICE

BY

R. MAWSON, M.R.C.S., L.R.C.P.

Stimulated by the work of Grant Waugh on intra-articular injection of lactic acid with procaine into the hip we recently gave his treatment to a series of 26 patients. The series comprised 6 cases of arthritis of the hip, 18 of the knee, 2 of the wrist, and consisted of the following types: osteoarthritis of the elderly; post-traumatic; so-called "menopausal"; post-infective; and residual deformity stage of rheumatoid arthritis. The cases occurred in semirural general practice and were taken from all social strata in our valley community. Economic factors prevented serial radiographs and periodic estimation of the erythrocyte sedimentation rate, and the treatment was domiciliary in all instances but one.

Case records were kept as simple as possible, and notes of progress consisted of statements made by the patient and his relatives, together with my own observations. It was difficult always to assess evidence of progress or deterioration in objective terms except when the patient showed dramatically changed physical signs—e.g., doubled range of movement at the affected joint. A guide to progress in this disease is the distance that can be walked before and after treatment. This criterion was used in most cases. However, clinical medicine in general practice includes sundry observations of patients in the street, in shops, in the village dance hall, and in the public house. I should be more pleased to see an osteoarthritic at the end of a fish queue than a small change in her sedimentation rate. Grant Waugh's assessment of progress was based on the degree of overall disability, expressed as a percentage, taking 90 to 100% disabled to describe patients confined to bed, and 70 to 80% for those able to walk with considerable difficulty.

It is not easy to classify stages of recovery. So often psychological factors distort the clinical picture. It is well known that all forms of rheumatism and neurosis are often closely related, but it is not desirable here to add to the hothouse flavour of some of the observations made by certain workers in this field: "The cramped muscle reflects the cramped mind" is an illustration. Reference will be made subsequently to certain features of anxiety which obscure the slighter signs of recovery in this type of patient.

The clinical state before and after treatment was based on inquiry into the following: pain and tenderness; deformity and posture; anorexia and insomnia; maximum walking distance; stiffness and alteration of gait; effusion; mental distress.

Below are given a table showing results of treatment in the series of 26 and records of four typical cases.

Joints Affected	No. of Cases		Age Range	Av. No. of Injections	Results		
	M.	F.			Better	No Change	Worse
Hips ..	4	2	36-67	4	5	1	0
Knees ..	1	17	42-73	4	17	1	0
Wrists ..	0	2	—	2	1	1	0

Case Histories

Case 1.—Mr. D., a landworker aged 36, had suffered from osteoarthritis of the left hip for 7 years, the result of a fractured femoral neck in 1927 (4 months in plaster). His symptoms were pain, stiffness, and insomnia. Examination revealed quadriceps and gluteal wasting, concealed hip flexion, flexion 60°, extension *nil*, adduction and abduction very small, rotation *nil*. All movements were painful. Walking distance before treatment: 1/4 mile (400 metres) in great pain. He was on the point of giving up his employment. Treatment consisted of four injections. The first injection brought a dramatic improvement; the patient could not only walk easily, but was actually able to run—something he had not been able to attempt for several years. The third and fourth injections produced much pain, and work stopped for 10 weeks. He is now back at work, and can walk up to 4 to 5 miles (6.5 to 8 km.) a day. There is still pain in the joint, but he can tie his left shoe-lace. All movements have increased, particularly flexion—almost to 90°. There is no concealed hip flexion. A radiograph

FIG. 1.—Radiograph of Case 1, Feb. 14, 1946.

(Fig. 1) taken on Feb. 14, 1946, reveals loss of spacing, flattening of the head of the femur, sclerosis and broadening of the neck with buttress formation, suggesting an old injury to the neck of the femur and now showing changes of osteoarthritis. From the appearance this buttress should interfere with the movement of

FIG. 2.—Radiograph of Case 1 taken on March 21, 1946.

the hip-joint. It is not quite clear whether the buttress has fused with the lower part of the acetabulum. Fig. 2 shows the condition on March 21, 1946.

Case 2.—Miss P., a housekeeper aged 63, had her hips, knees, and wrists affected during a period of 9 years, the cause being rheumatoid arthritis. She had great pain, particularly in the right knee; and knee movements were almost *nil*. She suffered from loss of sleep and appetite, and had been taking eight aspirins daily. She could only hobble around with the aid of two sticks, and had not been outside the house for months. She received two injections in the right knee, which was manipulated under pentothal. There is now almost complete loss of pain in the knee, which shows greatly increased movement. She can carry a loaded tea-tray upstairs, and she can walk half a mile (800 metres) to the post office with the aid of a stick. Her general state—colour, appetite, mental outlook—is very much improved. She is almost weaned from salicylates.

Case 3.—Mr. W., a gardener aged 65, had his right hip-joint affected for 8 years. His symptoms before treatment were pain (which made him give up work), stiffness and diminished movement, and he could walk only about 200 yards. He was given two injections. Increased movements at the right hip resulted and there was far less pain. He resumed work after the second injection. Since then (March, 1946) he has worked 8 to 9 hours a day and is able to attend to his own garden in the evening.

Case 4.—Mrs. B., a housewife aged 69, had the shoulders, wrists, and small joints of the hands affected, especially the left wrist. The symptoms were pain, anorexia, insomnia, immobility of wrists, and wasting of the small muscles of the hands. She was unable to feed herself through inability to grip knife and fork. Her bed had been put downstairs because she could not manage the stairs. Her walking distance before treatment was *nil*. The patient received two injections in the left wrist. Movements at that joint began a few days after the first injection. She can now flex to 70° and dorsiflex 30°, and there is no pain or swelling. Her grip has improved, and she can feed herself and get upstairs.

Technique

From 15 to 20 ml. of a stable solution of 0.2% lactic acid with procaine, having a hydrogen-ion concentration of 5.2, is injected into and around the hip-joint on each occasion. A similar amount may be injected into the knee-joint, but the wrist-joint will only take amounts varying from 2 to 5 ml. Skin sterilization is effected by 1% "cetavlon" or 1:1,000 acriflavine. Skin, subcutis, muscle, and joint capsule are infiltrated with "novutox," the average amount required being 20 ml.; it is greatest in the case of the hip-joint in obese patients, and smallest in injection of the wrist. For the hip-joint a 6-in. (15 cm.) Record needle with wide bore and of high tensile strength is employed. For the other joints sturdy intramuscular needles will suffice.

Approach to the hip-joint is made by two methods. (i) *Anterior*: The needle is inserted midway between the great trochanter and the intersection of the femoral artery with Poupart's ligament; it is directed upwards, parallel with the femoral neck, inwards and backwards. (ii) *Posterior* (Fig. 3): The needle is inserted at a point just posterior to the great trochanter, and in an upward, inward, and forward direction.

In dealing with the knee-joint the needle is inserted 1 in. (2.5 cm.) to the outer side of the patella (Fig. 4).

Approach to the wrist-joint (Fig. 5) may be carried out either (i) immediately distal to the tip of the ulnar styloid process between flexor and extensor carpi ulnaris tendons, or (ii) dorsally between tendons of extensor pollicis longus and extensor indicis.

Miscellaneous Observations.—An intra-articular injection of the hip-joint requires a strong needle with a sharp short bevel, as the capsule in chronic cases of osteoarthritis may be up to 1/2 in. (1.25 cm.) in thickness. The needle must be inserted slowly without undue force. Once inside the joint the solution will run in easily. If much pressure is necessary the needle is probably outside the joint cavity embedded in the fibres of the capsule. In two of the cases under consideration pain was experienced at the time of the injection (which had required more pressure than usual), and this pain increased as the effect of the procaine diminished. Pain of a deep boring character persisted for several days and required the administration of the stronger barbiturates during that time. The thigh was held in the position of rest—internal rotation, semi-flexion, and adduction. After the intra-articular injection the patient is asked to put the joint through all its movements by himself.

Usually it is found that there is an immediate loss of local discomfort after the injection. Gentle passive manipulation of the joint follows the patient's own active movements. For the next 48 hours the patient is advised to remain in bed so that the joint shall not suffer weight-bearing. If the services of a physiotherapist are available the wasted muscles around the joint and elsewhere can be treated. A second injection is given after 7 days, following exactly the same plan as before. Subsequent injections, up to 6, are given if needed.

Manipulation under "Pentothal" Anaesthesia

In certain cases (from whose aetiology tuberculosis can with certainty be excluded) it may be necessary to manipulate the stiff painful joint under pentothal. This procedure was found to be followed by some pain and anxiety. The former responded to salicylates only, and was of short duration. The anxiety

FIG. 3.
Posterior approach to the hip-joint.

FIG. 4.
Approach to the knee-joint.

FIG. 5.
Approach to the wrist-joint.

has its origin in the belief—so commonly held by the chronic arthritic—that the affected joint will never move freely again. The patient's return to consciousness is awaited and the joint manipulated and flexed to its fullest extent, so that on opening his eyes the patient sees the degree of angulation possible in his joint. Quadriceps weakness makes it impossible for the patient to flex the hip completely at this stage, but there is always an improvement on the original degree of movement. Quadriceps drill, patient masseur, and encouragement from the family will do the rest.

Stages of Recovery

The first symptom to disappear is pain, followed by a changed mental outlook. To patients who previously were lethargic, and pessimistic new hope is given. Sleep and appetite improve. As atrophied muscles are stimulated by massage the patient

ventures to do more about the house, and gradually leave is taken of the bedroom. The difficulty of stairs is overcome, and provided progress is maintained the patient is soon in the garden. Fresh air and exercise enhance the sense of well-being, which in turn promotes recovery of the local condition.

Summary

A series of 26 cases of mixed forms of arthritis is reviewed after treatment by intra-articular acid injection. The technique of injection of hip, knee, and wrist is described.

Of the patients, 6 were wage-earners. Recovery was sufficient in all six to enable them to return to their previous employment.

Of the remaining 20, mostly housewives, 18 were able to resume or do much more housework, with considerable relief of symptoms. Of the other 2 cases, one declined to have more than one injection, for domestic reasons; the other, with active rheumatoid arthritis of the wrists, showed no improvement.

BIBLIOGRAPHY

- British Medical Journal*, 1946, 1, 134.
 British Orthopaedic Association Report, May 24, 1946.
 Miles, A., and Wilkie, D. P. D. (1936). *Operative Surgery*, Oxford Univ. Press, London.
 Waugh, W. G. (1936) *Lancet*, 2, 976.
 — (1938). *Ibid.*, 1, 487.
 — (1939). *Rheumatism*, London.
 — (1945). *British Medical Journal*, 1, 873.

DISAPPEARANCE OF SECONDARY SARCOMATOUS DEPOSITS IN THE LUNGS AFTER STILBOESTROL THERAPY

BY

J. GUNN ROBERTS, M.B., Ch.B.

Assistant, Surgical Unit, Welsh National School of Medicine

The beneficial effects of stilboestrol in the treatment of carcinoma of the prostate are well known, where not only the primary growth but also secondary growths in lung and bone have been kept under control and in several instances have disappeared completely. Less permanent and less dramatic effects have been observed in several cases of carcinoma of the breast treated similarly. Few successes have been claimed in any other type of growth.

In this communication a case of malignant endothelioma is recorded where secondary growths in the lung disappeared radiologically after stilboestrol therapy and where the patient was apparently free of active tumour two years later.

FIG. 1.—Skiagram of chest (June, 1944) showing multiple secondaries in both lung fields.