he states that I have attempted "to divorce it (the thalamic dysfunction syndrome) . . . from environmental factors"; whereas in my paper I stated clearly that "the thalamic dysfunction state may be regarded as a response of the organism to stresses of various kinds" (i.e., environmental stress, conflicts, toxins, and so on).

With regard to his remarks about the morphine and cocaine addiction syndromes, if Dr. Kirman were to go to the trouble of consulting any of the standard psychiatric textbooks dealing with the subject, he would find it stated that alkaloid addiction differs in a number of important respects from alcoholism. This last-named toxin, incidentally, is a cortical depressant rather than a true euphoriant of the type described in my article, in addition to being chemically and pharmacologically unrelated to the agents under consideration. Finally, while admitting "the limited value of present methods" (i.e., of psychiatric treatment), he is apparently unable to suggest any alternative or more effective form of treatment than that suggested in my article. In other words, his criticism is the purely destructive kind which questions everything but decides nothing, containing not one single constructive contribution to the problem under discussion. I should, of course, welcome any constructive suggestions or comments on the new concept of affective disorders put forward in my paper and the pharmacotherapy of these conditions. But it is a pity that Dr. Kirman did not take the trouble to read the paper through properly before erupting so impulsively into print.—I am, etc.,

London, S.E.6.

G. TAYLEUR STOCKINGS.

Bornholm Disease in the Tropics

SIR,—Only a few weeks before the outbreak of Bornholm disease at Aden described by Drs. W. M. Jamieson and D. M. Prinsley (July 12, p. 47) there was an outbreak of an unusual nature in a Naval barracks at Singapore (June, 1946). The distinctive features were as follows: (1) It was larger than any recorded to my knowledge (over 100 cases). (2) It was explosive—more than seventy of the cases occurring in the first twenty-four hours. (3) It was sharply limited to European ratings in the Naval establishment concerned, the majority of cases occurring among men who fed at one end of a mess hall, but who worked and slept in different places over a wide area. (4) The primary lesion appeared to us to be in the parietal pleura, and in several cases a well-marked pleural rub ensued. Features 2 and 3 do not seem consistent with a contact infection, and contamination of food was suspected, possibly by flies. Otherwise signs, symptoms, course, and complications were as normally described.

There had been an outbreak of poliomyelitis at Singapore earlier in the year, during which a few civilians admitted to the fever hospital were regarded as probably suffering from Bornholm disease, but previously this does not appear to have been recognized in Malaya. A Dutch colleague, however, stated that he was familiar with the symptoms in sporadic cases in the Netherlands East Indies.

I was recalled to U.K. before our outbreak had entirely subsided, but a colleague remained to follow up the cases and we intended to produce a joint report for publication on his return. Unforeseen circumstances have so far prevented this partnership from delivering the goods.—I am, etc.,

Milsted, Kent.

J. VINCENT SMITH.

The "Costoclavicular Syndrome"

SIR,—The greatest number of cases complaining of this condition are women, and the production of minor degrees of this condition may be by the shoulder strap of the brassiere. I have had three such cases in recent months, in which symptoms have been functionally relieved by doing away with the shoulder strap. The present passion for "uplift," no doubt encouraged by film stars, is not without its physical disadvantages.—I am, etc.,

Sunderland.

W. GRANT WAUGH.

What is Folic Acid?

SIR,—In the article by Prof. L. S. P. Davidson and Dr. R. H. Girdwood entitled "Folic Acid as a Therapeutic Agent" (May 3, p. 587) it is stated that free folic acid is pteroylglutamic acid, and in its conjugated form is pteroylheptaglutamic acid. Your leading article (p. 604) stated, "Folic acid does not occur

free in foods but as a conjugate with hexaglutamic acid, known as pteroylhexaglutamylglutamic acid. . . ." Wilkinson *et al.* in their article in the *Lancet*, 1946, quoting from Angier *et al.* (1946), hold that synthetic folic acid is a compound of 2-amino-4-hydroxy-6-methyl pteridine and *p*-aminobenzoylglutamic acid.

There seems, therefore, to be a difference of opinion as to what folic acid really is, both in its free and conjugated form.—I am, etc.,

Dublin.

ALFRED R. PARSONS.

REFERENCES

- Angier, R. B., *et al.* (1945). *Science*, **102**, 227.
— (1946). *Ibid.*, **103**, 667.
Wilkinson, J. F., *et al.* (1946). *Lancet*, **2**, 156.

Hypertrichosis of the Lanugo Hair in Malnutrition

SIR,—With reference to Lieut.-Col. S. K. Roy's very interesting letter on achromotrichia in tropical malnutrition (March 22, p. 392), I should like to be permitted to call attention to a little-known condition also found in some cases of malnutrition, especially in children—viz., hypertrichosis of the lanugo hair.

This condition generally affects the lanugo hair of the legs, especially the extensor surfaces. The hair becomes much thicker, longer, and at times darker. It is often seen in times of war in children who, used to an abundant, well-balanced diet, have been placed suddenly, for reasons of necessity, on an insufficient diet particularly poor in animal proteins and fat. On giving the child again a proper diet the condition rapidly ameliorates and finally disappears.

It is interesting to note that the condition is well known to women of the working classes in the South of Europe. A woman who, for reasons of work or some other reason, has put her children in charge of neighbours or relations, will often complain on her return home that the children have not been well looked after and have been starved, "as shown by the thick hair on the children's shins."

The subject of lanugo hypertrichosis in general, as well as lanugo alopecias, is an interesting one, but is beyond the scope of this letter. I might mention, however, a peculiar form of lanugo hypertrichosis of the arms and legs which occurs in certain individuals every summer (summer lanugo hypertrichosis), in others (very few) every winter (winter lanugo hypertrichosis). The subject has been dealt with briefly in my little book *Diseases of Africa* recently published (Apollon and Co., Rome).—I am, etc.,

Lisbon.

ALDO CASTELLANI.

Treatment of Hydrocele by Injection

SIR,—For many years several sclerosing substances have been used in the treatment of hydroceles with varying results. In a recent follow-up of patients suffering from hydroceles treated at this hospital in 1944, 1945, and 1946, 46 patients have been treated by injection of quinine urethane. Six patients have not been traced. Of the 40 patients traced all (with the exception of one still under treatment) have been successfully treated. Some of those treated have had hydroceles of long standing, some had previously been treated by repeated aspirations, and nearly all attended hospital seeking operative treatment. The average age of patients is 55 years, the youngest being 38, the oldest 68.

TECHNIQUE

The technique used is of the simplest kind and could be carried out in a doctor's surgery using ordinary aseptic precautions. The patient lies on the operating table, the scrotum is exposed, and the portion of the skin to be punctured is sterilized with either spirit or iodine. With a 20-ml. syringe and a size 24-in. (6.3-cm.) 18 B.W.G. bored needle the hydrocele fluid is aspirated, the utmost care being taken that the point of the needle does not injure the testicle. When all the hydrocele fluid has been withdrawn the syringe is removed leaving the needle *in situ*, the greatest care being taken that the point of the needle remains within the hydrocele sac. 1-2 ml. quinine urethane (already loaded in a 2-ml. syringe) is then introduced and the needle withdrawn. In the majority of patients little or no pain is experienced. In sensitive patients a small wheel of local anaesthesia is used where the needle is to be introduced.

The patient is again seen in a week or a fortnight's time. The local reaction varies in different patients. In some it has been necessary to aspirate when the reaction has been severe. In six patients it has been necessary to repeat the injections, in four patients the injection has been repeated twice. Patients are seen at intervals as required until the treatment has been completed. When the treat-

ment has been completed it is difficult to tell on which side the hydrocele has been, and there are no painful after-effects.

COMMENT

(1) Technique is simple and can be carried out in a doctor's surgery.

(2) Care must be taken that there is no underlying pathological condition of the testicle, such as tumour or tuberculosis or other infections. Until experience has been gained, or when in doubt, transillumination to find the most favourable spot for introducing the needle should be used.

(3) No time is lost from work, and there is no danger associated with the treatment if carefully carried out. The hydrocele fluid should be withdrawn slowly and gently. A plea is entered on behalf of the patient to avoid aspiration by the antiquated and painful method of using the trocar and cannula.

(4) A warning is given to avoid using too much sclerosing agent—for instance, 2 ml. need not be used where 1 ml. is considered sufficient. Sclerosing agents which give too much pain or severe reaction should not be used. Quinine urethane gives the best results.—I am, etc.,

Manor House Hospital,
London, N.W.11.

J. W. NICOLSON.

Hemiplegia in Young Adults

SIR,—Dr. G. Joly Dixon in his article entitled "Hemiplegia in Young Adults" (July 12, p. 53) calls attention to the frequency of strokes in this age group and to the obscurity of their aetiology.

It may be of interest to mention two cases recorded recently (Hawkins and Rewell, *Guy's Hospital Reports*, 95, Nos. 3 & 4) where massive haemorrhage of the brain took place in two healthy young adults. Both of these accidents were unheralded by any previous symptoms and were fatal. In one case hemiplegia was present shortly before death. These would have been labelled "idiopathic" had not sections been taken of the surrounding brain tissue, which appeared to the naked eye as quite normal. Microscopy showed that a haemangioma was present in each case to account for the cerebral catastrophe.

Such vascular anomalies are probably rare, but it seems possible that examination of sections of the surrounding brain in the fatal cases and arteriography in those that survive might demonstrate these lesions to be a more frequent cause of strokes in young people than is commonly believed.—I am, etc.,

Birmingham.

C. F. HAWKINS.

Acute Non-specific Diarrhoea and Dysentery

SIR,—Having enjoyed a recent article on "Myth and Mumpsimus in Medicine" it was indeed a surprise to see another example which should be added to this collection. Dr. G. R. Kershaw (May 24, p. 717) has picked up an old seaman's belief and polished it up with technical language. He states that it is "a matter of common knowledge that *chilling of the abdomen* by the rapid ingestion of cold fluid in large quantities when the body is hot does often induce diarrhoea, and it is fairly well recognized that *chilling of the anterior abdominal wall* frequently does likewise." The first assumption is one with which many Americans, who experience extremely hot summers, evidently do not agree, as one may notice from the customary serving of iced water with meals and on other occasions. As for the second assumption, it is interesting to note the tradition of the cholera belt, the fear of leaving the abdomen exposed, and the shrinking from the open louver in the Royal Navy, and the lack of evidence to show any ill effects from the cooling of the abdomen.

It is true that the latent heat of steam (or should it be of water?) is largely responsible for the cooling of the body. It should be noted, however, that this is much more effective in a dry climate than in a moist climate. In the Imperial Valley in California I have been informed that the humidity is very low, but that the temperature often keeps above 120° F. (49° C.) for weeks. One resident of this area said that the only way he and his neighbours could get to sleep before they constructed cooling devices was to wrap themselves in a sheet soaked in water. He claims that the wet sheet would evaporate in about half an hour, when the heat would be so great as to

awaken him, and by repeating the process several times he would obtain a good night's sleep. I don't recall any history of diarrhoea, and yet the cooling effect would be much greater than from similar evaporation at sea.

In the Gulf of California on one trip in a Canadian Naval ship I had occasion to try an experiment with three willing subjects in the wardroom, who volunteered with me to sleep with the louver exposed on our abdomens. Unfortunately the numbers were not sufficient to be of statistical significance, but although we experienced no diarrhoea we could not convince the other members that our action was other than folly.

The author of the article states that cooks "were from time to time victims of the outbreaks themselves," and that "cockroaches abounded at all times in the galleys and pantries." Surely one's knowledge of modern hygiene and the germ theory of disease would make one think carefully before suggesting a mechanical cause for a subsequent outbreak of diarrhoea, although as the Editor remarks in the annotation (p. 729): "any condition which lowers the resistance of the individual will favour the onset of infectious disease."

I will agree with Dr. Kershaw in his plea for experimental research. Surely this may be settled by some Naval medical officer in the Tropics who can divide his messes equally and study the effects of one group leaving the bare belly exposed to the evening miasma while the other group methodically buttons on the cholera belt.—I am, etc.,

Winnipeg.

G. L. MORGAN SMITH.

Salaries of Specialists in N.H.S.

SIR,—I was perturbed recently to see that in the advertisement for specialists under the Iraq Government a different scale of remuneration was offered for psychiatrists and, I think, for radiologists. This was considerably less than that offered for surgeons and general physicians. It has occurred to me that there is a possibility that such a hierarchy of specialists, with the surgeons at the top and the lesser breeds at the base, might be contemplated for the new State medical scheme. If this is so then it would be a wise thing to state our objections to it at once.

I would claim that all specialists have the same responsibility and that if one is properly qualified he is worthy of remuneration equal to the others. It has always provided me with sardonic amusement how someone with half the experience and qualifications will jeer at one particular specialty—for example, psychiatry—without the appreciation that the work expended would have taken the specialist to the top of any other branch of medicine. It seems to me that the M.D. and the M.R.C.P. should be considered equal to the M.S. and the F.R.C.S., and if a D.P.M. is appended to the medical degrees it should not lead to poorer remuneration. If any branch of medicine is to be regarded as inferior, then this will lead to its neglect, and those who are suitably qualified will seek a more creditable type of work.

It would be interesting to know what other psychiatrists and radiologists think of this point of view. It seems to me to be of the utmost importance that this should be clarified before the scheme becomes solidified into law.—I am, etc.,

London, W.1.

CLIFFORD ALLEN.

Medicine in the U.S.A.

SIR,—Having also spent 2½ years in an American medical school in Philadelphia I wish to associate myself wholeheartedly and in detail with everything Dr. N. K. Connolly says in his letter (July 19, p. 108). I would like to emphasize further just one point.

The teachers feel under the obligation to tell the students, in so far as they can, not only the name of the disease the patient is believed to be suffering from, but *what is going on in the patient's disordered body*. The emphasis is continuously on pathological physiology. When clinical signs have elucidated as far as possible the pathological anatomy, and the history provided guesses as to physiological changes, physiological methods are used to obtain further information. These methods are partly "clinical"—such tests as circulation time and venous pressure—and partly biochemical—serum cholesterol and insulin tolerance. One hears the latter—biochemical—methods referred to—often scathingly—as "lab. tests." The extensive