Correspondence

State Medical Service in New Zealand

SIR,—The letter from the pen of Sir Ernest Graham-Little (May 3, p. 611) invites comment. Your heading of "State Medical Service in New Zealand" is unfortunate and quite inaccurate. We have had a State salaried service in this country for many years looking after the mental hospitals, all of which, except one, are under State control. The scheme to which your correspondent refers, however, is simply a compulsory universal prepayed system whereby there is a more equitable distribution of that costly article, medical care. The only part the State plays is to collect the premiums and pay the doctor 7s. 6d. for each service rendered. It also pays the chemist for any drug ordered by the doctor providing it is one listed in the *B.P.* and *Addenda* or *B.P.C.*, and will even pay a portion of the cost of a proprietary equal to the sum payable for the official equivalent if such exists. It also pays all public hospital bills and similar sums to private hospitals, and all x-ray and laboratory facilities. Finally, it pays for immunization of school-children and provides the material free, so that now it is almost impossible to find a child who has not been immunized.

Nowadays people can get medical care for all conditions instead of having to self-diagnose and self-prescribe a patent medicine. Nowadays the doctors can and do order all sorts of laboratory procedures without wondering whether the patient can pay. And at last the profession, for the first time, have learned to think and prescribe in terms of official nomenclature and not in terms of a trade jargon covering a bewildering jumble of proprietary medicines, about the exact composition of which so many were profoundly ignorant. Yet, in spite of these and other great advances, we are told that the standards have dropped. Other people think differently. We are also told that patients cannot get attention at night and during the week-ends. I am sure that hundreds of G.P.s in this country will join with me in saying that this is grossly inaccurate and quite unjust. Certainly it is not true of the South. Finally, a dearth of specialists, if such really exists, could be readily explained by the increased amount of medical work consequent on discovering illness in people who previously could not afford medical attention. At any rate, as the general practitioner scheme has only been in operation for six years, it cannot have played any part in the alleged shortage. Furthermore, the difficulties of taking a postgraduate course in Great Britain at the present time are too real to need any stressing.

It is becoming rather tiresome to see stalwarts of organized medicine in America and Great Britain labelling this country as the Ugly Duckling in reports which are essentially superficial, misleading, and biased. If they must survey the New Zealand scene, then it is only fair that they do it thoroughly and conscientiously. It will then be found that, while the present scheme is neither wholly good nor wholly bad, yet on points it is much in advance of the old system, and that the establishment of the sociological principle that medical care shall be free and universal is here to stay.—I am, etc.,

Wanaka, New Zealand.

ALBERT D. G. BLANC.

SIR,—The following extract from a letter which has just reached me from a university professor (non-medical) in New Zealand may be found interesting and suggestive:

"If you introduce your own State Service in England, you will probably be able to profit from our mistakes. A common objection is that a young practitioner without much ability can earn a good salary so soon that he has no incentive to specialize. . . . Moreover the system does not discriminate between men of superior and inferior qualities. The general opinion is that our best medical practitioners will tend to leave New Zealand to an even greater extent than before."

I think you will agree that the above stresses a tendency which needs to be borne in mind.—I am, etc.,

Weybridge, Surrey.

A. C. LANKESTER.

Adder Bite

SIR,—In view of the fact that we have had two cases of adder bite within the last three months, we feel their case histories may be of interest. In both cases we tried the effect of "benadryl" to counter any allergic symptoms which might have arisen; in both cases this appears to have been effective. In the second case we were able to obtain anti-viperine serum, but injection was not possible until several hours after the bite, and its full value could not be assessed. In both cases there was a marked degree of shock, vomiting, and upper abdominal pain of considerable intensity. Recovery was uncomplicated, and both patients have reported no subsequent ill effects.

CASE REPORTS

Case 1.—De B., aged 60, bitten by unidentified snake—an adder from his description—at the junction of the middle and lower third left shin, at 5 p.m. on April 11, 1947. No tourniquet applied, and patient travelled thirty miles before being seen by one of us (H.H.M.) at 7.30 p.m. By this time he was considerably shocked and vomited several times, complaining of severe upper abdominal pain. A cruciate incision was made into the site of bite until blood flowed freely. No tourniquet was applied because of the time lapse (2½ hours) since the bite. The patient was given morphine 1/6 gr. (11 mg.), and the leg was immobilized between pillows. That night the patient developed swelling of lips and face; he was given benadryl 50 mg. by mouth, and the swelling subsided by morning. The following day, April 12, his leg began to swell, became brawny, and finally turned mottled blue and black up to and including the groin. This continued for one week with adenitis of left inguinal glands. On the second night his face and lips again swelled, and he was given 50 mg. of benadryl with good effect. His general condition continued satisfactory, the pain and swelling of leg after an initial three days' increase gradually subsiding, and he was discharged upon the tenth day with slight oedema of leg. On May 9 slight oedema still persisted at end of day.

Case 2.—P. T., aged 17, on June 17 found a snake coiled up under an earth bank. He picked it up, thinking it was a grass snake, whereupon it bit him on the dorsum of his left thumb over the interphalangeal joint, at 6 p.m. His young brother tied a handkerchief tourniquet around base of thumb, but apparently not very tightly. He walked one and a half miles home, where he arrived in a collapsed condition. He was immediately taken to hospital. One of us (G.E.F.) saw him at 6.45 p.m. He was then in a very shocked condition, grey in colour, pulse very rapid and weak, vomiting repeatedly and complaining of severe upper abdominal pain. A cruciate incision was made into the site of the bite and intermittent venous occlusion of upper arm was commenced. By this time there was considerable local swelling and pain. Benadryl (100 mg. in 4 oz. (114 ml.) of saline was injected P.R. and 1/6 gr. morphine given. General condition improved, and at 1 a.m. on June 17 10 ml. of anti-viperine serum was given intramuscularly into left deltoid. After a restless night sulphadiazine (2 g. stat., 1 g. 6-hourly) was commenced and continued to a total of 20 g. Benadryl 50 mg. t.d.s. was given for two days. No allergic symptoms were reported. Brawny swelling of hand, forearm, and upper arm developed, with temperature up to 100° F. (37.8° C.) on first day. Swelling became maximum on third day, involving lateral chest wall and scapular region with axillary adenitis. Region of bite became bluish, with serous discharge. Recovery began on fourth day, and patient was discharged home with slight swelling but no pain on eighth day. June 30, reported fully recovered.

—We are, etc.,

H. H. MATHIAS,
G. E. FRENCH.

Tenby, Pembrokeshire.

Choice of Drugs in Treatment of Duodenal Ulcer

SIR,—I have read with interest the article by Dr. A. H. Douthwaite (July 12, p. 43) and in particular his remarks on the use of belladonna in the treatment of peptic ulceration. With him I agree that its effectiveness is undoubted, as indicated in my article on "Dyspepsia and its Surgical Significance" in the *Journal* of June 3, 1933 (p. 954). From my own studies of its effects its main action lies in the prevention of coeliac stimuli reaching the intestinal musculature, in particular the pyloric and other junctional sphincters. By doing so it relieves all stomach excitability. The unfortunate feature in its administration, as pointed out in my article, is the low concentration in the tissues which produces the distressing subjective symptoms of dry mouth and ocular disturbances. An adequate and long course of therapy is required if permanent cure is to be reached—I have not met the patient who will submit to it.