after a meal. Some such relationship is, however, a natural consequence of the fact that the peak incidence of perforation is in the middle of the afternoon and that the majority of patients have dinner at noon. A better idea of the relationship between perforation and the last meal is obtained by considering separately those patients who perforated (a) between 6 a.m. and noon, (b) between noon and 6 p.m., and (c) between 6 p.m. and midnight. These groups are represented in Fig. 7, where it will be seen that there

FIG. 7.—Number of hours between perforation and last meal in patients perforating between (a) 6 a.m. and noon; (b) noon and 6 p.m.; (c) 6 p.m. and midnight.

is no correspondence among them as regards a peak incidence related to the preceding meal. We concluded that there was no evidence to suggest that perforation occurs at any particular phase of digestion.

Finally we investigated the possibility that perforation might be precipitated by strenuous bodily exertion. We were able to ascertain what the patients were doing at the precise moment of perforation in 504 cases, thus:

No strain (364 cases):			
Standing ..	92	Sleeping ..	42
Sitting ..	89	Washing, shaving ..	16
Walking ..	80	Kneeling ..	2
Lying awake ..	43		
Strain doubtful or mild (98 cases):			
Stooping ..	27	Eating ..	16
Crouching ..	3	Drinking ..	16
Act of sitting ..	8	Turning in bed ..	5
Act of rising ..	11	Having barium meal ..	1
Stretching up ..	11		
Straining (42 cases):			
Pushing, pulling, carrying, hammering ..	30	Playing football ..	1
At stool ..	6	Jumping ..	1
Coughing ..	1	Leaning on stomach to ease pain ..	3

From the foregoing it is clear that severe exertion plays little part in precipitating perforation. Indeed, there is a strong impression, especially on reading the detailed records, that the patients were engaged in all the normal activities of daily life when the pain of perforation struck them.

Summary

Attention is drawn to the short-term variations in frequency of perforated peptic ulcer as seen in a large series of cases treated in Glasgow hospitals in the period 1924-45.

1. Perforations are relatively uncommon in August, September, and October, and unduly common in December. The beginning of the autumn fall corresponds with the local holiday period at the end of July, and may be due to the period of rest. The December peak is not due to over-indulgence at the Christmas period.

2. Perforations are relatively uncommon on Sundays and Mondays, and become more common towards the end of the week. This may be related to rest at the week-end.

3. Perforations are uncommon during the night. The frequency of perforation increases towards the end of the morning, drops in the early afternoon, and again increases notably towards the end of the afternoon. These changes may well be related to periods of rest and fatigue.

4. No evidence was found that perforation was related to the phase of digestion or occurred during strenuous bodily exertion.

SOME CLINICAL OBSERVATIONS ON THE PRESENT OUTBREAK OF ACUTE POLIOMYELITIS

BY

W. HOWLETT KELLEHER, M.D., D.P.H.*

Medical Superintendent, Western Hospital, London

When many cases of poliomyelitis are occurring, as in the present epidemic, a varied clinical picture is presented. Confusion may arise because, with emphasis on damage to the cranial nerves, other forms of meningo-encephalitis of unknown aetiology may be simulated. The percentage of bulbar cases varies in different epidemics. In the hospital of which I am medical superintendent we have been struck by the comparatively high percentage of cases showing bulbar involvement as compared with our previous experience of the disease. Had the symptoms remained entirely brain-stem, with drowsiness a marked feature and an absence of pareses or palsies of the usual asymmetrical distribution of poliomyelitis, there would have been justification for believing that intermixed with unmistakable cases of poliomyelitis were examples of a virus meningo-encephalitis, possibly in no way related in its causal agent to poliomyelitis. From a close study of some of our cases, however, there appears to be little doubt that the cases with which we are dealing, with marked meningeal and bulbar inflammation, are in fact cases of true poliomyelitis. In only a small number was there drowsiness at onset, or even during the course of the illness. Clouding of consciousness and confusion, which appeared in a few cases, amounted to no more than the delirium seen in many acute toxic illnesses. With close observation of the progress of the cases and the ultimate finding of paresis and paralysis there was little doubt that the cause of the illness was the poliomyelitis virus.

Degrees of Attack and Age Incidence

All degrees of attack are experienced, from the case in which there is only a mild transient illness, simulating other minor illnesses such as the common cold, to the most extreme forms of paralysis involving vital centres in the medulla and causing death within a few hours. The encephalitic or bulbar types may be missed in their minor form but for the occurrence of a frank case of poliomyelitis in another member of the family.

The following cases might be quoted as exemplifying variation in degree of attack:

Two cases were admitted from one family—a woman aged 31, and her child aged 4. Had the child's been an isolated case, with no known contacts, much difficulty might have been caused in diagnosis. The child, 12 days before admission, complained of headache and pain in right leg and abdomen and had vomited. On admission temperature was normal and the child was apparently well and active, but there was distinct weakness of the right facial muscles. There was no nuchal rigidity or stiffness of spine, and Kernig's sign was absent. More complete examination, however, showed weakness of the right masseter and right sternomastoid. Cerebrospinal fluid and colloidal gold reaction—no abnormality. Diagnosis was made easier by the admission on the same day of the mother with definite poliomyelitis. She had almost complete flaccid paralysis of both lower limbs and lower part of rectus abdominis. Her symptoms had begun with abdominal pain and backache three days before admission; 24 hours before admission she had

* EDITOR'S NOTE.—Dr. Kelleher, who has had special experience of poliomyelitis during the present epidemic, has, at a time of great pressure, kindly consented to write this article at very short notice in order to afford some guidance to medical men with less experience. For any incompleteness of presentation editorial exigency must be pleaded.—ED., *B.M.J.*

headache, chilliness, and weakness of legs. On day of admission headache was severe, vomiting was fairly frequent, and there was considerable pain and rigidity of posterior cervical muscles. Cerebrospinal fluid showed typical changes: increased protein and moderate increase of cells, mainly lymphocytes. The woman was in the seventh month of pregnancy.¹

This disease should no longer be called "infantile paralysis," because frequently the adult is attacked and the attack, as in this instance, may be more severe than in the child. According to some New York statistics, whereas 40 years ago 85% of the victims were young children or infants, the proportion in more recent cases has dropped to something like 30%.

It would be wrong to deduce age incidence on the basis of the fewer than 70 cases in which poliomyelitis has been confirmed at the Western Hospital during the last two months, but it may be mentioned that 33% of the patients were under 5; 40% between 5 and 15; and 37% over 15. The disease is rare after middle age. The oldest in the present series was a man aged 41, who had a very severe form of the disease involving flaccid quadriplegia and respiratory failure. The youngest was an infant of 8 months. There have been three deaths in the series—one of them the man just mentioned; another a boy of 11, with an attack of the bulbar type and respiratory involvement; and the third an infant, with no extreme paralysis, death resulting probably not from poliomyelitis but from an intercurrent gastro-enteritis, associated at first with feeding difficulties. This child was one of twins, both bottle-fed, the other twin apparently escaping entirely.

Cranial Nerve Involvement

As an indication of the emphasis on cranial nerve involvement in the present epidemic the following instances may be given:

In a public school of mixed boys and girls there was an outbreak of illness just before the summer holidays. Many children had high temperatures and headache, some sore throat, and vomiting. One girl aged 13 complained of difficulty in swallowing and pain at back of neck. On admission she was found to have weakness of both sternomastoids and general weakness of upper extremities, but no absolute paralysis; lower limbs and trunk muscles not involved.

A boy from the same school was admitted a few days later with bulbar signs and symptoms. He had a right facial paralysis, palate affected, weak sternomastoids, and within a few days developed various other pareses and paralyzes involving both upper and lower limbs.

From a residential nursery for children under 5 a nursery trainee aged 17 was admitted on a provisional diagnosis of poliomyelitis. Two days before admission she had headache and some pain at back of eyes; on day before admission she had vomited. She was found to have a slight stiff neck, temperature 100° F. (37.8° C.), persisting for less than 24 hours; no other suggestive signs. Cerebrospinal fluid showed protein within normal limits, an increase in globulin, cells, mostly lymphocytes, 23 per c.mm.

Another young girl from the same school was admitted five days later with a history of 4 days' illness, beginning with sore throat, followed by vomiting, and accompanied by pain in the back and neck and diplopia. Slight left facial paralysis noted prior to admission. Temperature, 102° F. (38.9° C.), persisted for two days. Double facial palsy rapidly developed. Slight pharyngeal symptoms on admission developed almost to complete dysfunction, with considerable collection of mucus. Muscle weakness went on to complete flaccidity of right lower limb and left upper limb, and to a lesser degree right shoulder girdle and left lower limb.

¹ Poliomyelitis in the pregnant woman is not uncommon. One woman admitted to a London hospital this year was almost at full-term, and rapidly developed a severe form of poliomyelitis. The child was born alive after caesarean section. The woman died of respiratory and cardiac failure.

From the same institution two other trainees have been admitted with suspicious symptoms, but to date it has been impossible to confirm or exclude the possibility of very mild abortive attacks.

Considering the number of cases in which there have been cranial nerve lesions—in other words, bulbar types of poliomyelitis—it is rather remarkable that we have not had more than eight cases of involvement of respiratory function. Three cases are still being treated in respirators—an infant of 14 months, a child of 3 years, and a male adult.

Need for Systematic Examination

A detailed examination for muscle weakness should never be omitted. A group of muscles or a single muscle may be singled out by the disease. Too frequently it happens that diagnosis is made only retrospectively, the child having been allowed to walk about for a week or two after the onset of illness until a limp or drop-foot was noticed, or, being stripped for the bath, his scapula was seen to be winged owing to muscle weakness. Not only are there cases which are obvious through the manifestation of a definite paralysis, even though of minor degree, for long after the onset of the initial symptom or symptoms, but there are others with only what amounts to a weakness or paresis of a muscle group. As these weaknesses may be rapidly recovered from, they may be entirely missed unless a systematic examination is made of every case in which there is a possibility of poliomyelitis, with suggestive onset, history of contacts, possibly even signs of meningeal irritation, pointing to an acute infection involving the central nervous system.

What are the symptoms and signs which may lead to a suspicion of poliomyelitis infection, and at what stage after the infection has been contracted do they arise? Experimentally, where a virus has been transmitted to monkeys a period of from 5 to 10 days may elapse before the onset of definite paralytic disease. In the human it is difficult to be certain. Most authorities agree that the period is between 7 and 14 or possibly up to 21 days; commonly it is between 10 and 12 days.

Prodromal Illness

Initially the symptoms may in no way indicate a specific infectious disease of this nature. Those who have had the opportunity of observing large epidemics and studying family contacts have noticed an initial period of illness which may be described as prodromal or precursory, appearing within one to four days after exposure. Symptoms may be fever, general weakness, malaise, sore throat, diarrhoea, and general pains. Few practitioners in the absence of avowed cases in the neighbourhood or family would associate such a group of symptoms with the onset of poliomyelitis attack. Certain American observers¹ who have studied family and school contacts over a period after exposure and have based their diagnosis of an abortive or "sub-clinical" attack on raised temperature, state that of 22 children who had had intimate contacts with cases 14 had a febrile illness between 14 and 25 days after exposure. One of these 14 had frank poliomyelitis with transient paralysis; 2 had frank poliomyelitis without paralysis but with stiff neck, etc.; 4 had fever and mild symptoms, but without stiff neck, stiff back, or paralysis; and the other 7, although they had axillary temperatures of 98.8 to 99.8° F. (37.1–37.7° C.), seemed to have no symptoms.

Measures to be taken in such cases are the isolation of the most intimate child contacts over a period of 5 to 15 days after exposure, and the taking of temperatures daily

¹ Casey *et al.* "Sub-clinical Poliomyelitis," *Amer. J. Dis. Child.*, 1946, 72, 661.

during that period. The determination of cerebrospinal fluid protein for 3 to 5 weeks after the onset of the fever is also recommended in suspected cases.

The Pre-paralytic Phase

The prodromal illness may not be noted in every case in which poliomyelitis arises in a clinical form, and the case may proceed to the stage at which frank meningeal signs and symptoms are apparent, and these may be followed by paralysis. In this pre-paralytic phase of the disease, or in what may be called, in the cases in which paralysis does not afterwards develop, the meningitic or abortive form of poliomyelitis, the symptoms are usually considerable pyrexia, headache, vomiting, irritability, pain in the posterior cervical muscles, and, not uncommonly, exaggerated tendon and skin reflexes. Except in those rare instances in which the virus attacks the cells of the cortex, or where the patient is profoundly ill as a result of the attack of the virus on the vital centres in the medulla, the patients are conscious and alert.

In the presence of suggestive symptoms one of the earliest signs of involvement of the central nervous system is nuchal rigidity. This may be elicited in the usual fashion by placing the hand under the occiput and gently attempting flexion of the head; but a proportion of cases in which a minimal degree of nuchal rigidity is present may be missed when the examination is made in that fashion, and I have found it more useful to examine for rigidity in this group of muscles by pressure over the mid-cervical region rather than on the occiput. Occasionally the patient may suffer very little pain even when considerable stretching is caused to these muscles. When asked if it hurts, children more especially may indicate that the back of the thighs or knees is the area in which pain is produced.

In the absence of definite rigidity in the cervical region the patient may be directed to sit up in bed. The degree of weakness or rigidity that may be present—and it is important to remember that spasm of the muscle may also be responsible for dysfunction—may be estimated from the difficulty in performing this act. Where the stiffness is minimal it may be elicited by asking the patient who is sitting up in bed without apparent difficulty to attempt to kiss his bent knees.

A young man admitted to hospital with suspected poliomyelitis had had some of the group of symptoms I have described above as suggestive of the pre-paralytic or meningitic phase. He was a well-developed youth, conscious and alert, and probably a little apprehensive. There was no weakness of muscle or nuchal rigidity. Kernig's sign was absent. All reflexes were markedly exaggerated. On the following day there was slight neck stiffness, Kernig's sign was positive, and the patient could no longer kiss his bent knees in the sitting posture. Twenty-four hours later he was unable to lift his head from the bed as a result of weakness of the sternomastoid muscles, both of which appeared equally affected. The right masseter muscle was also found to be weak. Weakness of the elevators at the shoulders and of the left deltoid and right triceps was later noted. When transferred to the orthopaedic hospital a few weeks later he showed considerable improvement in the affected muscles.

Rigidity of spinal muscles may be marked, and no flexion may be possible. When the patient is raised from the supine position by a hand placed between the scapulae I have noted a sign which may be of value. As he approaches the sitting position considerable tension is thrown upon the hamstring muscles, which are obviously in spasm, and to relieve tension he will flex his knees and incline them to the side. A sign of moderate spinal rigidity may be elicited when the patient, even though able to get himself into the sitting posture, cannot support himself therein and places his hands on the bed behind him—the

so-called tripod position. Spasm of the muscles of the back of the thigh and knee may be shown in varying degrees by the reaction to attempts to elicit Kernig's sign.

Occasionally, even in this pre-paralytic phase of the disease, where there may be no obvious paralysis present, diminished tendon reflexes may be apparent; also there may be flushing of the skin, vasomotor disturbance, and pain and tenderness in certain muscle groups.

The Paralytic Phase

The paralytic phase of the disease may develop within a few hours of the first symptom. The central nervous system should be systematically examined, bearing in mind minor degrees of involvement. Examination of cranial nerves may reveal a slight facial paralysis, unilateral or bilateral. Nystagmus may be noted, and I have come across a few cases of diplopia in the present series; more rarely there may be a recognizable squint. A not uncommon finding has been weakness of the pharyngeal muscles, with difficulty in swallowing and, with the palate affected, difficulty in phonation, due to some extent to involvement of the orbicularis oris combined with the pharyngeal weakness. Careful watch should be kept for any difficulty in swallowing buccal secretions. Paralysis here may develop rapidly and give rise to much distress and even to partial or complete asphyxiation. Prompt recognition of this complication and its correction by removal of the products either directly or by postural drainage is of great importance. Careful search must be made for any abnormality of respiratory function.

Weakness of sternomastoids is shown not uncommonly in patients with other cranial nerve weaknesses. It may be elicited by asking the patient to turn his head to either side against the resistance of the observer's hand, when weakness may be noted or contraction of the muscle be made visible, especially on comparing one side with the other. The deltoid is one of the most commonly affected muscles. The patient may be quite unable to abduct his arm, or the weakness may be appreciated only by comparison with the other side.

In the determination of the degree of involvement of a group of muscles the suspected limb must first be placed in the neutral position and the patient's power to move the limb when supported in that position be observed. Next the patient's ability to move the limb against gravity may be tested, and finally his ability to move it against resistance from pressure. The movement of elbow and wrist, not forgetting the intrinsic muscles of the hand, should be tested. A limb should not be pronounced free from disability until comparative tests have been made on the opposing limb. A mild degree of weakness of the triceps, for example, may only be elicited by placing the patient's arm in the fully flexed position and then asking him to extend it against resistance. No examination of the upper limb is complete without a test for the power of elevation of the shoulders and for weakness of the pectoral and scapular muscles.

In the lower limb the muscles most commonly involved in paralysis or paresis are the quadriceps and the dorsiflexors of the ankle. Initial tests should be made for gross weakness and then more detailed examination for comparative disabilities. Where the movement is present in all directions it is most important, particularly when estimating the power of dorsiflexion of the foot, to estimate the degree of weakness on resisting movement with various degrees of pressure.

No examination, of course, is complete without attention to the muscles of the trunk, especially the abdominal and spinal. Retention of urine, although very common with

extreme paralysis, particularly when respiratory muscles are involved, may also be met with in cases without extensive paralysis, especially when there is involvement of the lower limbs. The finding of a distended bladder should lead to detailed examination for other signs of poliomyelitis.

The response of tendon reflexes should never be omitted in any complete examination, and valuable information may thus be obtained. The absence of reflexes, although the rule in severely paralysed limbs, does not by any means necessarily imply that serious damage has occurred or will follow. Cases are not rare in which, for example, with a paralysis of one or both upper limbs there may be diminished or absent knee or ankle jerks and yet no sign of muscle weakness in the lower limbs. After a considerable time during which no obvious alteration in power in the legs occurs these reflexes become fully responsive.

The usual tests for muscle function in infants and young children may be difficult, and one may easily be misled into believing that no weakness is present. An alteration in reflex response, either absence or diminution, especially in comparison with the opposite limb, should help to direct attention to a possible weakness which may be found after a patient examination. In infants particularly, by the use of stimuli as in stroking the sole of the foot with a pencil-point, muscle action may be seen, such as flexion of the thigh or dorsiflexion of the foot, where other tests had led one to believe that no function was present.

Pseudo-poliomyelitis

The present publicity given to this disease may lead to an occasional case in which some paralysis or paresis may be a manifestation of a hysterical condition.

A male aged 19, on short leave from the Forces, was taken ill with headache and vomiting, and complained of a weak and tired feeling in legs and feet. He also complained of backache and general pains in the limbs. On the day preceding admission he was unable to move his arms and had stiff neck. On admission, four days after onset of initial symptoms, he had a temperature of 99° F. (37.2° C.). He looked remarkably well. There was no trace of neck or back stiffness, he could sit up in bed without difficulty, but he had apparently a profound symmetrical paralysis of both lower limbs; there was no sensory loss; tendon reflexes were normal. There were strong suspicions that the paralysis was of hysterical origin. Improvement at first was slow, but became rapid when arrangements were made for his transfer to a Service hospital, a decision which he received with distaste.

A recent admission was a young man whose illness was ushered in by headache, sore throat, malaise, and, on day of admission, apparent inability to use right leg. He was found to have a marked streptococcal angina and a high temperature; no sign of nuchal rigidity; Kernig's sign absent. Dorsiflexion of right foot was weak. Tendon reflexes were normal. Pseudo-paralysis or paresis was suspected, confirmed by complete recovery of limb on following day.

A very high percentage of cases may prove to be abortive attacks. The American observers already quoted state that perhaps 98% of poliomyelitis is a mild, widespread, highly communicable disease, mainly of young children, leaving no residual paralysis, and the fatality rate is mentioned as not exceeding 0.7%.

Prophylaxis

Drugs, sera, and vaccines are of no proved value. There is still a good deal of misconception about the value of a serum, but once the virus has made biological union with the cell it cannot be dissociated by a so-called specific antibody. There is no proof that serum given to patients after onset of symptoms, even preliminary and non-neurological, will prevent the development of paralysis. It is possible that serum given to a contact directly after exposure may be preventive—that is, that the virus may be inactivated

before it has reached the central nervous system, though as the vast majority of contacts escape the disease without any specific therapy it is impossible to judge the value of any method of preventing the disease among those exposed.

Prophylaxis can normally only take the form of avoidance of contacts, discouragement of crowding, avoidance of severe physical strain by contacts, care in the handling of food and its protection from flies, good chlorination of public swimming-pools, and like measures. There is undeniable evidence that recent tonsillectomy and perhaps tooth extraction increase the risk. Operations on ear, nose, and throat, and possibly, in view of the fact that the virus is present in the intestinal tract, gastro-intestinal operations, are better avoided, if this is practicable, during epidemic prevalence. An intact mucous membrane is the best barrier against the entrance of the virus. In two of the patients recently admitted to this hospital, one of whom died with a severe form of bulbar poliomyelitis, and the other had an attack involving the respiratory muscles, teeth had been extracted two or three weeks before the attack, and in another case appendectomy had been performed 12 days before onset.

General Treatment

The three essentials of treatment are muscular rest, maintenance of good posture, and re-educational measures for muscles. The treatment of the case in hospital, apart from symptomatic treatment for pain, tenderness, and so forth, is mainly the province of the orthopaedic surgeon. Concentration of cases in a special hospital with facilities for physiotherapy is of prime importance. During the acute stage there should be avoidance of stimuli; action and movement of a limb may set up spasm and pain. Rest, physical and mental, is indicated. Sedatives may be necessary where there is much irritability and pain and tenderness. The patient should lie on a firm mattress under which should be laid a fracture board. I will not here enter upon a discussion as to the value or otherwise of rigid splinting as against the complete absence of splinting practised by the followers of Sister Elizabeth Kenny. Undoubtedly much harm can be done by too rigid splinting, and there is a good deal to be said in support of the management which includes good limb posture with avoidance of rigid splinting; also for passive movements at an early stage after disappearance of pain and tenderness, and carefully graded active exercises under adequate supervision in due course. During the acute stage the limb should be kept at rest in a neutral position, the lower limbs being so placed as to correspond to the normal erect posture, slightly flexed at the knees, and supported by small pads in that position. The management of the upper limbs depends on the degree of involvement. On the question of suitable support and posture so as to prevent movement of the limbs during the painful stage and tension on the affected muscles the opinion of the orthopaedic surgeon should, of course, be obtained. Massage is most usefully employed in cases in which muscles are completely paralysed or where no other treatment is available, but massage is no adequate substitute for passive and active exercises. Electrical treatment is of doubtful value at any stage. Where meningeal symptoms are prominent lumbar puncture may be of value.

Treatment of Respiratory Paralysis

The detection of minor degrees of respiratory involvement is sometimes very difficult. Involvement of the diaphragm may be partial, unilateral, or complete. Help in the detection of diaphragmatic weakness may be gained by immobilizing the lower ribs by lateral pressure, in which case the excursion of the epigastrium during downward

movement of the diaphragm may be more easily detected, or an inverse movement of the epigastrium—that is, a sucking in by that area during inspiration—may be noted. In cases of doubt I have found it of assistance to place the patient on his side, the abdominal wall being then in a neutral position, when it is easy to detect lesser degrees of excursion. Relative weaknesses of intercostal movement may be difficult to assess, but it should be recognized that there may be involvement of upper or of lower intercostals only or of all intercostal muscles.

Minor degrees of weakness may not require any accessory mechanical aid. When insufficient pulmonary ventilation occurs, however, respiratory treatment should be instituted, and the earlier the better if the involvement is of any degree. It is remarkable how promptly, if the respiratory treatment is going to do any good, the patient responds to it, falling asleep almost at once. A large number of respirators of the "Both" Nuffield cabinet type are distributed in the country, and this is the type of machine which is likely to be used for these cases. Patients with severe bulbar paralysis, having great respiratory difficulty, inconvenienced by collections of mucus in the pharynx, are extremely bad subjects for respirator treatment; their condition may even be worsened by putting them in respirators. But when paralysis is due to involvement of the phrenic or to a dorsal lesion, causing respectively diaphragmatic or intercostal weakness or paralysis, the respirator is of considerable value. Cases with extreme forms of respiratory paralysis of the spinal type may, of course, be kept alive for long periods in a respirator, but the end-result is often very disappointing. The intercostal muscles in particular seem prone to permanent weakness, and diaphragmatic recovery is often only partial.

The management of the patient in the respirator is a matter of careful detail into which the present space will not allow me to go. Much of it is a matter of highly skilled nursing. When there is complete respiratory failure owing to lesions in the cord the problem may arise, as it has in some of our cases, of keeping the patient alive and conscious during the brief periods when the cabinet is opened for change of bedclothes, skin treatment, catheterization, etc. Some form of accessory artificial respiration has to be provided, and for this the McKesson resuscitator is of value. It rhythmically inflates and deflates the lungs through a face mask closely applied, as in an anaesthetic apparatus, using pure oxygen or a mixture of CO₂ and O₂. The eventual weaning of the patient from the respirator may be a long and wearisome business, extending over months. When recovery is taking place he is allowed to breathe for gradually increasing periods without respirator aid, but sudden dyspnoea may occur and a rapid return to the machine may be necessary.

Statistical Summary

Of the 66 cases on which this article is based the age grouping is as follows:

Ages	Cases	Ages	Cases
0-5	22	26-30	5
6-10	15	31-35	4
11-15	11	36-40	0
16-20	6	41-45	1
21-25	2		

Cases showing disturbance of cranial nerves, 19; nuchal rigidity, 35; inability to sit up, 16; abortive attacks, 9.

Symptoms shown after onset and before paralysis (if any): vomiting, 37; headache, 46; sore throat, 11; catarrhal conditions, 11; rigor and shivering, 7; pain in limbs, 18; backache, 17; constipation, 6 cases.

Unusual manifestations: vertigo, 4; double vision, 3; nystagmus, 3; inequality of pupils, 2; facial paralysis, mostly unilateral, 5; diaphragm affected, 9; intercostal muscles affected, 7; masseter muscle weak or paralysed, 6; sternomastoid, one or both affected, 10; paralysis or paresis of pharynx, 9; retention of urine, 8 cases.

Of 42 cases with paralysis or paresis of limb or limbs: one upper extremity only affected, 9; one lower extremity only affected, 13; upper and lower extremities affected, 8; both upper extremities affected, 6; both lower extremities affected, 6 cases.

ARTIFICIAL CIRCULATION PRODUCED BY ROCKING

ITS USE IN DROWNING AND ANAESTHETIC EMERGENCIES

BY

F. C. EVE, M.D., F.R.C.P.

Senior Consulting Physician, Royal Infirmary, Hull

Valuable lessons can be learnt from the following case of drowning treated by manual rocking. In this method the victim, instead of being rocked on a stretcher resting on a trestle, is supported face downwards on the forearms of two operators, facing each other, who tilt the body 45 degrees each way alternately. This method I have fully described and illustrated in a booklet (Eve, 1946). There I had to admit that this new method was only on trial, because it had been tested in but one case—that of a drowned child, successfully revived but inadequately observed.

Case Report

The present case was treated by Mr. A. Akrill, a highly experienced first-aid-er and an acute observer who has been most useful to me in working out this manual method of rocking. He tells me that, in the Beverley shipyard where he is employed, a boy of 14 fell into the swollen River Hull (temperature 45° F. (7.2° C.)) on a cold November day in 1946. He soon sank, and his body was dragged out by grapnels 15 to 20 minutes later, 100 yards downstream. Mr. Akrill, with two trained helpers, arrived in two to three minutes and at once started manual rocking. The first head-down tilt drained out water freely, but during rocking the face soon became congested and blue. Schäfer's method was therefore tried—then and for brief periods later. It stopped congestion but proved less effective than rocking for ventilation and in expelling mucus. (A curious feature was that the lower jaw was propped slightly open by the tongue but was rigid till forced open.)

After about 30 minutes of mainly rocking the face changed from cyanosis to a natural pink colour, showing that an artificial circulation of oxygenated blood had been created. The boy seemed warmer too, necessitating further precautions against heat-loss. Great hopes of success were thereby raised, and after 65 to 70 minutes from the start Dr. W. H. James arrived. He tells me he found the face pale but not cyanosed, the heart inaudible, the pupils widely dilated and inactive, the eyeballs flaccid, and there was no current in the distended jugulars. Hence further efforts were abandoned. The long submersion in such cold water would make recovery highly improbable. Mr. Akrill found that he could be more useful at the head end instead of at the feet—as pictured in my booklet. For there a third man could, by his two thumbs, hold the rubber hot-water bottle to the neck, wipe mucus from the nose frequently, and help rocking with his fingers under the armpits. The boy in his wet clothes would weigh about 7 st. (45 kg.); above 10 st. (63.5 kg.) a rocking stretcher would be needed, and would always be preferable. The new Akrill thigh-grip proved efficient and labour-saving, and also kept the legs straight. These new achievements may mark a mile-