Clinic. Surgeons for the most part accept the lower figure, but Maingot in a recent article has stated that in about 10% of the cases of cancer of the stomach that come to him the change has been implanted on a previous gastric ulcer.

I personally believe that the number is very low indeed—so low that I should never use the argument of impending malignancy to bully a patient with an ulcer that had all the clinical indications of innocence into an operation she did not want to have. No surgeon in a lifetime has enough personal cases to allow him to lay down the law, but I must place on record that in twenty years of gastric surgery I have never seen a gastric ulcer become malignant. I feel that Maingot's experience is a piece of unparalleled bad luck in the practice of an otherwise very lucky man.

Cancer mimicking ulcer, on the other hand, is seen by every surgeon several times a year. It is necessary to remind ourselves that cancer of the stomach, while it is usually a very rapid disease, may be as slow as a scirrhus tumour of the breast, going on for ten years with little change; that patients with slowly progressing cancers may have a good appetite and free hydrochloric acid, and that they may lose their symptoms for a time and put on weight on a gastric diet. These slow cancers are digested in their central necrotic parts by the gastric juices, and, whether at operation or necropsy, may have very much the appearance of an indurated peptic ulcer. But in these cases, if we go into the history carefully, we find that the gastric symptoms have started suspiciously late in life, that though they may have gone for some years they have been steadily and slowly progressive, and that the remissions have not been periods of complete health but of lessened discomfort. The true ulcer usually starts in the twenties, and, in its long history before its final cure by medicine or surgery, has many remissions of months' or years' duration. I know that prolonged irritations and the presence of scar tissue may give rise to a cancer, but in a lifetime I have seen little evidence that they do so except in squamous epithelium.

On the other hand we need not be too philosophic. A large chronic gastric ulcer causes constant misery even if it does not cause cancer, and gastrectomy is one of the most satisfactory and, in practised hands, one of the safest operations in surgery. An ulcer in the bucket has never been known to become malignant.

We must suspect malignancy or a premalignant character in all gastric ulcers first appearing after the age of 40; in all those in the prepyloric inch, and in other parts of the stomach that are not on or adjacent to the lesser curvature; in those exceeding an inch in diameter; in those that do not heal after a month's strict medical treatment in bed and in which occult blood is still found in the stools after such treatment. All these should be submitted to surgery without further delay.

Conclusion

I would conclude by repeating that the early diagnosis of cancer in any internal organ depends, in the main, on clinical judgment, on the early confirmation of what can, in most cases, be no more than a suspicion. We can all remember instances of a diagnosis of malignancy made on clinical grounds, of repeated examinations by experts giving repeated denials, of confirmation coming too late for curative treatment. When cancer is in question we should accept the negative warning of a penny weighing-machine in preference to the positive reassurance of the ten-guinea Harley Street expert. And if our suspicions continue, if the weight continues to fall, we must insist on an exploration. There is more rejoicing in heaven over the one laparotomy that fails to find cancer than over the ninety-and-nine (positive ones) that find it too late.

PUBLIC HEALTH—BERLIN, 1946

BY

JAMES MELVIN, C.B.E., M.C., T.D., M.B.

Chief Control Officer, Public Health Branch, Military Government, British Troops, Berlin

[ABRIDGED]

From the entry into Berlin in July, 1945, to the end of the winter of 1945-6 the concern of the Military Government from the health and welfare point of view was almost entirely that of first-aid to a demoralized, apathetic, depressed, and needy population. A factual account of the conditions to the end of 1945 has been given by P. G. Horsburgh and H. A. Raeburn.¹ History has shown that in the train of the devastation caused by wars come epidemics which have usually claimed more victims than the wars themselves. A winter plan of epidemic prevention was therefore of first importance. The situation was complicated by vast movements of population, refugees in tens of thousands moving in all directions, but chiefly from east to west. Small groups of these, infected with typhus, dysentery, and typhoid, were a potential menace to all concerned. Barriers in the form of camps for refugees had to be established. Medical teams, hospital arrangements, disinfestation, feeding, and transport all had to be organized, and organized quickly.

Public Health Control

The Allied Control Authority takes the place of the Government in Germany, and the Public Health Control Officer of the Military Government therefore acts in the capacity of an official of the Ministry of Health, except that he has overriding powers to ensure that certain aspects of local government are carried out as ordered. Berlin is under a quadripartite government, and each of the Allies has a Public Health Branch of Military Government controlling all the medical and health services of the 20 "Bezirks" (corresponding to metropolitan boroughs in London) into which Berlin is divided. The Public Health Branch of the British Military Government consists of four officers assisted by clerks, and all the medical and auxiliary services in the four Bezirke which it includes come under its control. Direction of the doctor is within its province, and on the whole the doctors, instead of resenting this, rather look to the Branch to help them. In the British sector there is one doctor to 850 population, but in an Allied sector only one to 2,500; so that, allowing for the fact that more of the hospitals in proportion to the population are in the British sector, some general practitioners must be ordered to work in other Bezirke. There will be no compulsion upon the doctor to move his home to the other Bezirk, but only his field of employment, and the direction will be limited to "squatters" and the newly qualified. The effective measure for securing direction is through control of insurance contracts, 90% of the population being insured.

The Berlin Hospitals

The hospitals of Berlin had been reduced by May, 1945, to some 30,000 beds. By August they were up again to 38,000. The winter target was 50,000, and by the end of December 46,000 had been reached. For the British sector the target was 10,000, and the figure at the moment of writing is over 11,000. There are now practically twice as many hospital beds as in peacetime, but there are nearly three times as many sick—this with a population depleted

¹ *British Medical Journal*, 1946, 1, 423.

FIG. 1.—The Zoo Bunker, a large reinforced command post due to be blown up under the Potsdam agreement; destruction delayed that it may serve as an annexe to the Robert Koch Hospital.

to 70% of the pre-war figure. The proportion of hospital beds to the Berlin population is 15–17 per thousand, but only about one-third are in the condition which would be expected in a modern hospital. By December, 1946, temporary repairs were completed to 89% of the requirement, and permanent repairs to 70%. Some 38% of the hospitals were formerly schools, and plans are now being made to return these to their former use. The Grosse Halle at Spandau, a fine group of pavilion-like buildings, will form a hospital of about 2,000 beds divided into departments of medicine, surgery, gynaecology, and orthopaedics (Fig. 2).

The staffing of the hospitals has been a difficulty, because many of the supervisory doctors, being Nazis, have been removed. The British Military Government has made strenuous efforts over a period of 15 months to re-establish an organization of physicians and also associations of nurses and other auxiliaries, but so far this has been fruitless because quadripartite agreement could not be obtained. It would have been easy to establish these organizations in the British sector of Berlin, but it was obviously necessary to run Berlin as a city, and not as four separate cities, and eventual agreement, if and when obtained, will be well worth the time lost.

To visit the hospitals, however, and to see their work is one of the recompenses of a harassing day. Such courtesy and such keenness to listen, to demonstrate, and to co-operate are met with that considerable pleasure as well as educational value is found in these visits. The lack of sound administrators as well as mature specialists is a big problem. As denazification proceeds more and more doctors of the

able type are thrown out, and we are left with the elderly who resisted Hitler, but whose work was so valuable that the Nazis did not get rid of them. There are in the hospitals and in the specialties some young men who will be good specialists in the future, but they are few and far between at present.

Equipment and Supplies

Hospital equipment and medical supplies when the Allies took over were, of course, in a state of great shortage. With the usual German initiative hundreds of small firms came into being, some of which made pharmaceutical products in their cellars. If it had not been for the activity of these "Rucksack" firms, as the Germans called them, in

the early days our difficulties would have been much greater; the larger firms, owing to the disruption of their more intricate organization, were semi-paralysed. All this trade, however, must be brought to an end, at first by the Military Government, but at the earliest moment by the German representatives of the pharmaceutical trade and the local government Health Offices. In addition to pharmaceutical products large amounts of equipment and appliances have been brought in from the British zone, but, because of the policy of treating Berlin as one unit, they have been handed over to the central department and not allocated to the British sector only.

The Germans have been so cut off from the rest of the world that their medical men have been unaware of recent medical advances. German scientists, chemists, and doctors have not stood still, and they have several things to show for the betterment of mankind; but penicillin was generally unknown in Germany, and such discoveries as the Rh factor had not reached them. Extracts from medical literature have been presented to the Health Offices so that translations may be made for the medical profession. An article on the modern treatment of dysentery has probably contributed to the significant drop in dysentery incidence in Berlin.

Nutrition

The nutritional state of Berlin has been a cause of grave anxiety to the Allies. Cross-sections of the population have been examined at intervals of about three months. The rations in use in Berlin to-day are shown in the following table:

Ration Cards	Calories per Day	Entitled Persons
Group I	2,498	Heavy workers. Persons with active tuberculosis or severe anaemia, plus 0.5 litre milk daily
" II	1,999	Moderately heavy workers. Persons with inactive tuberculosis, convalescents from serious illness, invalids in hospital, the last receiving also 0.5 litre milk daily
" III	1,608	Normal consumers
" IVa	1,388 (without milk)	Children up to 6 years
" IVb	1,786 (with milk)	Children up to 1 year
" IVc	1,653 (with milk)	Children from 1 to 6 years
" IVd	1,487 (without milk)	Children from 6 to 9 years
" IVe	1,619 (with milk)	Children from 9 to 14 years
" V	1,559	(Discontinued since March 1, 1947)

Diabetics receive extra fat and meat, and less bread and sugar. Pregnant women from the fifth month are in Group II and receive half a litre of milk daily. Classes of the population which still give concern are young adolescents, mothers of young children, and people over 60, and efforts are being made to bring these up to a

FIG. 2.—A corner of the Grosse Halle at Spandau, which is to form a 2,000-bed hospital.

higher category of ration. The International and the Swedish Red Cross have made valuable contributions in the form of additional supplies through the medium of communal kitchens catering for children and young persons. The British Commandant in Berlin (Major-General E. P. Nares) caused the distribution of one million Arctic packs (which had been designed for an invasion of Norway) on a city-wide basis to augment the ordinary ration of school-children and adolescents in the two winters. The packs contain pemmican, biscuits, chocolate, cheese, and other concentrated foods.

Tuberculosis

Pressing problems from the public health point of view are tuberculosis, venereal diseases, and diphtheria (at the moment adult diphtheria). Unless the increasing incidence of tuberculosis can be brought to a standstill the outlook for Berlin, and indeed for Germany and even for Europe, in the next ten years is going to be serious. There are in Berlin 70,000 cases, of which between 20,000 and 30,000 are active and infectious. The British Military Government has handed over the Luftkriegsakademie at Gatow for a tuberculosis sanatorium of 1,700 beds (Fig. 3), and the U.S.

FIG. 3.—Part of the Luftkriegsakademie, formerly a Service barracks, to be converted into a 1,700-bed tuberculosis hospital.

Government has handed over the Air School building at Wannsee, which will have a 2,000-bed capacity. The tuberculosis notifications in the British Sector and Berlin in 1946 were as follows:

Notifications of Tuberculosis in 1946

	British Sector						Berlin					
	Tb. Lung		Tb. Other		Total		Tb. Lung		Tb. Other		Total	
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases	Deaths
January	364	136	5	8	369	144	1,208	753	75	63	1,283	816
Per 10,000 pop. ..	77	29	1	2	78	30	46	29	3	2	49	31
February	335	120	3	6	338	126	1,146	725	78	51	1,224	776
Per 10,000 pop. ..	78	28	1	1	78	29	48	31	3	2	51	33
March	292	128	21	7	313	135	1,401	803	180	60	1,581	863
Per 10,000 pop. ..	61	27	4	1	65	28	54	31	7	2	61	33
April	250	122	30	14	280	136	1,229	786	138	79	1,367	865
Per 10,000 pop. ..	54	26	6	3	60	29	48	31	5	3	53	34
May	248	133	27	18	275	151	1,332	731	160	90	1,492	821
Per 10,000 pop. ..	50	27	5	4	55	30	50	28	6	3	56	31
June	271	117	46	11	317	128	1,144	661	152	75	1,296	736
Per 10,000 pop. ..	56	24	9	2	65	26	45	26	6	3	51	29
July	378	110	49	16	427	126	1,412	587	228	73	1,640	660
Per 10,000 pop. ..	74	22	10	3	84	25	53	22	9	3	62	25
August	363	76	65	17	428	93	1,716	509	424	70	2,140	579
Per 10,000 pop. ..	71	15	13	3	85	18	64	19	16	3	80	22
September	275	68	59	11	334	79	1,230	427	281	46	1,511	473
Per 10,000 pop. ..	55	14	12	2	67	16	47	16	11	2	58	18
October	229	64	51	6	280	70	1,288	431	252	56	1,540	487
Per 10,000 pop. ..	44	12	10	1	54	14	48	16	9	2	57	18
November	271	69	58	8	329	77	1,252	454	230	43	1,482	497
Per 10,000 pop. ..	54	14	12	2	66	15	48	17	9	2	57	19
December	229	79	47	—	276	79	1,033	532	190	57	1,223	589
Per 10,000 pop. ..	44	15	9	—	53	15	38	20	7	2	45	22
Total	3,505	1,222	461	122	3,966	1,344	15,391	7,399	2,388	763	17,779	8,162
Per 10,000 pop. ..	60	21	8	2	68	23	49	24	8	2	57	26

FIG. 4.—Patients leaving to begin "Operation Sanatorium"—a three-months stay in hospital in Schleswig-Holstein.

A contribution from the British Public Health Branch to the tuberculosis problem in Berlin was the making of an arrangement with the Military Government authorities in Schleswig-Holstein to accept convalescent and sanatorium patients from Berlin in the summer of 1946. In July some 500 patients left Berlin in an ambulance train for a three-months stay in the sanatorium (Figs. 4 and 5). This operation was repeated monthly until October.

Other Communicable Diseases

Possibly the next most serious problem is venereal disease. In the latter months of 1946 the figures declined, but some of this decline is no doubt due to the treatment of gonorrhoea by penicillin. The steadily rising rate of syphilis gives a truer indication of the incidence. The number of fresh cases of syphilis in the British sector in January, 1946, was 165; this figure fell in the second quarter of the year, but rose again almost to the same level in the last months.

The incidence of diphtheria was highest in the British sector in January, 1946, with 291 cases and 25 deaths; in December of that year the cases still stood at 203 and the deaths numbered 16. In Berlin as a whole cases and deaths showed a general decline. There is compulsory inoculation of all children under 5. Inoculation on a voluntary basis after Schick-testing has been arranged for any of the adult

The term cases indicates fresh notifications. By the end of the year the total notified cases in Berlin amounted to approximately 75,000. These are the figures produced by the Statistical Office and are therefore final and correct.

FIG. 5.—“Operation Sanatorium”: Setting out on the ambulance train.

population particularly exposed to risk of infection.

Typhoid has steadily declined since the winter of 1945-6. It was decided that compulsory inoculation was essential, and a certificate of inoculation was made a qualification for the issue of a ration card. Typhus has disappeared from Berlin, which, considering the vast influx of refugees and displaced persons which took place throughout 1946, is a striking tribute to the effectiveness of the control measures and the efficacy of

D.D.T. powder dusting. Dysentery is still present and will remain until the sanitary conditions of Berlin have considerably improved. The death rate has been alarmingly high in relation to the total incidence, probably owing to the general state of under-nutrition and lowered resistance of the population, but partly also to the defective knowledge that German doctors generally have of the modern treatment of bacillary dysentery. A decided improvement has taken place in the death rate in the British sector and in Berlin generally since the middle of 1946. Owing to the lack of fuel and soap the incidence of scabies has shown a progressive rise. Bathing establishments and treatment centres have been set up in each Bezirk of the British sector.

A further problem awaiting solution at the moment is represented by the large number of orthopaedic cases in the city. These amount to some 70,000, of whom 30,000 are amputees (the result of war wounds or air-raid casualties). The material for orthopaedic appliances and prostheses for amputees is not obtainable at the moment. An enterprising firm is ingeniously converting German aeroplane parts into a light and serviceable type of artificial limb, but this work is held up by lack of leather and special steel to make the fittings and joints.

FIG. 6.—Repaired hospital roof converted to a solarium.

Vital Statistics

The vital statistics for each month since the entrance of the Allies into Berlin (the figures for July, 1945, are not available) are given in the following table:

	Birth Rate per 1,000		Death Rate per 1,000		Infant Mortality per 1,000 Live Births	
	British Sector	Berlin	British Sector	Berlin	British Sector	Berlin
1945						
Aug. . .	15.0	12.2	54.0	62.9	231	658
Sept. . .	15.0	13.5	54.0	55.3	231	285
Oct. . .	13.0	10.0	54.0	51.5	183	245
Nov. . .	12.0	7.9	46.0	50.7	246	190
Dec. . .	7.5	6.7	46.0	48.9	246	158
1946						
Jan. . .	4.8	5.2	38.8	41.9	118	152
Feb. . .	5.1	5.3	32.9	37.0	94	152
Mar. . .	5.4	5.2	30.7	36.1	82	136
April . .	5.0	5.0	25.8	29.1	66	97
May . .	5.3	5.1	21.2	24.1	86	111
June . .	6.2	6.4	18.6	20.2	92	103
July . .	8.4	7.9	16.1	17.9	91	100
Aug. . .	8.7	9.1	15.0	15.9	100	106
Sept. . .	11.0	10.8	15.3	16.9	102	113
Oct. . .	9.7	10.2	16.4	18.6	116	114
Nov. . .	10.6	10.1	15.7	18.6	105	114
Dec. . .	9.9	9.8	20.6	23.2	98	160

The suicides in the British sector in 1946 numbered 285, and in Berlin 653. A large increase in the number of suicides in Berlin as a whole took place in December, 1946, and January, 1947; the total number in those two months was 252. The increase is said to have been due to the accentuation of misery caused by the cold.

No description of the work of Health Control in Berlin would be complete without reference to the Commander, Brigadier W. R. N. Hinde, D.S.O., D.D.M.G. Military Government, and to his staff and our colleagues in other departments of the Military Government. The British element in Military Control in Berlin is in fact a well-knit and hard-working team. Tribute must also be paid to our Allies, with whom we spend at least one whole day twice weekly. Last, but not least, a word must be said about the sound, democratically minded German official or civilian with whom we have to work. In a difficult position he is upright, hard-working, and trustworthy. There are others, it is true; but it is our job, and in fact our mission, to bring forward and encourage the former.

A council has lately been formed for the investigation of spastic paralysis and the welfare of the often forgotten victims of this condition. The British Council for the Welfare of Spastics proposes to act as an advisory and co-ordinating centre for all activities in England and Wales directed towards the treatment, the education, and the training of these afflicted people. Prof. J. M. Mackintosh is its chairman; Lord Horder is its medical vice-president, the other vice-presidents being Sir Richard Livingstone, the educationist, and Mr. Fred Messer, M.P. On its council the Royal Colleges, the British Medical Association, and a large number of other medical bodies and bodies ancillary to medicine are represented, and six Government departments have observers. In close association is the Scottish Council for the Care of Spastics, the president of which is Sir John Fraser. The Ministry of Education has drawn the attention of local education authorities to the new council, stating that “the problem of the education and treatment of children suffering from spastic paralysis has not hitherto been attacked in this country except as part of the provision for physically handicapped children, and in one or two instances in schools or units specially provided or organized for the purpose. The Minister is of opinion that there is scope for investigation, research, and treatment which the British Council is well equipped through its several committees to undertake.”

No precise estimate of the number of children in this country suffering from spastic paralysis is available, but one figure mentioned on the basis of certain surveys is 10,000, and of this number it is supposed that rather more than a quarter are in the “mild” category and able to attend ordinary schools. In the U.S.A., where complete investigations have been carried out, it is said that, on the average, seven children with cerebral palsy are born annually in every 100,000 of the population.

The address of the British Council for the Welfare of Spastics is 34, Eccleston Square, London, S.W.1, and of the Scottish Council, County Buildings, Inverness.