"eternal triangle" which gives the "second-hand surgeon" so much trouble when operating for "benign strictures." When this dissection is properly done, enough serosa flaps always are found to cover the cystic duct and at least the lower part of the liver bed. When grafting is required, omentum is by far the best tissue. Only in one case, where the region of the cystic duct was blocked by an overlying tumour consisting of a bladder-colon fistula, I left the whole region untouched and was satisfied with the radical electrocoagulation of the opened bladder from inside without tying the cystic duct. The result, in a woman of 50 years of age handicapped by a severe thyrotoxicosis, was excellent.

May I be permitted to give a very brief summary of my experiences with electrosurgical cholecystectomy? This should positively be the operation of choice in badly infected cases in preference to the old cholecystectomy. It sterilizes the whole infected field and prevents the spreading of peritonitis beyond the operating field, still the main cause of a fatal outcome in most statistics. This can best be illustrated by the fact that not one single case of fatal issue through post-operative peritonitis was observed since 1928. The sterilizing effect of the operation can further be seen by the fact, since confirmed by all observers, that the abdominal wall can be closed also in badly infected cases.

Electrosurgical operation is the best method of dealing with liver abscesses and is, in poor-risk cases, well able to replace cholecystostomy, an operation which I have abandoned since the development of the electrosurgical cholecystectomy.—I am, etc.,

New York City.

B. O. C. PRIBRAM.

Penicillin and Blood Coagulation

SIR,—When penicillin powder was introduced for insufflation into wounds after incision I noticed that a wound which had been rendered "dry" began to bleed again after the application of penicillin powder, and occasionally I even found it necessary to secure points which, before the application of the powder, had not been bleeding. Sometimes this bleeding was very troublesome in, for instance, extensive limb injuries with much muscle damage.

It was therefore with considerable interest I read the article by Sir Alexander Fleming and Dr. E. W. Fish (Aug. 16, p. 242) and which explains a phenomenon which I am sure many surgeons operating on war wounds must have observed.—I am, etc.,

Whiston, Lancs.

W. N. CHISHOLM.

Acute Non-specific Diarrhoea and Dysentery

SIR,—I should like to associate myself with the remarks of Dr. G. L. Morgan Smith (Aug. 2, p. 189). During six years' service in India (one year in the Punjab, two in the N.W.F.P., and three in Hyderabad, Sind), and four years in Hong Kong, I never saw a case of diarrhoea which could be unequivocally ascribed to chilling of the abdomen. Personally, when the weather was very hot I invariably slept naked under a fan, and never suffered any ill effects, in spite of the warnings of some of my colleagues and of old "Koi Lais" who informed me that my demise under such conditions would be rapid.

I found that the greatest incidence of diarrhoea occurred (a) in the recent arrivals from temperate climates, and (b) during the winter and early spring, when flies were most prevalent; and I am quite convinced that food or water contamination is responsible for these outbreaks. Anyone who has had any experience of the hygienic conditions of the average cookhouse in India and other tropical countries can only be amazed that gastro-intestinal infections are not more frequent. (This particularly applies to officers' messes, which are generally the least inspected and most unhygienic of any.)

The attitude of mind of those who perpetuate this myth of "chilling of the abdomen" is further exemplified by the (until recently) prevalent ideas about so-called "sunstroke." I have never met anyone who could claim to have seen a fully authenticated case of sunstroke, though I have myself seen several cases of heat-stroke, which usually occurred in the evening when the humidity increased. While in the Tropics I never wore a *topi* except in uniform, when it was compulsory, and

frequently went about without a hat at all, in spite of the many warnings I received. I feel sure that a lot of harm has been done to the European resident in the Tropics by Kipling's lines: "But the worst of your foes is the sun overhead," because it caused unnecessary emphasis to be placed on the *topi* and spine-pad.

In my opinion, the value of a *topi* lies not in the material from which it is made but in the better ventilation it affords by reason of the corrugated head-band around the rim. The criterion I used, both as regards heat and cold, was comfort. Any type of hat, if worn for long periods in the hot sun, will cause sweating and discomfort. Its temporary removal allows evaporation of the sweat, and therefore cooling. In the same way a fan allowed to play on the naked body in bed permits one to obtain that necessary amount of sleep which is essential. Should the temperature drop during the night the bodily reactions will very soon warn one, and one can either turn off the fan or pull up a sheet.

Provided one is in reasonable health, Nature, through our own sensations, will indicate our necessities.—I am, etc.,

H. G. ROBERTSON,
Lieutenant-Colonel, R.A.M.C.

An Infants' Pharyngoscope

SIR,—An infants' pharyngoscope, which I designed several years ago, has since proved extremely efficient. It is simply a pharyngoscope blade that can be adapted to any auriscope. This makes it a valuable adjunct to a practitioner's kit. The pharyngoscope blade has a bayonet catch for use on any electric auriscope, the long length is 3 inches, the short length 2 inches, and the cut-out portion 3/8 inch. It is nickel-plated, made by Down Bros. and Mayer and Phelps.

It fits into the auriscope in the same way that the different sized aural funnels do, and the light from the auriscope bulb, projected along the hollow blade, gives an excellent illumination of the infant's larynx. With the aid of any cranked sinus forceps (I use St. Clair Thomson's cranked forceps), a size No. 4 Jacques rubber catheter can be inserted easily into the trachea. If this is connected up to a low pressure oxygen supply, which is bubbling at the rate of forty bubbles a minute, it will keep sufficient oxygen in the bronchioles to sustain the child until it can recover from its shock and take over breathing on its own initiative. This is useful in cases of asphyxia pallida, especially after operative delivery or prolonged labour where the limp infant refuses to make any attempt at respiration.—I am, etc.,

Littleover, Derby.

LINDSAY O. WATT.

Care and Treatment of Elderly and Infirm

SIR,—The report of the B.M.A. Committee on the Care and Treatment of the Elderly and Infirm (*Supplement*, June 21, p. 133) is very promising, and a step in the right direction. One important group of elderly patients which does not seem to be provided for is that of the old pulmonary tuberculous patient with a positive sputum. The curtailment of the disease generally depends a lot upon the early diagnosis and isolation of these patients. It is hoped that separate accommodation, or communities, suitably equipped for these old patients, will be provided in the proposed geriatric service, their medical treatment being under the supervision of tuberculosis officers.—I am, etc.,

Stafford.

CHARLES D. ROSS.