BRITISH MEDICAL JOURNAL

LONDON SATURDAY SEPTEMBER 27 1947

HAEMATEMESIS AND MELAENA*

WITH SPECIAL REFERENCE TO BLEEDING PEPTIC ULCER

BY

F. AVERY JONES, M.D., F.R.C.P.

Physician in Charge of Dietetic Department, Central Middlesex County Hospital;
Consultant in Gastro-enterology, British Postgraduate Medical School

[WITH PHOTOGRAVURE PLATE]

Effect of Early Feeding and Liberal Transfusion on Mortality

The more recent statistics, particularly from America and Scandinavia, show a considerable fall in mortality in the past ten years. The Scandinavian figures were never as high as the English and American were, possibly because there is easier admission to hospital and a lower proportion of chronic ulcers. The present series shows a definite fall, and if allowance is made for the older age distribution the reduction is impressive. The figures in this series are compared with those of Cullinan and Price (1932), whose paper is the best-documented one in the literature. Their patients were treated on a restricted fluid intake and only small transfusions. They took a five-year period, 1925-9, and the total number of cases of haematemesis and melaena was 165, but in many the bleeding was associated with diseases other than peptic ulcer, such as cirrhosis of the liver, carcinoma of the stomach, and other morbid conditions. These were rejected and so were cases of melaena only. The total number of cases of haematemesis from proved or probable peptic ulcer in the series was 105. To obtain a comparable series the peptic ulcer group has been taken and cases with melaena only were excluded. The scales are probably weighted against the present series, as 56% had radiological evidence of peptic ulcer, against 44% in Cullinan and Price's series. The amount of surgery performed is not sufficient to affect the main issues of comparison.

A comparison of the two series reveals an impressive fall in the mortality in the men, particularly under 60; it is less obvious, though present, in the female cases. The difference in the age distribution in the two series is well demonstrated.

Reasons for Fall in Mortality

There are important reasons for the reduction in mortality. Recurrent haemorrhages are fewer and, thanks to early feeding and liberal transfusions, the patients are in a much stronger position to withstand further bleeding. With drip transfusions fewer patients become anoxaemic, making them restless and difficult to nurse; to-day, patients no longer die from therapeutic dehydration.

The incidence of recurrent bleeding in this series may be compared with Cullinan and Price's cases. Taking patients admitted with haematemesis and showing evidence of brisk haemorrhage after admission, the figures shown in Table XIV are obtained.

TABLE XIV.—Recurrent Haemorrhages

	Cullinan and Price (1932)		Present Series (1947)	
	No.	Deaths	No.	Deaths
All admissions	109	19 (18%)	530	42 (8%)
Those which recurred	39	16 (40%)	103	26 (26%)
Those which recurred more than once	21	12 (60%)	57	17 (30%)

TABLE XIII.—Proportion of Fatal Cases in each Age Group

Age	Cullinan and Price (1932)		Present Series (1947) (Haematemesis only)	
	Males	Females	Males	Females
10-19 ..	0/1	—	0/1	—
20-29 ..	0/9	0/3	1/21	0/8
30-39 ..	2/7 (30%)	1/8	0/56	1/16
40-49 ..	5/24 (20%)	1/5	2/88 (2%)	1/30
50-59 ..	5/25 (20%)	1/12	6/79 (8%)	1/49
60-69 ..	4/8 (50%)	0/3	9/63 (14%)	6/46
70-79 ..	—	—	9/32	3/29
80+ ..	—	—	1/6	2/6
Total ..	16/74 (21%)	3/31 (10%)	28/346 (8%)	14/184 (7%)

Males: $\chi^2 = 11.875$, $n = 1$, $P = < 0.01$ (highly significant).
Females: $\chi^2 = 1.0554$, $n = 1$, $P = 0.3$ (not significant).

*The second Goulstonian Lecture given at the Royal College of Physicians of London on March 20, 1947. Lecture I appeared in last week's issue.

TABLE XV.—Statistical Analysis: Comparison between the Two Series ($n=1$)

	χ^2	P	Comment
(1) Frequency of recurrent haemorrhage	13.9757	0.01	Highly significant
(2) Frequency of multiple haemorrhages	6.1115	0.014	Significant
(3) Mortality (all admissions)	9.4624	0.01	Highly significant
(4) Mortality (single recurrence)	3.3831	0.075	On the border of significance
(5) Mortality (multiple recurrences) ..	4.9034	0.03	Significant

Table XV shows very clearly that liberal drip transfusion and early feeding actually reduce the risk of further bleeding and the mortality of recurrent haemorrhage.

The greater interest in feeding of patients has eliminated fatalities resulting from inadequate fluid intake. An examination of the post-mortem records of fatal cases of

haemorrhage from simple peptic ulcer (1931-5) showed that in 18 out of 39 cases the haemorrhage had ceased at the time of death and that many had not bled for some days previously (Avery Jones, 1939b). In seven of these cases the blood or cerebrospinal fluid urea was determined at death, and all showed uraemia, the readings being 860, 710, 540, 380, 368, 330, and 272 mg. per 100 ml. Except in one case, the kidneys did not show sufficient changes to account for the azotaemia. These patients did not die exsanguinated; nor were they old enfeebled people, for five were under 50 and one was only 28. They died from uraemia between the eighth and the sixteenth day after admission. The common factor in these cases was dehydration, and an estimate of their fluid balance from their charts showed a gross deficit. They became dehydrated because the routine treatment provided insufficient fluids, and their symptoms were probably masked by heavy sedation.

The giving of a purée diet after haemorrhage from a peptic ulcer has the advantage of supplying available water as well as calories. Although much of the improvement in mortality may be due to the elimination of dehydration the better nourishment at a time of crisis must also have helped.

Clinical Observations

The diagnosis of haematemesis and melaena would not appear to present any difficulties, but nevertheless errors can arise. Patients have been admitted with the diagnosis of congestive heart failure, angina of effort, or coronary thrombosis. These patients were mainly elderly. Disorientation from cerebral anoxia may have made it impossible to get an adequate history, and melaena has been first discovered in hospital. Pallor may have been masked by dark skin or by wartime "black-out" conditions of lighting, if admitted at night. An ill-cared-for pigmented senile individual with severe dyspnoea, oedema, and distended jugular veins may legitimately deceive the admitting physician. Cardiac pain from recent acute anaemia twice led to an erroneous initial diagnosis. Patients have been admitted direct to a surgical ward because of red blood passed per rectum, and haemorrhoids were presumed. Dark-red fluid stools have often been observed after a brisk recurrent haemorrhage in the ward, and the absence of the classical "tarry" characteristic may be misleading if observed on arrival at hospital. Schiff *et al.* (1942) have made observations on the effect of oral administration of citrated blood in man, and they found that it was necessary to give 100-150 ml. of blood to get a tarry stool. They then studied the effect of giving 1,000-2,000 ml. through a Rehfuß tube into the fasting stomach in a period of 30-60 min. The 2,000 ml. was given in three doses—700, 700, and 600 ml.—at four-hourly intervals. Three out of the six subjects had stools entirely bloody and never tarry, presumably from intestinal hypermotility. None of the subjects vomited the blood.

It was of interest to note that, whereas haematemesis occurred in all the fatal duodenal ulcer patients, melaena only was present in six of the chronic gastric ulcer patients. A common feature of these six cases was their extremely poor general condition. Three had extensive oedema, two were irrational on admission, and one had severe purulent bronchitis. It almost seemed as though they were too weak to vomit. It has been assumed in many previous papers that melaena only is less serious than haematemesis, but taking the whole peptic ulcer group there is no real difference in mortality. Thompson *et al.* (1946) noted a mortality for melaena similar to that for haematemesis in their series.

The appearance of the patient on admission soon after haemorrhage is extraordinarily misleading. Profound

circulatory collapse and deathly pallor of the lips may disappear within an hour, with great clinical improvement in those whose subsequent course is uneventful and in whom the haemoglobin had fallen only to 70%. The initial pallor is probably due to the intense vasoconstriction which may occur as a response to haemorrhage. The blood pressure during this initial phase may be moderately increased (Avery Jones, 1939a). Following a haemorrhage the pulse may be unusually full and bounding, the diastolic blood pressure cannot be registered as the sounds are still audible at zero, and capillary pulsation may be demonstrated. These changes are usually particularly marked the day after a haemorrhage and may persist for several days. The haemodynamics of the circulation after haemorrhage has been recently studied by Howarth and Sharpey-Schafer (1947).

Fever after gastro-duodenal haemorrhage is common and may persist for some days. In one fatal case the temperature rose steadily over four days. Schiff (1944) has shown that the fever is not due to the presence of blood in the intestine, and it is possibly related to the endogenous breakdown of body protein which may occur after haemorrhage (Black, 1942). Leucocytosis may accompany the fever, which is usually found after any gastro-duodenal bleeding, except in cases of cirrhosis hepatis. Leucocytosis after haemorrhage in Banti's syndrome has been observed in this series.

Persistence of pain after admission for haemorrhage is well known to be uncommon, and patients may appreciate the relief which the bleeding has brought them. The mechanism of this relief is unknown. Bonney and Pickering (1946) suggest that it may be due to blood in the crater increasing the thickness of the protective layer of slough, which may cover the pain-nerve-endings and hinder their excitation by changes in acidity in the cavity of the stomach. Gastroscopically, the slough may be obscured by blood soon after bleeding, but another explanation of the initial loss of pain may be that the large protein meal with delayed emptying which occurs after bleeding (Van Lier *et al.*, 1936) may effectively neutralize the acid in the stomach. For seven to ten days after a haemorrhage, gastroscopy usually reveals only a small haemorrhagic area in the stomach, presumably the site of the vessel; very occasionally a vessel or an aneurysmal dilatation on a vessel may be identified. At this stage pain is still very uncommon and may not be precipitated by meat, which will often bring on pain in the early days after admission for an uncomplicated gastric ulcer. Pain persisting after admission is a bad prognostic sign, and severe persistent pain has been seen only in cases which have proved fatal. Discomfort may be present if there is an associated stenosis with continued pain. Two possibilities always to be considered are subacute perforation (a plain radiograph of the diaphragm may confirm this by the presence of air under the diaphragm) and coronary thrombosis.

Coronary thrombosis was precipitated by the gastro-duodenal haemorrhage and caused death in two cases. The first of these patients had severe pain after admission, but lapsed into a coma and died on the third day. The second had persistent pain for ten days, and during this time he had three transfusions of 1,080 ml., but his haemoglobin did not rise above 43%. Following his third transfusion he developed acute left heart failure and died. Necropsy showed complete occlusion of the right coronary artery. This complication has been reported by McLaughlin *et al.* (1940), Blumgart *et al.* (1941), McKinlay (1943), and Kinney *et al.* (1945). Electrocardiographic changes may occur after acute haemorrhage without subsequent evidence of coronary artery disease. Rasmussen and Foss

(1942) report flattening, isoelectric, or negative T-waves, and lowering of the S-T interval in about half the cases.

Amaurosis is a well-recognized and tragic complication of gastro-duodenal haemorrhage. One patient aged 70 twice had a transient amaurosis lasting ten minutes following recurrences of bleeding from a duodenal ulcer. No changes were seen in the fundus. A second patient, aged 56, was admitted with brisk haemorrhage from a duodenal ulcer; he had severe recurrent haemorrhages and became blind in his only functioning eye on the eighth day. Amaurosis has been described following severe haemorrhage after parturition, abortion, and epistaxis, but it is almost unknown after trauma in primary anaemias. It is probable that recurrent haemorrhage in an already anaemic subject is the first condition necessary for its development. The vasoconstriction following bleeding may so severely reduce the blood supply to the optic nerve that irreversible damage from anoxia ensues.

The association of acute emotional stress with the haemorrhage, as described by Davies and Wilson (1939), has often been observed and cases are readily recalled to mind: the young man so perturbed by his wife's confinement that he had a severe haemorrhage just before and soon after the birth of a son; the man blown into the air by a "land-mine" who bled a few days later; the patient due to attend court over a possession order on the day of admission; the woman badly upset by a horrific film. Cases of repeated episodes of haemorrhage at times of particular stress, as noticed by Gainsborough and Slater (1946), have also been seen. Under wartime conditions of work it was impossible to make a psychological assessment of every case, but the impression left after fairly detailed inquiries in many cases is that the emotional facet was an impressive factor in only a minority of admissions. The work of Wolf and Wolff (1943), showing the engorging effect of sustained resentment, frustration, and anxiety on the gastric mucosa, gives a clue to the mechanism whereby emotional factors can promote ulceration. Such a background acting over weeks may possibly lead to peptic ulceration. The precipitation of bleeding or perforation by the final environmental upset may be through the same mechanism, but the acute anxiety often causes missed meals and extra smoking or drinking, which may be enough to cause a sudden extension of the ulceration.

Pregnancy is known to have a beneficial effect on peptic ulcer, and symptoms of active ulceration are extremely rare during that period. Sandweiss *et al.* (1943) reported only one death, from a perforated ulcer, among 70,310 women during pregnancy. Over the period of the present series there were approximately 10,000 women who attended the antenatal clinic and were confined in the hospital. Only one had a slight haematemesis, and no ulceration could be demonstrated. Among the very few referred to the dietetic clinic on account of dyspepsia none were found to have peptic ulceration. During the same time over 2,000 women were admitted with abortions and only one had haematemesis; she had an associated acute renal failure. The acute anxiety of pregnancy in unmarried women does not precipitate gastro-duodenal haemorrhage.

Certain isolated cases stand out in reviewing this series: the patient with proved duodenal ulcer whose haemoglobin did not rise satisfactorily and whose stools remained positive for occult blood and who eventually was found to have a carcinoid tumour of the caecum; a man with recurring episodes of bleeding who developed marked koilonychia after each bout of haemorrhage; a woman with Simmonds's disease who had haemorrhage from gastric carcinoma. In spite of the extreme collapse of many of the patients no case of pituitary cachexia has been observed in the follow-up

clinic. It is possible that the cases following post-partum obstetric shock are conditioned by vascular changes present in the pituitary as the result of pregnancy.

Gastroscopic Study of Patients without Radiological Proof of Peptic Ulcer

The "acute-lesion group" included 217 admissions, all of whom showed no radiological evidence of gastric ulcer and all but three had no changes in the duodenal cap. These three patients also had acute gastric ulcers, and are included for simplicity of classification. It is well recognized that a good proportion of admissions for haematemesis and melaena fail to show any evidence of peptic ulcer when a barium meal is given during convalescence. It has been tacitly assumed that these cases are due to acute ulceration or gastrostaxis, as sometimes no abnormality may be noticed at necropsy. The mortality of these "acute-lesion" patients is low; only four died—all women, aged 51, 69, 73, and 84—two showing very small acute ulcers and two not revealing any adequate source of bleeding, but autolytic post-mortem changes had occurred in the stomach.

An attempt has been made to investigate the cause of bleeding in this group, and early gastroscopy has been undertaken with great caution. This has been performed mainly between the third and the tenth day, but sometimes in the second or third week, after the cessation of bleeding. Gastroscopy is not recommended as a routine investigation soon after the haemorrhage and without previous barium-meal examination, but in this series it has been undertaken as a definite clinical investigation. No ill effects have been observed from this cautious investigation. Gastroscopy was successful in 116 out of the 217 cases. The results are shown in Table XVI.

TABLE XVI.—*Acute-lesion Group (X-ray Negative)*

	Peptic Ulcer Symptoms		Vague Dyspepsia		No Dyspepsia	
	Male	Female	Male	Female	Male	Female
No. of admissions	63	52	20	20	31	31
No. of successful gastroscopies	49	25	6	6	16	14
Gastroscopic findings:						
Gastric ulcer	34	17	3	3	5	3
Acute gastritis	—	—	3	—	2	1
Atrophic mucosa only	1	—	—	3	1	—
Hypertrophic mucosa only	5	2	—	—	2	1
Normal mucosa	9	6	—	—	6	9

The patients have been divided into three classes: (1) those with symptoms suggesting a peptic ulcer, particularly a remitting dyspepsia with epigastric discomfort or pain related to food and relieved by alkalis; (2) patients with a vague indefinite mild dyspepsia; and (3) those with no digestive symptoms at all. It will be seen that the gastroscopy contributed information of special interest in 86 out of the 116 cases, and a gastric ulcer was found in 65 cases.

Gastroscopically, every gradation of ulcer from a small superficial red-rimmed ulcer to a deep penetrating crater can be found in this group of x-ray-negative cases. For convenience they can be divided arbitrarily into four groups.

Group I.—These are superficial flat ulcers, mainly less than 1 cm. in diameter, often with a red rim of hyperaemia but little or no oedema. Usually there is no generalized change in the gastric mucosa. These ulcers would almost certainly not be demonstrated radiologically even if x-rayed the same day.

Group II.—These are larger ulcers, probably 1–2 cm. in diameter, with the slough apparently a little depressed and with moderate or gross redness and swelling around the ulcer. These would probably be demonstrated radiologically if radiographed the same day, but are not seen one to three weeks later on account of rapid healing.

Group III.—These are still larger ulcers, probably more than 2 cm. in diameter, with much swelling around them and marked depression of the slough. They correspond with "chronic" gastric ulcers seen normally, but they heal rapidly after the haematemesis, and may not be observed radiologically one to three weeks after the gastroscopy.

Group IV.—These are scars of ulcers. These patients have either had gastroscopy later than usual or the bleeding has come from a duodenal ulcer.

The majority of these acute ulcers are found in Group I; five patients had multiple ulcers—two having three ulcers and three having two. In three others there was an unassociated scar of an ulcer. These small acute ulcers were particularly common on the posterior wall. Healing might occur even within five to twenty days, and six out of 13 left a visible scar. The important findings are summarized in Table XVII.

TABLE XVII.—"Acute" Gastric Ulcers

	Group I	Group II	Group III	Group IV
No.	38	14	5	8
Male	21	11	4	6
M/F ratio	1.2:1	4:1	4:1	3:1
Multiple ulcers	5	2	—	—
Position:				
Anterior wall	4	1	—	—
Posterior wall	27	8	1	2
Lesser curve	11	9	4	5
Greater curve	—	—	—	—
Pyloric antrum	1	—	—	1
Atrophy of mucosa	16	1	—	1
	(13 female)			
Final scar	6	5	3	8
No scar visible	7	2	—	—

Evidence of atrophy of the mucosa was present in 18 of the 65 cases, 17 had a pre-histamine achlorhydria, and five were histamine-fast. These acute ulcers occurring in atrophic mucosae with achlorhydria have been previously reported by Rodgers and Avery Jones (1938) and confirmed by Hurst and Lintott (1939). The remaining cases all had acid curves rising not higher than 60 units of free acid.

Case Reports

The following cases are illustrative of these acute ulcers causing haemorrhage.

Group I

Male aged 58. No dyspepsia. Admitted after sudden haematemesis and melaena. Haemoglobin fell to 38%, and he received 540 ml. of blood. Gastroscopy on the eighth day showed a small flat ulcer on the posterior wall; this had completely healed by the 13th day (Plate, Fig. 1). A barium meal on the 24th day showed normal stomach and duodenum. A fractional test meal revealed free acid rising to 22 units.

Female aged 61. For many years slight indigestion, with tight feeling in chest after meals; never any pain. No recent symptoms. Sudden haematemesis. Recurrent bleeding on first and third days. Blood transfusions totalled 2,160 ml. Gastroscopy on the 11th day showed three flat ulcers—one on the lesser curve, one on the posterior wall, and one in the region of the greater curve, but situated on a mid-gastric constriction ring (Plate, Fig. 2). On the 25th day the first two ulcers had healed and the other two were much smaller; these had healed by the 39th day, but the ring constriction persisted, and was seen radiologically as an incisura on the 31st day. The acid curve rose to 32 units.

Female aged 48. History of 30 years' remittent dyspepsia with epigastric pain one hour after meals. She had had four haematemeses, the last one 16 years ago. An attack of dyspepsia began two weeks before admission and haematemesis three days before. Gastroscopy on the second day after admission showed an organic hour-glass constriction with an acute ulcer just above it, and there was gross atrophy of the mucosa. The lower sac was seen (using the Hermon Taylor instrument) and no further ulcer found. On the 16th day the

ulcer had healed, leaving a small scar. A radiograph on the 11th day showed a mid-gastric stenosis only. Haemoglobin fell to 50%.

Group II

Male aged 39. Three days' left epigastric pain. Sudden haematemesis. Haemoglobin fell to 40%. Gastroscopy on the eighth day showed an ulcer just above the angulus (Plate, Fig. 3), and this had healed by the 22nd day (Plate, Fig. 4). A radiograph on the 19th day showed normal stomach and duodenum.

Female aged 72. For 20 years she had had occasional attacks of indigestion lasting two weeks. Epigastric pain occurred after meals for the past six weeks. She was admitted with haematemesis and had a recurrent haemorrhage next day. Haemoglobin fell to 30%, so 1,620 ml. of blood was given. Gastroscopy on the seventh day showed a medium-sized ulcer high up on the lesser curve (Plate, Fig. 5), which had almost healed by the 21st day (Plate, Fig. 6). A test meal revealed a pre-histamine achlorhydria, and a barium meal in the fourth week showed normal stomach and duodenum.

Group III

Male aged 68. Eight years' history of recurring bouts of epigastric pain after meals, attacks lasting one to two weeks. Haematemesis seven years and three years previously. On the second occasion he was admitted to hospital, and gastroscopy on the third day showed a small ulcer on the posterior wall (Plate, Fig. 7). This was almost healed two weeks later and did not appear radiologically. He had had pain again, this time severe for one month, and a radiograph had shown a gastric ulcer high up on the posterior wall. He was admitted with haematemesis, and had four severe recurrent bleedings in the next four days and at one time was profoundly anoxaemic. He had 4,860 ml. of blood. Gastroscopy on the 19th day after admission showed a large active ulcer high up on the posterior wall (Plate, Fig. 8), but the radiologist, though aware of its presence, was quite unable to demonstrate it the following day. The patient had a second severe episode of repeated bleeding in the sixth week and received a further 4,320 ml. of blood, but the ulcer eventually healed. He was a frail, wizened man, and operation would have been impracticable because of his poor general condition.

It is possible that the acute ulcer may sometimes be the result of and not the cause of the bleeding, which has occurred as a diapedesis from the stomach mucosa or from intramucosal haemorrhages. Ulcers have twice been seen to develop from intramucosal haemorrhages, and this occurred in the following case:

Female aged 49. Two years previously she had had a haematemesis without previous dyspepsia and the barium meal was negative. No pain since. Admitted with sudden melaena and haematemesis. She had brisk recurrent bleeding on the first and fourth days. On the third day gastroscopy showed multiple intramucosal haemorrhages, and on the 10th day these had all disappeared, but there was a flat acute ulcer (0.5 cm.) on the posterior wall lower down where there had been a particularly large haemorrhage (Plate, Figs. 9 and 10). The mucosa on the anterior wall had the appearance of being atrophic. Barium meal on the 40th day showed a normal stomach and duodenum. The capillary fragility platelet count, bleeding time, and clotting time were all normal on the fourth day. A fractional test meal showed acid rising to 36 units. The haemoglobin fell to 38%, and she was given 1,080 ml. of blood.

Gastritis

There are many difficulties in the diagnosis of gastroscopic gastritis (Avery Jones, 1944), and a bias against the diagnosis has become greater with increasing experience of gastroscopy. It is true to say that minor changes in the gastric mucosa have been discounted. An appearance which with the Wolf-Schindler gastroscope would suggest mucopurulent exudate may often be demonstrated as fine froth with the better vision and greater mobility of the

Hermion Taylor instrument. Nevertheless, five patients did show an apparent generalized acute inflammatory reaction with hyperaemia, oedema, and erosions. The differential diagnosis from a submucosal infiltrating carcinoma is important, but was excluded in these cases by the follow-up. One case is described below.

Male aged 53. In hospital for treatment for infected hydrocele and urinary infection. Sudden haematemesis occurred. Lowest haemoglobin 55%. Gastroscopy on the eighth day showed an intense erosive gastritis, particularly in the upper part of the stomach. By the 15th day the erosions had nearly healed and there was much less mucosal swelling; by the 29th day the mucosa appeared normal. A barium meal showed no abnormality. At no time did he have any abdominal discomfort or pain.

In two cases of acute ulcer the gross irregularity of mammillation with patchy atrophy suggested a diffuse chronic gastritis. A number showed some degree of atrophy, but the appearance did not suggest a previous inflammatory reaction.

These gastroscopic studies on the cases which do not reveal any radiological evidence of ulceration have clearly shown that acute rapidly healing ulcers are a common cause of haemorrhage and that acute gastritis is rare. As many of the patients have had precisely similar bouts of epigastric pain lasting from a few days to three weeks, but without haemorrhage, it is reasonable to assume that these episodes have been due to a previous similar acute ulcer and that a negative radiograph in such patients must not be declared as evidence against peptic ulcer.

Comparison of Acute and Chronic Gastric Ulcers

In view of the much better prognosis of acute ulcers it is necessary to be able to distinguish them so far as is possible by the bedside, especially when the question of surgical intervention arises. The acute rapidly healing ulcers in Group I, together with the two fatal cases of acute ulcer, have been contrasted with forty consecutive cases in the chronic gastric ulcer group.

The ratio of males to females was 3.5:1 for chronic gastric ulcer and 1:1 for acute ulcer. There was no appreciable difference in age distribution or in the length of history of dyspepsia. The acute ulcer cases did not give histories of protracted spells of pain, but gave short intermittent attacks lasting a few days to three weeks. The recent pain, if present, is never severe with acute ulcers and has seldom lasted more than two weeks. Absence of pain is a common history (Table XVIII). Recurrence of bleed-

TABLE XVIII.—*Period of Recent Pain before Haemorrhage*

	Chronic Gastric Ulcer (40)	Acute Gastric Ulcer (40)
3-12 months	2	—
4-13 weeks	10	1
2-4 "	11	4
1-2 "	9	12
1 week	4	7
None	4	14
Not recorded	—	2
Character of recent pain:		
Severe pain	14	0

ing after admission is just as common with acute as with chronic ulcer. Recurrence after the third day is in favour of a chronic ulcer (Table XIX). Bleeding may be extremely severe from the acute ulcers, but the average severity is greater with chronic ulcer, judging by the fall in the haemoglobin and the amount of blood transfused.

The same aetiological factors seemed to be at play in these acute ulcer patients as in those with chronic lesions.

Many had had excessive fatigue, anxiety, sustained resentment, or frustration. In a few an acute emotional upset had precipitated the bleeding. The general impression left was that these acute ulcers were essentially due to the same factors as chronic ulcer, but as they occurred in people

TABLE XIX.—*Recurrence of Bleeding after Admission to Hospital*

	Chronic Gastric Ulcer (40)	Acute Gastric Ulcer (40)
No. with recurrence	13	11
Total No. of brisk recurrent bleedings	31	39
Days of first recurrence:		
1st	4	5
2nd	2	5
3rd	1	1
4th	2	—
5th	1	—
Later	3	—

with a low or absent acid secretion the ulcer developed in a medium favourable for healing and therefore resolved quickly.

Management

Early reassurance of the patient is essential and lessens the anxiety which may reflect itself adversely on the stomach. Sedation with morphine or sodium phenobarbitone should be used only to reinforce mental calmness, and heavy sedation must be avoided. The patient should be made comfortable, with sufficient pillows to suit his personal wishes. A detailed history should be taken as soon as possible. A severely ill patient must not be considered an excuse for delaying this essential step. It is necessary to interview relatives at once, as essential points can often be elicited. In hospital practice there is a tendency to focus attention entirely on the patient; but relatives also need reassurance, and the opportunity to enrol them for the blood bank must not be missed.

Having obtained a full history, it is essential to decide whether operation should be considered if the patient bleeds again. The nursing staff should keep an hourly pulse chart, and blood-pressure readings should be recorded when the patient is seen by the medical staff. In addition 5 ml. of blood should be obtained for initial haemoglobin, blood urea, and blood-grouping tests, including Rh-screening, and plasma for direct testing of blood if transfusion becomes necessary. Caution must be exercised in interpreting the initial haemoglobin, as haemodilution may not have been complete.

The patient should be allowed a semi-solid purée diet with two-hourly feeds or 7-oz. (200-ml.) milk feeds if preferred. The great majority enjoy a soft diet from the onset. Thirst may be entirely prevented by allowing as much one-third normal saline, flavoured with fruit juice, as desired, and this should be readily available at the bedside for the first three days. Antacids are not essential, but nevertheless it is reasonable to diminish proteolytic activity in the stomach, and aluminium hydroxide may depress acidity and also peptic activity—1/2 oz. (14 g.) (6% emulsion) two-hourly by day and a double dose at 10 p.m. The bowels are usually constive for several days, and no steps other than reassurance are necessary for at least four days, when a simple enema may be given.

For severely ill patients there are certain procedures which may be of value. In cases with a long history of duodenal ulcer a continuous milk drip with 6 pints (3.4 litres) daily may be of value initially, or, alternatively, 2 pints (1.1 litres) may be given by drip during the night. In very ill patients in whom the vomiting reflex is decreased and the stomach is very distended with blood the emptying of it with an oesophageal tube and a Senoran's evacuator may much improve the general condition.

Cellular respiratory processes may be greatly depressed by anoxia and irreversible changes sometimes develop. Efficient oxygen therapy and the administration of vitamins B and C may be of some value and are recommended—for example, ascorbic acid, 500 mg.; aneurin, 25 mg.; nicotinic acid amide, 200 mg.; riboflavin, 10 mg. daily. These may be given intramuscularly or in a saline transfusion.

Apart from its use in gravely ill patients, ascorbic acid is a valuable routine therapy, as the patients may have had a deficient intake, which impairs the healing reaction of their tissues. The dosage recommended is 200 mg. for five days and then 50 mg. daily. Other medication will usually include "fersolate," 3 gr. (0.2 g.) thrice daily, or iron and ammonium citrate, 30 gr. (2 g.) thrice daily, when the stools have become light in colour again. Sedatives—for example, phenobarbitone, 1/2–1 gr. (32–65 mg.) two or three times a day—are usually advisable for patients with probable duodenal ulceration.

Conclusions

The general observations which may be drawn from this study are fourfold. First, the potentialities of municipal hospitals as a source of large-scale clinical investigations are demonstrated. Secondly, the number of cases reflects the magnitude of the morbidity from peptic ulcer in the general population. Survey studies in industry and transport by my colleague Dr. R. Doll show that 5–7% of these working-men have, or have had, peptic ulcer. A morbidity of this degree from a recurring disease certainly demands further research into its causation and improved facilities for its treatment. Thirdly, there are considerable advantages in concentrating the gastro-enterological emergencies into special wards. The senior nursing staff become particularly expert in treating them, and an especially interested physician has the opportunity of rapidly acquiring a wide experience of the particular medical problems. Cases of acute perforation are also admitted to these wards and post-operative care is undertaken there after elective surgery. A complete perspective of peptic ulcer is therefore obtained.

Fourthly, the conclusions concerning haematemesis and melaena are: (1) there are fallacies in many of the past statistics on bleeding and peptic ulcer; (2) there is still an appreciable mortality (8%) among cases admitted to a municipal hospital; (3) associated complications, giant ulcers, and old age are important factors contributing to death; (4) it is important to consider prognosis in relation to type of lesion and age of patient (the prognosis in different age groups in this series of 687 patients may be ascertained from the tables); (5) the incidence of brisk recurrent bleeding after admission, and also the mortality, have fallen since the introduction of early feeding and drip blood transfusion; (6) there is scope for emergency partial gastrectomy in the management of haematemesis and melaena; (7) acute gastric ulcer is an important cause of bleeding and can be demonstrated in 60% of the patients who have no x-ray evidence of peptic ulcer and in whom gastroscopy has been carried out; and (8) these acute gastric ulcers may cause severe and recurrent bleeding, but they carry a very low mortality.

This work has not been done alone, and I wish to express my thanks to my assistant, Dr. H. Pollak, for the able help he has given me; to my medical colleagues for allowing me to treat all admissions for haematemesis and melaena; to my surgical colleagues for their excellent co-operation; to my pathology colleague, Dr. Pagel; to Dr. Discombe for statistical advice; to Dr. R. Doll, Dr. S. Ransom, and Dr. Antia for help in the preparation of these lectures; to Dr. G. Graham for valuable criticism; and to Prof. L. J. Witts for the initial stimulus which he gave me in this problem. I should also point out that these two articles have been presented in an abridged form. Full details about the surgical cases will be published elsewhere.

REFERENCES

- Aitken, R. S. (1934). *Lancet*, 1, 839.
 Babey, A. M., and Hurst, A. F. (1936). *Guy's Hosp. Rep.*, 86, 129.
 Baker, H. L., and Halley, H. (1946). *Ann. Surg.*, 123, 1067.
 Black, D. A. K. (1942). *Quart. J. Med.*, 35, 77.
 Blumgart, H. L., Schlesinger, M. J., and Zoll, P. M. (1941). *Arch. intern. Med.*, 68, 181.
 Bolton, C. (1913). *Ulcer of the Stomach*. Arnold, London.
 Bonney, G. L. W., and Pickering, G. W. (1946). *Clin. Sci.*, 6, 63.
 Bulmer, E. (1927). *Lancet*, 2, 169.
 — (1932). *Ibid.*, 2, 720.
 Burger, G., and Hartfall, S. J. (1934). *Guy's Hosp. Rep.*, 84, 197.
 Chiesman, W. E. (1932). *Lancet*, 2, 722.
 Clemmesen, J., and Lund, K. (1939). *Nord. med. Tidskr.*, 2, 1729.
 Crohn, B. B. (1927). *Affections of the Stomach*. Saunders, Philadelphia and London.
 Cullinan, E. R., and Price, R. K. (1932). *St. Bart's Hosp. Rep.*, 65, 185.
 Davies, D. T., and Wilson, A. T. M. (1939). *Lancet*, 2, 723.
 Dudley, H. D. (1934). *Surg. Clin. N. Amer.*, 14, 1331.
 Dicks, R., Knudsen, A. F., and Franco, S. C. (1942). *Arch. intern. Med.*, 70, 121.
 Finsterer, H. (1939). *Surg. Gynec. Obstet.*, 69, 291.
 Frank, W. (1946). *Gastroenterology*, 7, 231.
 Gainsborough, H., and Slater, E. (1946). *British Medical Journal*, 2, 253.
 Goldman, L. (1936). *J. Amer. med. Ass.*, 107, 1537.
 Gordon-Taylor, G. (1937). *Brit. J. Surg.*, 25, 403.
 — (1943). *British Medical Journal*, 1, 504.
 Griggs, D. E., and Baker, M. Q. (1941). *Amer. J. digest. Dis.*, 8, 344.
 Hanno, H. A., and Mench, M. (1944). *Ann. Surg.*, 120, 199.
 Hawkins, H. P. (1907). *Med.-chir. Trans.*, 90, 268.
 Hellier, F. F. (1934). *Lancet*, 2, 1271.
 Howarth, S., and Sharpey-Schafer, E. P. (1947). *Ibid.*, 1, 18.
 Hurst, A. (1924). *Proc. roy. Soc. Med.*, 17, Sect. Surg., Med., and Therap., 13.
 — and Lintott, G. A. M. (1939). *Guy's Hosp. Rep.*, 89, 173, 177.
 — and Ryle, J. A. (1937). *Lancet*, 1, 1.
 Illingworth, C. F. W., Scott, L. D. W., and Jamieson, R. A. (1944). *British Medical Journal*, 2, 617.
 Jennings, D. (1940). *Lancet*, 1, 395.
 Jones, F. Avery (1939a). *British Medical Journal*, 1, 915.
 — (1939b). *Ibid.*, 2, 332.
 — (1944). *Proc. roy. Soc. Med.*, 38, 82.
 Kinney, T. D., and Mallory, G. K. (1945). *New Engl. J. Med.*, 232, 215.
 Kirsner, J. B., and Palmer, W. L. (1939). *Int. Clin.*, 4, 105.
 Klingenstein, P. (1938). *J. Mt. Sinai Hosp., N.Y.*, 4, 972.
 McKinlay, C. A. (1943). *Journal Lancet*, 63, 31.
 McLaughlin, C. W., Baker, C. P., and Sharpe, J. C. (1940). *Nebraska med. J.*, 25, 266.
 Marriott, H. L., and Kekwick, A. (1935). *Lancet*, 1, 977.
 Meulengracht, E. (1935). *Ibid.*, 2, 1220.
 — (1936). *Wien. klin. Wschr.*, 49, 1481.
 — (1939). *British Medical Journal*, 2, 321.
 Morris, J. N., and Titmuss, R. M. (1944). *Lancet*, 2, 841.
 Murphy, B. (1942). *Ibid.*, 1, 704.
 Paterson, H. J. (1924). *Proc. roy. Soc. Med.*, 17, Sect. Surg., Med., and Therap., 1.
 Rasberry, E. A., and Miller, T. B. (1943). *Gastroenterology*, 1, 911.
 Rasmussen, H., and Foss, M. (1942). *Acta med. scand.*, 111, 420.
 Rodgers, H. W., and Jones, F. Avery (1938). *St. Bart's Hosp. Rep.*, 71, 141.
 Ryle, J. A. (1937). See Hurst and Ryle (1937).
 Sandweiss, D. J., Podolsky, H. M., Saltzstein, H. C., and Farberman, A. A. (1943). *Amer. J. Obstet. Gynec.*, 45, 131.
 Schiff, L. (1944). *Sth. med. J., Nashville*, 37, 335.
 — Shapiro, N., and Stevens, R. J. (1944). *Amer. J. med. Sci.*, 207, 465.
 — Stevens, R. J., Shapiro, N., and Goodman, S. (1942). *Ibid.*, 203, 409.
 Scott, L. D. W. (1940). *Edinb. med. J.*, 47, 49.
 Segal, H. L., Scott, W. J. M., and Watson, J. S. (1945). *J. Amer. med. Ass.*, 129, 116.
 Sharpey-Schafer, E. P. (1945). *Lancet*, 2, 296.
 Smith, D. (1945). *Glass. med. J.*, 144, 129.
 Thompson, H. L., Oyster, J. M., Heide, J. B., and Morgan, F. M. (1946). *Gastroenterology*, 7, 320.
 Tidy, H. L. (1944). *British Medical Journal*, 1, 677.
 Van Liere, E. J., Sleeth, C. K., and Northrup, D. (1936). *Amer. J. Physiol.*, 117, 226.
 Wiener, A. S., and Gordon, E. B. S. (1947). *Amer. J. clin. Path.*, 17, 67.
 Witts, L. J. (1937). *British Medical Journal*, 1, 847.
 Wolf, S., and Wolff, H. G. (1943). *Human Gastric Function*. Oxford Univ. Press, London and New York.

To further its campaign against poverty, ignorance, and disease Unesco has published a book, *Fundamental Education: Common Ground for All Peoples*, written by authorities of various nations who worked in the Preparatory Commission of Unesco during 1946. The book gives an account of problems in different parts of the world and describes the notable measures taken against ignorance and illiteracy in such countries as the U.S.S.R. and Mexico.

FIG. 1.—Small acute ulcer, healed five days later.

FIG. 5.—Subacute ulcer.

FIG. 9.—Multiple intramucosal haemorrhages.

FIG. 2.—Acute ulcer on a mid-gastric constriction-ring.

FIG. 6.—Same ulcer as Fig. 5, almost healed after two weeks.

FIG. 10.—Haemorrhages disappeared in one week, leaving small acute ulcer.

FIG. 3.—Subacute ulcer.

FIG. 7.—Subacute ulcer, posterior wall.

FIG. 4.—Same ulcer as Fig. 3, healed after two weeks.

FIG. 8.—Large ulcer, posterior wall.

CASE OF PINEAL CYST:
SIMON SEVITT AND J. SCHORSTEIN

5

FIG. 1.—Thin glial layer externally (A), middle epithelial layer of pineal parenchyma (B), and internal pigment-containing gliotic layer (C).

FIG. 2.—High-power view of B, showing epithelial cells and one corpus arenaceum.