Without submitting any ready solution of Africa's problems I suggest that it should be borne in mind that the population problem is one of balance, and that the economy of grossly underdeveloped countries can be improved more readily and more to the general advantage of the world by an increase in the supportive capacity of the land than by a restriction of population, though ultimately that must come. Colonial development is not "whistling in the wind" but the essential background of human welfare, which should be at least as much the concern of your readers as the policy of despair you advocate. The ignorant and poverty-stricken peasant is admittedly an inadequate foundation for an enlightened state of society, or for the production of goods to be exchanged with industrial countries. The remedy may lie in the cure of his ignorance and poverty rather than in the restriction of his numbers.—I am, etc.,

London, W.C.1.

G. MACDONALD.

REFERENCES

- Baker, S. J. K. (1937). *Africa*, 10, 37.
 — and White, A. T. (1947). *Geographical Journal*, 108, 198.
 Dixey, F. (1926). *Geographical Review*, 18, 274.
 Gillman, C. (1936). *Ibid.*, 26, 353.

Poliomyelitis

SIR,—Three out of a total number of 33 cases of acute anterior poliomyelitis under my care have developed symptoms which were regarded as due to paralytic ileus. The three cases were severe, with widespread paralyses, and the symptoms of bowel paralysis appeared three days after the first onset of paralyses in the voluntary muscles.

In one of the cases there was no abdominal pain, but in the other two complaint was made of abdominal aching. In all the cases the abdomen began to swell progressively; the swelling was tympanitic and there was absolute constipation. No flatus was passed. In one of the cases, a woman of 50, the swelling rapidly became enormous, but she complained of no feeling of discomfort. The main danger of the abdominal distension is embarrassment to the heart and respirations in a patient with some degree of diaphragmatic paralysis.

Treatment consisted of 1 ml. of "prostigmin" followed four hours later by a further 1 ml. and a turpentine enema. This resulted in slight colicky pain, followed by the passage of flatus and faeces, with disappearance of the abdominal distension. In children the dose of prostigmin used was 0.5 ml. After the subsidence of the abdominal swelling constipation again followed in one of the cases, but prostigmin 0.5 ml. daily relieved this.

The treatment I have used in all cases with paralysis has been: vitamin B₁ tablets, 1 t.d.s., together with strychnine, 1/32 gr. (2 mg.) b.i.d. in the case of an adult, and 1/60 gr. (1 mg.) in the case of a child.—I am, etc.,

Newport, Isle of Wight.

JOHN MILLS.

SIR,—Though there is no specific treatment as yet for this disease, for some six or seven years I have treated the comparatively few patients I have seen in the very early stages of the attack by intramuscular injections of their own cerebrospinal fluid, 5 ml. given daily or on alternate days according to the severity of the illness, three or four injections being the usual amount. As many patients recover completely without any treatment, and as the general tendency is towards at least partial recovery, it is impossible to assess the value of this or any other form of treatment. In a number of cases it appeared to be of some value towards arresting the progress of the attack.

Two patients were treated by intramuscular injections of blood taken from people who had had the disease; some benefit appeared to result. Is it possible that convalescent serum (such as is given to abort or attenuate an attack of measles) might be of use? Obviously this form of prophylactic treatment could only have a very limited application.—I am, etc.,

Purbrook, Hants.

GEOFFREY EDEN.

SIR,—In Prof. H. J. Seddon's recent paper on the early treatment of poliomyelitis (Aug. 30, p. 319) it is very depressing to read that no really early treatment is possible. During the past few years letters have appeared in the *British Medical Journal* advocating early treatment for herpes zoster—another virus disease which attacks the central nervous system—publishing records of cases showing successful results. In my own

practice treatment of herpes zoster by injections of vitamin B₁, 25 mg., and liver extract and "contramine," 0.125 g., repeated if necessary, appears to have yielded satisfactory results. After a personal attack I can vouch for the fact that these injections are only uncomfortable.

In the present epidemic many cases will probably be diagnosed in the pre-paralytic stage, and it would appear possible that this line of treatment may be worthy of a trial, particularly when there is no alternative. Whether the present widespread sub-optimum intake of vitamin B₁ is a factor in the spread of poliomyelitis is another question which might be asked of those qualified to give an authoritative answer—I am, etc.,

St. Faiths, Norfolk.

J. N. GALE.

SIR,—It was with great interest that I read the note by Dr. J. N. McIntosh (Gloucester) (Sept. 6, p. 398) on the treatment of poliomyelitis by complete rest in bed, sulphonamide therapy, and isolation, and I would like to emphasize the necessity of giving the sulphonamides in massive doses in the early stage of the illness, especially in the more severe type of case, if subsequent paralysis is to be avoided. In such cases the drug should be given intravenously, due regard being paid to the fluid intake, alkalis, and the cytology of the blood. Initially I give for a young adult with a severe attack 4 g. of a soluble preparation of sulphonamide by the intravenous route, followed by 3 g. at intervals of twelve hours for two or three days.

In a series of over twenty cases of the severe type treated on these lines there has been either no paralysis supervening, or, where treatment was delayed and paralysis had already commenced, there has been no further extension of it.—I am, etc.,

Harrogate.

SINCLAIR MILLER.

Early Diagnosis of Tuberculous Meningitis

SIR,—Now that centres have been established for the treatment of military tuberculosis and tuberculous meningitis by streptomycin, a readjustment is necessary in our ideas about these forms of tuberculosis. Previously a diagnosis of tuberculous meningitis has been a sentence of death, only to be accepted when there is no longer any possibility of error. The possibility of cure has altered that, and early diagnosis has become a matter of particular importance, for it is not to be expected that patients will recover completely once damage to the brain has occurred. Therefore, it is necessary to study the early diagnosis of tuberculous meningitis. As the result of more than twenty years' experience in the treatment of children with tuberculosis I have formed the following opinions.

In many cases there is a prodromal period during which the disease is establishing itself and during which the signs and symptoms are very variable. The chief aids to diagnosis are clinical and neurological examination, examination of the cerebrospinal fluid, and radiography of the lungs. Any one of these may precede the others in providing evidence on which to base the diagnosis. Gross evidence, either clinical, laboratory, or radiological, means that the disease is becoming established, and brain damage may be already present. Fever of unknown origin and persistent headache in a child, especially one who is a tuberculosis contact or has some overt tuberculous disease, merit lumbar puncture and chest radiography. Even when neurological signs are absent, and when the cerebrospinal fluid is normal except possibly for a slight rise in the lymphocyte count, early military tuberculosis may be seen in the lungs; or, if the chest x-ray examination and neurological signs are negative, the cells in the cerebrospinal fluid may be definitely increased.

Early neurological signs may be equivocal and it is unwise to delay until they become well established. If there is present a combination of fever of unknown origin, persistent headache, and some definite evidence, however slight, from any of the triad, clinical, laboratory, or radiological, then streptomycin should be started. A specimen of the cerebrospinal fluid, injected before treatment into a guinea-pig, will probably reveal later whether the patient has been cured of tuberculous meningitis or an error in diagnosis has been made. In either case the patient's life will have been safeguarded to the best of the doctor's ability.—I am, etc.,

Braintree, Essex.

M. C. WILKINSON.