The predominance among males is well seen, the only specifically mentioned female cases being direct contacts of meningitis. (Mangiaracine and Pollen give no indication of the sex of five of their patients.) In nearly half the cases both eyes were involved, either simultaneously or at short intervals; corneal ulceration is mentioned in one-sixth of cases; in no case was the eye permanently damaged by the infection. Meningitis is an unusual but possible sequel.

Summary

Attention is drawn to the importance of recognizing the meningococcus as a primary cause of conjunctivitis.

Two cases in Leeds school-children are described, the first being treated eventually by local sulphacetamide and the second by local penicillin, with good results.

Some aspects of the literature are discussed.

We wish to thank Mr. John Foster, honorary ophthalmic surgeon to the Leeds General Infirmary, who was in charge of these cases, for his interest and encouragement and for his permission to publish this account. Our thanks are also due to Prof. J. W. McLeod, professor of bacteriology in the University of Leeds, to our colleagues in the department, and to Prof. A. Loewenstein for much helpful advice and criticism.

REFERENCES

- *Bennett, J. A. (1940). *J. R.A.M.C.*, 75, 264.
- *Clifton, F., and Laird, S. M. (1941). *Ibid.*, 77, 318.
- Cushing, R. W. (1940). *British Medical Journal*, 2, 439.
- Duke-Elder, Sir W. S. (1938). *Text Book of Ophthalmology*, 2, 1547, London.
- *Gifford, S. R., and Day, A. A. (1935). *Arch. Ophthalm.*, Chicago, 13, 1038.
- *Hayden, A. F., and Hayden, A. F. (1939). *Brit. J. vener. Dis.*, 15, 45.
- Herrell, W. E. (1945). *Penicillin and Other Antibiotic Agents*, p. 142, Philadelphia.
- *Mangiaracine, A. B., and Pollen, A. (1944). *Arch. Ophthalm.*, Chicago, 31, 284.
- *Reese, F. M. (1936). *Amer. J. Ophthalm.*, 19, 780.
- *Reid, R. D., and Bronstein, L. H. (1944). *J. Amer. med. Ass.*, 124, 703.
- *Smith, D. (1905). *Arch. Ophthalm.*, New York, 34, 481.
- *Theodore, F. H., and Kost, P. F. (1944). *Arch. Ophthalm.*, Chicago, 31, 245.
- *Thygeson, P. (1944). *Amer. J. Ophthalm.*, 27, 400.

Those marked with a * were used in the compilation of the Table.

THE BIOCHEMISTRY OF SENESCENCE*

BY

C. P. STEWART, M.Sc., Ph.D.

Reader in Clinical Chemistry, University of Edinburgh;
Senior Biochemist, Royal Infirmary, Edinburgh

Ray Lankester differentiated between "specific longevity" (the expectation of life at birth of a normal individual) and "potential longevity" (the duration of life which might be expected under an ideal environment). Knowledge of the processes involved in senescence, by which term we usually imply the generalized weakening of the vitality that accompanies growing old, may be expected to narrow the gap between specific and potential longevity. No doubt a gap must always remain, for even if we knew enough to provide the ideal physiological environment with complete protection from disease, and if we had the wit to apply our knowledge, there would still be occasional accidents.

The use of the term "potential longevity" implies a belief that the life of an organism is in fact limited and that senescence with ultimate death is its necessary fate. It is perfectly possible to argue that certain single-celled organisms are potentially immortal, since, given the

proper environment, they appear to be capable of reproducing themselves indefinitely. In one experiment extending over five years a single cell of paramecium was shown to be capable of producing 2^{3029} similar cells (equivalent to 10^{1000} times the volume of the earth), and the last cultures showed no signs of senility. But even if we admit that a complex organism such as man may be regarded as potentially immortal in the special sense that he can apparently produce children who in turn can reproduce their species, and so on *ad infinitum*, we have no reason for believing that the individual man can escape death. In studying senescence, therefore, we are not looking for the *elixir vitae*; we are seeking, from the point of view of abstract knowledge, to understand the changes produced by the passage of time; and by applying such knowledge we may hope to achieve a closer approximation between "specific" and "potential" longevity, with, in particular, an extension of the period of active and useful life. To believe in the possibility of such an approximation and to attempt its achievement is not to deny the proposition that heredity plays an important part in determining longevity. Whatever part heredity may play, be it great or small, the actual duration of life is the resultant of the effects of inheritance and environment. We may at least seek by knowledge and the application of knowledge to influence the latter.

Changes in Basal Metabolic Rate

There are many scattered references in the literature to quantitative changes associated with old age, in the composition of body fluids, in the estimated need for particular nutrients, and in the functional capacity both of individual organs and of the body as a whole. Perhaps the best-known of these is the change in the basal metabolic rate. The "prediction tables," produced by estimating the B.M.R. of considerable groups of subjects of different ages, indicate that the B.M.R. is practically constant during adult life to the age of 40, but thereafter falls each decade by approximately 1 calorie per sq. metre per hour. The differences between the age groups are statistically significant. Since this relationship between age and metabolic rate may be of fundamental importance it behoves us to examine it very carefully before accepting it at its face value. We may believe that the average B.M.R. of 100 men in the age group 60-70 is lower than that of a similar number of men aged 30-40. But, even though gross disease be excluded, a group of 100 men may be expected to include some with pathological changes not detectable by anything short of a most rigorous clinical examination—and perhaps not even then. The number is likely to be greater among the old than among the young. Is, then, the lower average B.M.R. of the older men due not to their greater age *per se* but to the greater incidence of pathological changes which may occur at any age but are commoner among the old? This is, of course, a difficulty of interpretation which must occur in any measurement comparing the old and young when the subjects are chosen only as "apparently healthy." It is obviously a serious difficulty and one which must be faced if we are to disentangle the changes inherent in the ageing process from those due to coincident disease. It is possible to get evidence on this point. In a small number of individuals the B.M.R. has been measured at frequent intervals over a considerable number of years. In two such cases the measurements extended roughly from age 40 to 60, during which period there were no more than inconsiderable changes in height and weight; in both the B.M.R. was significantly lower at 60 than at 40, though both individuals remained in good health. In a third case, over approximately the same period, the B.M.R. did not fall, and in this case the body weight was greater and the general

*Lecture to the Postgraduate Class of Internal Medicine, Edinburgh, June 10, 1947.

health was better at 60 than at 40 years of age. These results, though one would have preferred them from a greater number of cases, undoubtedly support the view that the B.M.R. is an expression of "vitality" and that in the absence of detectable disease it decreases appreciably in later life.

If, then, we accept the view that the observed decrease in the B.M.R. is an expression of the process of senescence, some at least of the other changes which have been reported may be related to it. For if the total energy production is decreased, a decrease in energy usage for certain of the individual processes must be postulated—there must be reduced activity in some of the directions in which energy is expended. Thus, although the secretion of gastric juice is not strictly a "basal" activity, the reduced cellular metabolic rate in old age may well be associated with the much greater incidence of hypochlorhydria (or even achlorhydria). It is stated that whereas achlorhydria occurs in about 4% of young adults it is found in some 30% of persons over 60—and one cannot imagine this phenomenon being accounted for by a differential death rate in favour of achlorhydrics.

Other Changes

Similarly, we might consider the lowered rate of energy production as contributing to the change in glucose tolerance which appears to accompany old age. The range of the response to the ingestion of 50 g. of glucose is very different for young adults and for old. The old usually show a blood-sugar maximum later (and higher) than the young, and the subsequent fall is slower, so that in many cases the fasting level (itself higher on the average than that of young people) is not attained within the usual period of two hours. It is by no means unusual for the blood-sugar curve to be such that in a young person it would be regarded as frankly diabetic—and this in the complete absence of any of the symptoms indicative of diabetes. Apart from the question obviously raised—What is diabetes?—these findings could be explained on the supposition that in old age both absorption of glucose from the alimentary tract and glycogen formation have become less efficient, and, since both these processes involve the expenditure of energy, they could be regarded as associated with the more sluggish metabolism expressed in the lowered B.M.R. It is true, however, that other interpretations of the tendency towards reduced glucose tolerance are possible, and we shall have occasion later to refer to at least one of these.

Again, decreased metabolic activity may be considered a causative agent in the decreased concentration of plasma proteins which, with the expected accompaniment of an increased blood sedimentation rate, has been found in old age.

Some of the changes which have been reported as accompanying old age are almost certainly not factors in the process of "growing old"—that is, in the essential involutionary processes—but are a statistical result of the more frequent occurrence among old people of diseases which may be found at any age. For example, it has been stated that the plasma non-protein nitrogen tends to rise in old age, that (as one would expect if the preceding statement were true) the blood urea is also raised, and that the urea clearance diminishes. Examination of the data shows, however, that there is no general progressive change of the blood urea or the urea clearance with increasing age; there is a tendency for the blood urea to be more often increased above the average for young people and for the urea clearance to be more often decreased. The "spread" becomes greater, suggesting strongly that the data merely express the increasing frequency of renal functional impairment.

Nevertheless, the very fact that renal failure and other "pathological" changes are much commoner in the aged than in the young indicates that, secondary though they may be to the fundamental changes involved in ageing, they may be closely bound up with those changes. Detailed examination of such functional disturbances and their age distribution might well yield valuable contributions to the solution of the major problem. The more information we can gain of the changes in old age, whether they form part of the essential process of ageing or whether they are secondary to these, the nearer we shall come to fitting together our jigsaw puzzle.

Nutritional Problems

In any comprehensive study of the ageing process nutritional problems must have an important part. We know a certain amount (though less than would sometimes appear) about the nutritional requirements during the period of growth. We know less about the requirements in later life. This is partly because growth itself is a phenomenon so obvious and easily measured that nutritional studies are much easier during that period than in adult life, when the criteria of good nutrition are difficult to define. Partly it is due to the impression that proper nutrition during the growth period presented a problem more urgent than that of nutrition in adult life, when, superficially at least, there seems to be greater tolerance of dietary variations. There are, however, some suggestions that more intensive investigation is required of the nutritional environment ideal for later life. Nor ought we to forget that senescence and growth are intimately related: a famous aphorism has it that senescence begins at birth.

Determination of the vitamin content of tissues has been possible for only a few years (and the results must still be accepted with caution), but there are indications of changes associated with age, although so far these have been demonstrated only by comparison of adult animals with young growing animals and with embryos. Extension of such investigations to old age is required, with further work on the influence of the nutritional environment on the tissue content of vitamins.

It has been shown in rats that a diet qualitatively as well balanced as possible but quantitatively insufficient for rapid growth promoted longevity and postponed the changes associated with old age. This, of course, is in keeping with the greater "expectation of life" of middle-aged people who are slightly below the average weight. Are we, then, wrong in providing (or attempting to provide) for our children a diet which not only contains the essential nutrients but is so generous in quantity that it induces rapid growth? A high-fat high-carbohydrate intake (which is in fact provided by any ordinary high-calorie diet) is by no means an unmixed blessing—it is no new idea that overfeeding may be a factor in the causation of diabetes, and various facts support the idea. Much evidence relates atherosclerosis to an excessive intake of fat and, especially, cholesterol; and there are data suggesting that the same factors may be concerned in the production of arteriosclerosis in man. Evidently research on the nutritional environment appropriate to maturity and old age may do much to diminish the incidence of serious metabolic disorders in older people, besides, in due course, contributing to our knowledge of the fundamental processes of growth and ageing. Elimination of disorders attributable to faulty nutrition would, apart altogether from incidental contributions to fundamental knowledge, undoubtedly reduce very considerably the gap between specific and potential longevity and so be of immense practical value.

Rate of Growth

A single-celled organism implanted in a suitable medium multiplies and so grows, the rate of growth continually increasing until a point is reached at which the supply of nutrients becomes insufficient to maintain the rate; thereafter growth slackens and then ceases, and the organism eventually dies from what is, in effect, starvation, or from the accumulation of the end-products of metabolism. Man also grows, and after a period during which growth has apparently ceased he dies. His cells have not lost the capacity for new growth—a wound heals in the aged as in the young, and in animals (rats) rapid growth has been induced at an age beyond the normal life-span by keeping the animals in a state of "arrested development" using a diet deficient in lysine and supplying the missing amino-acid only in extreme old age.

Now the complex and highly specialized organism man would appear to be in the position of the simple organism which is repeatedly transferred to a fresh medium (and so apparently maintained indefinitely in active growth), in that his supply of nutrients is continually renewed by his food and the excretory organs remove the end-products of his metabolism. If this were really so it would be necessary to attribute senescence and death to faulty environment. But the analogy between man and bacteria or fungi is a faulty one, for in the repeatedly subcultured colony we are really dealing with successive generations. It may be that, in man, changes in the intimate structure, which are natural in the sense that they form an essential part of the process of development, are harmful in that they ultimately affect adversely either the access of nutrient material to the cell (where alone metabolism occurs), and conversely the removal of waste products, or the enzyme systems which utilize this nutrient material.

Either of these hypotheses would account for many of the changes which have been reported as accompanying old age—the decreased basal metabolic rate, the tendency to hypochlorhydria, the decreased glucose tolerance, the increased requirements of certain of the vitamins which are concerned in enzymic activity, and so on. Equally, either would account for the increased incidence in old age of diseases which may be ascribed to metabolic disorders of particular organs—as, for example, hypertension may be the result of a failure of kidney metabolism.

Alteration in Cell Surfaces

Whether the process of senescence can ultimately be explained in either of these ways only much further research can tell. In the meantime there are a few "pointers" which indicate that slowly developing changes in the structure (and therefore the permeability) of the cell surfaces may be of fundamental importance. Recent work has suggested that although the cells, both red and white, are present in the same numbers in the blood of old people as of young, they are qualitatively altered in the aged. The red cells, though containing the "normal" amount of haemoglobin with the "normal" oxygen capacity, have, in old age, an altered stability towards hypotonic salt solutions which must be explained as due to alteration in the cell surfaces. These changes can be explained on the basis of a decreased rate of formation of the red cells (which fits the idea of a generalized decrease in cellular activity) and a corresponding decrease in the rate of cell destruction. This implies, in the red cell population at any given time, a greater number of "old" erythrocytes than in the blood of young people. The same suggestion of a lowered rate of cell formation and destruction is implicit in the finding that in the blood of aged people there is a tendency to an increased number of multi-lobular polymorphonuclear cells. This amounts, as has

been said, to no more than a "pointer," but it is one that is well worth following.

Were some change in the structure of cell surfaces demonstrable as an invariable concomitant of senescence, we should need much more knowledge than we now possess of this structure, and of the factors which influence it, before we could hope to delve further into the mysteries of senescence or to control it. We would have to be on our guard against ascribing too much to changes in the concentration of any one substance at the cell surfaces: any fundamental structural and functional changes there *may* be due to the alteration of a single constituent, but they may equally be of multiple origin. That structural changes do occur is suggested not only by the indirect evidence already outlined but by more direct evidence. Thus it has been demonstrated for certain plant leaves that, with increasing age of the leaf, electrical stimulation causes more calcium to leave the cell cortex and appear in the vacuole. (The conclusion was reached that with increasing age more calcium is concentrated at the cell surfaces, and the hypothesis was advanced that this, altering the permeability, favours the accumulation of toxic metabolites and so senescence.) In animal tissues, too, histological methods have been claimed to show increasing accumulation of calcium at the cell surfaces.

Conclusion

In this incomplete, and indeed sketchy, outline of some of the biochemical problems of senescence I have tried to indicate the kind of evidence which exists, the difficulties of interpreting it, and some of the directions which may profitably be taken by research. That the problems are complex is clear, but they are much more complex than would appear from this survey. For man, with his highly differentiated structure, is, in a sense, not one organism but many. His various tissues grow at different rates, in different ways, and, with different environmental demands, they age differently.

In research into the processes of ageing we must be clear as to our objects. One, of course, is the pursuit of knowledge for the sake of knowledge. Another is the application of such knowledge as we may gain. But our aim here is not merely to prolong life in terms of years; as has been wittily said: "It is for science not only to add years to life but, more important, to add life to years." There is the crux of our problem—so to *control* senescence that man's period of useful healthy life is prolonged, whether or not we succeed in increasing his total life-span. And I venture to suggest that such control will come, partly no doubt through study of the many degenerative diseases which more frequently affect the old, but chiefly and most completely through studies of the fundamental processes of cell metabolism and the effect of the nutritional environment on them. Among these latter studies an important place must be given to the problem of the structure and mechanism of the cell surfaces.

The Secretary of the Department of Health for Scotland, Sir George Henderson, speaking at the Annual Conference of the Scottish Association of Insurance Committees recently, said that there was no intention of setting up a huge bureaucratic machine to run hospitals as appendages of St. Andrew's House. The Secretary of State intended to leave the administration of the new hospital system in the hands of local people. New hospitals would be sited for maximum use of the widest area and not by accident of charity or the financial resources of a local authority. He hoped that outpatient departments would be so organized as to avoid the long queues which were common to-day. Under the National Health Service the work of the old insurance committees would be taken over by Executive Councils, which would represent the whole community and not simply the present insured population. By regrouping the areas covered the work of 54 insurance committees would be done by 25 Executive Councils.