divisions-extractions, by which the whole varicose vein could be extirpated through incisions—rarely more than 1/4 to 1/3 in. (0.63–0.85 cm.) in length—at intervals along the vein, the sites for incision carefully selected in the standing patient and indelibly marked before operation. The incisions, made parallel to the lines of tension of the skin, gape but little and do not require suturing. Only very few ligations of severed veins are ever necessary, and perforating veins and tributaries can be divided and ligated or, if small, torn off flush with the deep fascia without the use of a vein enucleator.

In some cases new varicose veins have appeared so soon after extirpation of the old as to raise inescapably the question of overloading in the deep veins and a new compensatory reflux through deep communicating veins, and the wisdom of severing all deep communications, if that were feasible.

Here I take opportunity to record my indebtedness to Mr. Park's painstaking anatomical studies (*Lond. Hosp. Gaz.*, 1945, 48, 221). He has demonstrated the number, complexity, and variability of the communications, deep and superficial, of the veins of the leg, and thus provided illuminating and irrefutable anatomical warning—a timely reinforcement of the conclusions many of us had come to from clinical studies—that it is not feasible on any occasion to deal with more than a very few of the deep communications.—I am, etc.,

London, E.16.

H. M. HANSCHALL.

SIR,—The article by Prof. A. M. Boyd and Mr. D. J. Robertson (Sept. 20, p. 452) with instructive phlebograms will I fear lead some to be less radical in their attacks upon varicose veins. It has always been accepted that solutions injected into varicose veins *must* inevitably enter the deep veins—and that very promptly, the patient noticing the taste of the solution within a few seconds of the injection. It has always been thought that the deep veins escaped damage because of the large volume of swift-flowing blood they contained diminishing internal damage, but most of all they were unaffected by sclerosing solutions because they were healthy normal veins.

I have encountered quite a number of Army pensioners with femoral thrombosis following ligatures and injections, but I have ascribed this complication to the long rest in bed (2–3 weeks) allowed after the operation of ligature or the occurrence of superficial thrombosis. Conditions in the Army apparently made it difficult to follow the ambulatory routine of civil life following ligatures and superficial thrombosis. I do not recall a single femoral thrombosis in a very large series of my own treated strictly on ambulatory lines. Others of the Army cases I have seen were due to operating on cases with bad thrombophilic histories, cases usually avoided for fear of stirring the hornets' nest.

Varicose veins are as hard to eradicate as rabbits, and if the article of Sept. 20 induces some to pull their punches I fear the pest of varicose veins will start to gain upon us again. For my part I will continue to inject downwards from the top ligature (if there is no valvular obstruction), upwards and downwards from the lower thigh ligature, downwards from the external saphenous ligature, and upwards from the excellent Dodds's ligature of the internal saphenous at the ankle. Up to a total of 100 ml. of 30% brine is used if the patient is anaesthetized, and less painful solutions if not. Following this all thrombosed veins are injected at weekly intervals. The foundation stone of success is to achieve a firm solid saphenous vein from groin to knee following the ligature.—I am, etc.,

London, W.1.

A. DICKSON WRIGHT.

Leprosy and its Problems

SIR,—In the leading article (June 7, p. 813) and in Dr. Bernard Moiser's letter (Aug. 30, p. 347) mention has been made of cockroaches as possible carriers of lepra bacilli. The idea that insect vectors play a role in transmission of the disease is evidently sound. But if an insect is to blame it could not be the cockroach, at least not exclusively. In this country we have had leprosy for centuries, but the disease is now practically extinct. It could not have been transmitted by cockroaches, because they did not exist in this country until a decade or two ago and are still practically unknown in the countryside, where leprosy has had its origin.

If an insect is a vector of this disease, lice or fleas are more likely to be instruments of transmission than any others. Leprosy is a disease of filth and disappears with cleanliness. It flourished in the Dark Ages, when the human body was neglected and when the old maxim of St. Jerome was still valid, that "*poenitens eo foetidior, eo pulcherrior*." The common belief that leprosy is transmitted by sexual intercourse might find its explanation in a transfer of phthirius inguinalis. In this connexion it may be mentioned that McCoy and Clegg found acid-fast bacilli which seemed to belong to the leprosy family in lice. The fact that savages are exempt might be explained by their nakedness and consequent inability to breed the body louse. When the savage begins to wear clothes and live in houses he will contract the disease, if the opportunity for infection is there. All this could fit in with the louse as vector.—I am, etc.,

Reykjavik.

NIELS DUNGAL.

Foetal Cries

SIR,—The prenatal cries of the Trowbridge foetus reported by Dr. E. Curphey (Sept. 27, p. 508) were anticipated by one at Strasbourg in 1821. A detailed account of the occurrence is given in a letter by Dr. William Lister to the *Edinburgh Medical and Surgical Journal* of October, 1822. There are remarkable points of similarity in the two cases.

The mother of the Strasbourg foetus had suffered repeated losses of liquor amnii up to the last month of her pregnancy. A sudden stumble, followed by considerable loss of liquor, occasioned unusually violent foetal movements and, to quote the author, "all at once the company present were alarmed by the cry of a child." The crying was repeated and "distinctly came from the womb of the patient. After a time it desisted." Two days later, in the eighth month of pregnancy, the mother went into labour and was delivered by cephalic presentation of a foetus of a premature infant. "The new born child," says the doctor, "gave a few weak tones" and died half an hour after birth. The author's comments on the aetiology of the event are worth repeating: "To judge from the evacuation of water, about the middle of the pregnancy, it may be well supposed that there had been some rent in the membranes by means of which the air was probably introduced. From the foregoing, it may then be well comprehended how the child cried; but how it could live 48 hours in the confined womb, surrounded with water in an amphibious manner is incomprehensible."

There may well have been other occasions between 1821 and 1947 when premature loss of amniotic fluid, air round the foetal face, and contraction of the uterus on the foetal chest have combined to produce cries loud enough to astonish the neighbours.—I am, etc.,

London, S.W.1.

W. HARTSTON.

The Extent of Neurosis

SIR,—I cannot let Dr. D. Yellowlees's letter (Sept. 20, p. 468) pass without comment. After reading the leading article on "Neurosis in Industry" (Aug. 16, p. 257) I had originally considered the results of my small survey in the form of a short article, but I did not think it worth the space and condensed the material into a letter (Sept. 6, p. 396). I agree that "more emotionally ill than actually so" is not altogether a very apt phrase, but it is not easy to define shortly the difference between an illness due to the invasion of external elements, or failure of internal mechanism, and regarded with no undue alarm by the patient, and the exaggeration of discomfort energized principally by an increase in already present strong basic anxiety, probably acting through endocrine channels and along pathways of outlet grooved from early childhood. By "actual" I meant "in terms of physical pathology" rather than and as different from a state of psychical activity.

I am in complete agreement with the fifteenth paragraph of Dr. John Rickman's very apposite article (Sept. 6, p. 363), which formulates the situation of the bromide-receiving woman perfectly, and Dr. Yellowlees will notice that I rate her as 100% ill. By "normal ailment" I meant one which is understandable and treatable in physical terms of microbial invasion and chemotherapy, and which is the normal sphere of the physician. To